

# True Decisions Inc.

An Independent Review Organization  
835 E. Lamar Blvd. #394  
Arlington, TX 76011  
Fax: 214-594-8680

Notice of Independent Review Decision

**DATE OF REVIEW:** DECEMBER 14, 2007

**IRO CASE #:**

**DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE**

Lumbar sacral (caudal) ESI #1 and Epidurography, radiological supervision.

**A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION**

Board Certified Orthopedic Surgeon

**REVIEW OUTCOME**

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

**INFORMATION PROVIDED TO THE IRO FOR REVIEW**

No ODG Guidelines  
Office note, Dr., 10/04/07  
Peer review, Dr., 10/29/07  
Peer review, Dr. , 11/09/07

**PATIENT CLINICAL HISTORY [SUMMARY]:**

This xx year-old male was injured on xx/xx/xx when he was bending over and a garage door fell on the back of his head. Dr. evaluated the claimant on 10/04/07 for cervical and low back pain. The claimant had moderate paravertebral muscle spasm, no tenderness and reduced lumbar flexion. Motor strength was intact and reflexes were symmetric. Dr. noted that a lumbar MRI was done on 10/28/05 that revealed an annular tear at L4-5. There was a broad based small protrusion at L4-5 and L5-S1 and a protrusion to the right at L4-5, L5-S1 contacting the L5-S1 nerve root. The impression was chronic cervicalgia and headaches with normal cervical MRI; chronic low back pain, intermittent

bilateral lower extremity pain and L4-5, L5-S1 herniated nucleus pulposus. Dr. recommended a lumbar epidural steroid injection.

**ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.**

This claimant is over two years post injury. His MRI was done two years ago and the report was not provided. There is no information regarding this claimant's treatment course to date or any indication that he has had any recent conservative treatment. It does not appear that there is pain in a particular dermatomal pattern or evidence of specific radiculopathy on exam. Given the nonspecific symptoms without a clear radicular pattern to the claimant's complaints or physical examination, the epidural steroid injection is not recommended in accordance with Official Disability Criteria.

Official Disability Guidelines Treatment in Worker's Comp 2007 Updates, Low Back: Epidural Steroid Injection.

Recommended as a possible option for short-term treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy) with use in conjunction with active rehab efforts. See specific criteria for use below. Radiculopathy symptoms are generally due to herniated nucleus pulposus or spinal stenosis, although ESIs have not been found to be as beneficial a treatment for the latter condition.

Short-term symptoms: The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months. (Armon, 2007) Epidural steroid injection can offer short-term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. There is little information on improved function or return to work. There is no high-level evidence to support the use of epidural injections of steroids, local anesthetics and/or opioids as a treatment for acute low back pain without radiculopathy. (Benzon, 1986) (ISIS, 1999) (DePalma, 2005) (Molloy, 2005) (Wilson-MacDonald, 2005)

Use for chronic pain: Chronic duration of symptoms (> 6 months) has also been found to decrease success rates with a threefold decrease found in patients with symptom duration > 24 months. The ideal time of either when to initiate treatment or when treatment is no longer thought to be effective has not been determined. (Hopwood, 1993) (Cyteval, 2006) Indications for repeating ESIs in patients with chronic pain at a level previously injected (> 24 months) include a symptom-free interval or indication of a new clinical presentation at the level.

Transforaminal approach: Some groups suggest that there may be a preference for a transforaminal approach as the technique allows for delivery of medication at the target tissue site, and an advantage for transforaminal injections in herniated nucleus pulposus over translaminar or caudal injections has been suggested in the best available studies. (Riew, 2000) (Vad, 2002) (Young, 2007) This approach may be particularly helpful in patients with large disc herniations, foraminal stenosis, and lateral disc herniations. (Colorado, 2001) (ICSI, 2004) (McLain, 2005) (Wilson-MacDonald, 2005)

Fluoroscopic guidance: Fluoroscopic guidance with use of contrast is recommended for all approaches as needle misplacement may be a cause of treatment failure. (Manchikanti, 1999) (Colorado, 2001) (ICSI, 2004) (Molloy, 2005) (Young, 2007)

Factors that decrease success: Decreased success rates have been found in patients who are unemployed due to pain, who smoke, have had previous back surgery, have pain that is not decreased by medication, and/or evidence of substance abuse, disability or litigation. (Jamison, 1991) (Abram, 1999) Research reporting effectiveness of ESIs in the past has been contradictory, but these discrepancies are felt to have been, in part, secondary to numerous methodological flaws in the early studies, including the lack of imaging and contrast administration. Success rates also may depend on the technical skill of the intervention list. (Carette, 1997) (Bigos, 1999) (Rozenberg, 1999) (Botwin, 2002) (Manchikanti, 2003) (CMS, 2004) (Delpont, 2004) (Khot, 2004) (Buttermann, 2004) (Buttermann2, 2004) (Samanta, 2004) (Cigna, 2004) (Benzon, 2005) (Dashfield, 2005) (Arden, 2005) (Price, 2005) (Resnick, 2005) (Abdi, 2007) (Boswell, 2007) Also see Epidural steroid injections, “series of three” and Epidural steroid injections, diagnostic. ESIs may be helpful with radicular symptoms not responsive to 2 to 6 weeks of conservative therapy. (Kinkade, 2007) As noted above, injections are recommended if they can facilitate a return to functionality (via activity & exercise). If post-injection physical therapy visits are required for instruction in these active self-performed exercise programs, these visits should be included within the overall recommendations under Physical therapy, or at least not require more than 2 additional visits to reinforce the home exercise program.

Criteria for the use of Epidural steroid injections:

Note: The purpose of ESI is to reduce pain and inflammation, restoring range of motion and thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit.

1. Radiculopathy must be documented. Objective findings on examination need to be present. For unequivocal evidence of radiculopathy, see AMA Guides, 5th Edition, page 382-383. (Andersson, 2000)
2. Initially unresponsive to conservative treatment (exercises, physical methods, NSAID's and muscle relaxants).
3. Injections should be performed using fluoroscopy (live x-ray) and injection of contrast for guidance.
4. At the time of initial use of an ESI (formally referred to as the “diagnostic phase” as initial injections indicate whether success will be obtained with this treatment intervention), a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. A second block is also not indicated if the first block is accurately placed unless: (a) there is a question of the pain generator; (b) there was possibility of inaccurate placement; or (c) there is evidence of multilevel pathology. In these cases a different level or approach might be proposed. There should be an interval of at least one to two weeks between injections. To be considered successful after this initial use of a block/blocks there should be documentation of at least 50-70% relief of pain from baseline and evidence of improved function for at least six to eight weeks after delivery.

5. No more than two nerve root levels should be injected using transforaminal blocks.
6. No more than one interlaminar level should be injected at one session.
7. In the therapeutic phase (the phase after the initial block/blocks were given and found to produce pain relief), repeat blocks should only be offered if there is at least 50-70% pain relief for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. (CMS, 2004) (Boswell, 2007)
8. Repeat injections should be based on continued objective documented pain relief, decreased need for pain medications, and functional response.
9. Current research does not support a routine use of a “series-of-three” injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections for the initial phase and rarely more than 2 for therapeutic treatment.
10. It is currently not recommended to perform epidural blocks on the same day of treatment as facet blocks or sacroiliac blocks or lumbar sympathetic blocks as this may lead to improper diagnosis or unnecessary treatment.
11. Cervical and lumbar epidural steroid injection should not be performed on the same day.

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:**

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES

- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES**
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR**
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS**
- TEXAS TACADA GUIDELINES**
- TMF SCREENING CRITERIA MANUAL**
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)**
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)**