

True Decisions Inc.

An Independent Review Organization
835 E. Lamar Blvd. #394
Arlington, TX 76011
Fax: 214-594-8680

Notice of Independent Review Decision

DATE OF REVIEW: January 16, 2008

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

Left Shoulder Arthroscopy and Rotator Cuff Repair

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION

Doctor of Medicine (M.D.)
Board Certified in Orthopaedic Surgery

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

INFORMATION PROVIDED TO THE IRO FOR REVIEW

Adverse Determination Letters, 11-1-07, 10-1-07
Medical records from Dr. 7-23-07 to 11-16-07
MRI shoulder 6-18-7

PATIENT CLINICAL HISTORY [SUMMARY]:

The patient suffered a work related injury to the left shoulder and right foot and ankle after a 6 foot fall. An MRI of the shoulder revealed a high riding humeral head and significant tearing and retraction of the entire rotator cuff.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

Dr. initial note describes the initial x-ray findings of severe arthritic changes at the AC joint and the glenohumeral joint. Therefore, this patient has a chronic rotator cuff tear and rotator cuff arthropathy. Rotator cuff repair is not indicated for this disease state. There are other options for the patient. Therefore, the Reviewer agrees with the non-authorization of surgery.

Surgery for rotator cuff repair	<p>Recommended as indicated below. Repair of the rotator cuff is indicated for significant tears that impair activities by causing weakness of arm elevation or rotation, particularly acutely in younger workers. However, rotator cuff tears are frequently partial-thickness or smaller full-thickness tears. For partial-thickness rotator cuff tears and small full-thickness tears presenting primarily as impingement, surgery is reserved for cases failing conservative therapy for three months. The preferred procedure is usually arthroscopic decompression, but the outcomes from open repair are as good or better. Surgery is not indicated for patients with mild symptoms or those who have no limitations of activities. (Ejnisman-Cochrane, 2004) (Grant, 2004) Lesions of the rotator cuff are best thought of as a continuum, from mild inflammation and degeneration to full avulsions. Studies of normal subjects document the universal presence of degenerative changes and conditions, including full avulsions without symptoms. Conservative treatment has results similar to surgical treatment but without surgical risks. Studies evaluating results of conservative treatment of full-thickness rotator cuff tears have shown an 82-86% success rate for patients presenting within three months of injury. The efficacy of arthroscopic decompression for full-thickness tears depends on the size of the tear; one study reported satisfactory results in 90% of patients with small tears. A prior study by the same group reported satisfactory results in 86% of patients who underwent open repair for larger tears. Surgical outcomes are much better in younger patients with a rotator cuff tear, than in older patients, who may be suffering from degenerative changes in the rotator cuff. Referral for surgical consultation may be indicated for patients who have: Activity limitation for more than three months, plus existence of a surgical lesion; Failure of exercise programs to increase range of motion and strength of the musculature around the shoulder, plus existence of a surgical lesion; Clear clinical and imaging evidence of a lesion that has been shown to benefit, in both the short and long term, from surgical repair; Red flag conditions (e.g., acute rotator cuff tear in a young worker, glenohumeral joint dislocation, etc.). Suspected acute tears of the rotator cuff in young workers may be surgically repaired acutely to restore function; in older workers, these tears are typically treated conservatively at first. Partial-thickness tears are treated the same as impingement syndrome regardless of MRI findings. Outpatient rotator cuff repair is a well accepted and cost effective procedure. (Cordasco, 2000) Difference between surgery & exercise was not significant. (Brox, 1999) There is significant variation in surgical decision-making and a lack of clinical agreement among orthopaedic surgeons about rotator cuff surgery. (Dunn, 2005)</p> <p><i>Revision rotator cuff repair:</i> The results of revision rotator cuff repair are inferior to those of primary repair. While pain relief may be achieved in most patients, selection criteria should include patients with an intact deltoid origin, good-quality rotator cuff tissue, preoperative elevation above the horizontal, and only one prior procedure. (Djurasovic, 2001)</p> <p><u>ODG Indications for Surgery™ -- Rotator cuff repair:</u> Criteria for rotator cuff repair with diagnosis of <u>full thickness</u> rotator cuff tear AND Cervical pathology and frozen shoulder syndrome have been ruled out: 1. Subjective Clinical Findings: Shoulder pain and inability to elevate the arm; tenderness over the greater tuberosity is common in acute cases. PLUS 2. Objective Clinical Findings: Patient may have weakness with abduction testing. May also demonstrate atrophy of shoulder musculature. Usually has full passive range of motion. PLUS</p>
---------------------------------	---

<p>3. Imaging Clinical Findings: Conventional x-rays, AP, and true lateral or axillary views. AND Gadolinium MRI, ultrasound, or arthrogram shows positive evidence of deficit in rotator cuff.</p> <p>Criteria for rotator cuff repair OR anterior acromioplasty with diagnosis of <u>partial thickness</u> rotator cuff repair OR acromial impingement syndrome (80% of these patients will get better without surgery.)</p> <p>1. Conservative Care: Recommend 3 to 6 months: Three months is adequate if treatment has been continuous, six months if treatment has been intermittent. Treatment must be directed toward gaining full ROM, which requires both stretching and strengthening to balance the musculature. PLUS</p> <p>2. Subjective Clinical Findings: Pain with active arc motion 90 to 130 degrees. AND Pain at night (Tenderness over the greater tuberosity is common in acute cases.) PLUS</p> <p>3. Objective Clinical Findings: Weak or absent abduction; may also demonstrate atrophy. AND Tenderness over rotator cuff or anterior acromial area. AND Positive impingement sign and temporary relief of pain with anesthetic injection (diagnostic injection test). PLUS</p> <p>4. Imaging Clinical Findings: Conventional x-rays, AP, and true lateral or axillary view. AND Gadolinium MRI, ultrasound, or arthrogram shows positive evidence of deficit in rotator cuff.</p> <p>(Washington, 2002)</p>
--

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR**
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS**
- TEXAS TACADA GUIDELINES**
- TMF SCREENING CRITERIA MANUAL**
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE**
 - **THE SHOULDER, ROCKWOOD & MATSEN**
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)**