

I-Decisions Inc.

An Independent Review Organization

71 Court Street

Belfast, Maine 04915

(207) 338-1141 (phone)

(866) 676-7547 (fax)

DATE OF REVIEW: OCTOBER 16, 2007

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

Anterior Cervical Discectomy and Fusion C4-5, C5-6, C6-7

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION

MD, Board Certified in Orthopedic Surgery

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

Provide a description of the review outcome that clearly states whether or not medical necessity exists for each of the health care services in dispute.

INFORMATION PROVIDED TO THE IRO FOR REVIEW

1. Case assignment with UR denials (9-17-07 and 8-24-07)
2. Letters from, MD 9-24-07, 5-22-07, 7-10-07, 8-14-07
3. Pain management notes 12-29-06,
4. Operative report cervical facet medial branch blocks 12-13-06
5. C-spine MRI report 4-24-06
6. FCE 3-27-07
7. Notes 3-30-07, 3-28-07, 3-23-07, 3-8-07
8. CT Myelogram report 4-17-07
9. ODG Fusion guidelines
10. Institute Behavioral Medicine Pre-Surgical Screening 7-26-07

11. *Discogram report 6-27-07*

PATIENT CLINICAL HISTORY [SUMMARY]:

The injured employee tripped over a cable at work and injured her neck. She developed significant, disabling neck pain due to disc dysfunction. She failed conservative care, including therapy, chiropractic, medications, and injections. A multilevel arthroplasty was recommended, but denied by the insurance company. The patient requested fusion for her pain after a cervical discogram showed concordant pain response at the requested levels.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

This request does not meet ODG criteria for cervical fusion. ODG criteria does not address fusion at more than two levels. To quote Dr. on July 10, 2007, "In my opinion, a three-level cervical fusion would not be in her best interest, it would leave her with restricted range of motion and the significant risk of adjacent level breakdown in a relatively short period of time". The denial for multi-level fusion should be upheld based on the requesting physician's initial opinion and the ODG guidelines.

Fusion, anterior cervical	<p>Recommended as an option in combination with anterior cervical discectomy for approved indications, although current evidence is conflicting about the benefit of fusion in general. (See Discectomy/laminectomy/laminoplasty.) Evidence is also conflicting as to whether autograft or allograft is preferable and/or what specific benefits are provided with fixation devices. Many patients have been found to have excellent outcomes while undergoing simple discectomy alone (for one- to two-level procedures), and have also been found to go on to develop spontaneous fusion after an anterior discectomy. (Bertalanffy, 1988) (Savolainen, 1998) (Donaldson, 2002) (Rosenorn, 1983) Cervical fusion for degenerative disease resulting in axial neck pain and no radiculopathy remains controversial and conservative therapy remains the choice if there is no evidence of instability. (Bambakidis, 2005) Conservative anterior cervical fusion techniques appear to be equally effective compared to techniques using allografts, plates or cages. (Savolainen, 1998) (Dowd, 1999) (Colorado, 2001) (Fouyas-Cochrane, 2002) (Goffin, 2003) This evidence was substantiated in a recent Cochrane review that stated that hard evidence for the need for a fusion procedure after discectomy was lacking, as outlined below:</p> <p>(1) <i>Anterior cervical discectomy compared to anterior cervical discectomy with interbody fusion with a bone graft or substitute:</i> Three of the six randomized controlled studies discussed in the 2004 Cochrane review found no difference between the two techniques and/or that fusion was not necessary. The Cochrane review felt there was conflicting evidence of the relative effectiveness of either procedure. Overall it was noted that patients with discectomy only had shorter hospital stays, and shorter length of operation. There was moderate evidence that pain relief after five to six weeks was higher for the patients who had discectomy with fusion. Return to work was higher early on (five weeks) in the patients with discectomy with fusion, but there was no significant difference at ten weeks. (Jacobs-Cochrane, 2004) (Abd-Alrahman, 1999) (Dowd, 1999) (Martins, 1976) (van den Bent, 1996) (Savolainen, 1998) One disadvantage of fusion appears to be abnormal kinematic strain on adjacent spinal levels. (Ragab, 2006) (Eck, 2002) (Matsunaga, 1999) (Katsuura, 2001) The advantage of fusion appears to be a decreased rate of kyphosis in the operated segments. (Yamamoto, 1991) (Abd-Alrahman, 1999)</p> <p>(2) <i>Fusion with autograft versus allograft:</i> The Cochrane review found limited evidence that</p>
---------------------------	---

the use of autograft provided better pain reduction than animal allograft. It also found that there was no difference between biocompatible osteoconductive polymer or autograft (limited evidence). ([Jacobs-Cochrane, 2004](#)) ([McConnell, 2003](#)) A problem with autograft is morbidity as related to the donor site including infection, prolonged drainage, hematomas, persistent pain and sensory loss. ([Younger, 1989](#)) ([Sawin, 1998](#)) ([Sasso, 2005](#)) Autograft is thought to increase fusion rates with less graft collapse. ([Deutsch, 2007](#)). See [Decompression, myelopathy](#).

(3) *Fusion with autograft with plate fixation versus allograft with plate fixation, Single level:* A recent retrospective review of patients who received allograft with plate fixation versus autograft with plate fixation at a single level found fusion rates in 100% versus 90.3% respectively. This was not statistically significant. Satisfactory outcomes were noted in all non-union patients. ([Samartzis, 2005](#))

(4) *Fusion with different types of autograft:* The Cochrane review did not find evidence that a vertebral body graft was superior to an iliac crest graft. ([McGuire, 1994](#))

(5) *Fusion with autograft versus fusion with autograft and additional instrumentation: Plate Fixation:* In single-level surgery there is limited evidence that there is any difference between the use of plates and fusion with autograft in terms of union rates. For two-level surgery, there was moderate evidence that there was more improvement in arm pain for patients treated with a plate than for those without a plate. Fusion rate is improved with plating in multi-level surgery. ([Wright, 2007](#)) See [Plate fixation, cervical spine surgery](#).

Cage: Donor site pain may be decreased with the use of a cage rather than a plate, but donor site pain was not presented in a standardized manner. At two years pseudoarthrosis rate has been found to be lower in the fusion group (15%) versus the cage group (44%). A six-year follow-up of the same study group revealed no significant difference in outcome variables between the two treatment groups (both groups had pain relief). In the subgroup of patients with the cage who attained fusion, the overall outcome was better than with fusion alone. Patients treated with cage instrumentation have less segmental kyphosis and better-preserved disc height. This only appears to affect outcome in a positive way in cage patients that achieve fusion (versus cage patients with pseudoarthrosis). ([Poelsson, 2007](#)) ([Varuch, 2002](#)) ([Hacker 2000](#)) See also [Adjacent segment disease/degeneration](#) (fusion).

(6) *Fusion with allograft alone versus with allograft and additional instrumentation:*

Plate Fixation: Retrospective studies indicate high levels of pseudoarthrosis rates (as high as 20% for one-level and 50% for two-level procedures) using allograft alone. In a recent comparative retrospective study examining fusion rate with plating, successful fusion was achieved in 96% of single-level cases and 91% of two-level procedures. This could be compared to a previous retrospective study by the same authors of non-plated cases that achieved successful fusion in 90% of single-level procedures and 72% of two-level procedures. ([Kaiser, 2002](#)) ([Martin, 1999](#)) See [Plate fixation, cervical spine surgery](#).

Complications:

Collapse of the grafted bone and loss of cervical lordosis: collapse of grafted bone has been found to be less likely in plated groups for patients with multiple-level fusion. Plating has been found to maintain cervical lordosis in both multi-level and one-level procedures. ([Trojanovich, 2002](#)) ([Herrmann, 2004](#)) ([Katsuura, 1996](#)) The significance on outcome of kyphosis or loss of cervical lordosis in terms of prediction of clinical outcome remains under investigation. ([Peolsson, 2004](#)) ([Haden, 2005](#)) ([Poelsson, 2007](#)) ([Hwang, 2007](#))

Pseudoarthrosis: This is recognized as an etiology of continued cervical pain and unsatisfactory outcome. Treatment options include a revision anterior approach vs. a posterior approach. Regardless of approach, there is a high rate of continued moderate to severe pain even after solid fusion is achieved. ([Kuhns, 2005](#)) ([Mummaneni, 2004](#)) ([Coric, 1997](#))

Anterior versus posterior fusion: In a study based on 932,009 hospital discharges associated with cervical spine surgery, anterior fusions were shown to have a much lower rate of complications compared to posterior fusions, with the overall percent of cases with complications being 2.40% for anterior decompression, 3.44% for anterior fusion, and 10.49% for posterior fusion. ([Wang, 2007](#)) *Predictors of outcome of ACDF:* Predictors of good outcome include non-smoking, a pre-operative lower pain level, soft disc disease, disease in one level, greater segmental kyphosis pre-operatively, radicular pain without additional neck or lumbar pain, short duration of symptoms, younger age, no use of analgesics, and normal ratings on biopsychosocial tests such as the Distress and Risk

Assessment Method (DRAM). Predictors of poor outcomes include non-specific neck pain, psychological distress, psychosomatic problems and poor general health. (Peolsson, 2006) (Peolsson, 2003) See Plate fixation, cervical spine surgery .
--

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)