

I-Decisions Inc.

An Independent Review Organization

71 Court Street

Belfast, Maine 04915

(207) 338-1141 (phone)

(866) 676-7547 (fax)

DATE OF REVIEW: OCTOBER 23, 2007

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

Posterior fusion, L5-S1 (22612, 63047, 22842, 20931) with 2-day inpatient stay

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION

Doctor of Medicine (M.D.)

Board Certified in Orthopaedic Surgery

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

Provide a description of the review outcome that clearly states whether or not medical necessity exists for each of the health care services in dispute.

The reviewer finds medical necessity exists for this patient for the following services: Posterior fusion, L5-S1 (22612, 63047, 22842, 20931) with 2-day inpatient stay.

INFORMATION PROVIDED TO THE IRO FOR REVIEW

Adverse Determination Letters, 8/29/07, 9/18/07

Letters, 8/28/07, 9/18/07

Official Disability Guidelines and Treatment Guidelines

Institute, Dr., 8/23/07, 7/6/07, 5/25/07, 3/27/07, 2/7/07, 3/10/06, 2/16/06, 1/6/06,

11/29/05, 7/27/04, 3/2/04, 2/23/04, 11/24/03, 10/7/03, 8/23/03, 8/4/03, 6/23/03,
 6/17/03, 6/3/03, 5/14/03,
 5/9/03, 4/2/03, 3/17/03
 Operative Reports, 3/8/07, 8/11/03, 2/23/06, 12/22/05
 MRI, 3/31/03

PATIENT CLINICAL HISTORY [SUMMARY]:

The injured employee complains of continued back pain s/p multiple lumbar surgeries for L4-S1 fusion and reoperation for pseudarthrosis. He has multiple injections and rhizotomies. Posterior fusion of the L5-S1 facets has been recommended. He has had excellent, but temporary relief from posterior facet injections / rhizotomies at L5/S1.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

The reviewer finds that the patient’s UR denials have been inappropriate. This patient is a candidate for surgery based on ODG criteria. Psychological screening has been appropriate and the patient has failed adequate conservative care.

The patient fits the following ODG criteria, *“Primary Mechanical Back Pain (i.e., pain aggravated by physical activity)/Functional Spinal Unit Failure/Instability, including one or two level segmental failure with progressive degenerative changes, loss of height, disc loading capability.”*

The reviewer finds medical necessity exists for this patient for the following services: Posterior fusion, L5-S1 (22612, 63047, 22842, 20931) with 2-day inpatient stay.

Fusion (spinal)	<p>Not recommended for patients who have less than six months of failed recommended conservative care unless there is objectively demonstrated severe structural instability and/or acute or progressive neurologic dysfunction, but recommended as an option for spinal fracture, dislocation, spondylolisthesis or frank neurogenic compromise, subject to the selection criteria outlined in the section below entitled, “Patient Selection Criteria for Lumbar Spinal Fusion,” after 6 months of conservative care. After screening for psychosocial variables, outcomes are improved and fusion may be recommended for degenerative disc disease with spinal segment collapse with or without neurologic compromise after 6 months of recommended conservative therapy. For complete references, see separate document with all studies focusing on Fusion (spinal). There is limited scientific evidence about the long-term effectiveness of fusion for degenerative disc disease compared with natural history, placebo, or conservative treatment. Studies conducted in order to compare different surgical techniques have shown success for fusion in carefully selected patients. (Gibson-Cochrane, 2000) (Savolainen, 1998) (Wetzel, 2001) (Molinari, 2001) (Bigos, 1999) (Washington, 1995) (DeBarard-Spine, 2001) (Fritzell-Spine, 2001) (Fritzell-Spine, 2002) (Deyo-NEJM, 2004) (Gibson-Cochrane/Spine, 2005) (Soegaard, 2005) (Glassman, 2006)</p>
-----------------	--

(Atlas, 2006) According to the recently released AANS/NASS Guidelines, lumbar fusion is recommended as a treatment for carefully selected patients with disabling low back pain due to one- or two-level degenerative disc disease after failure of an appropriate period of conservative care. This recommendation was based on one study that contained numerous flaws, including a lack of standardization of conservative care in the control group. At the time of the 2-year follow up it appeared that pain had significantly increased in the surgical group from year 1 to 2. Follow-up post study is still pending publication. In addition, there remains no direction regarding how to define the “carefully selected patient.”

(Resnick, 2005) (Fritzell, 2004) A recently published well respected international guideline, the “European Guidelines,” concluded that fusion surgery for nonspecific chronic LBP cannot be recommended unless 2 years of all other recommended conservative treatments – including multidisciplinary approaches with combined programs of cognitive intervention and exercises – have failed, or such combined programs are not available, and only then in carefully selected patients with maximum 2-level degenerative disc disease. (Airaksinen, 2006) For chronic LBP, exercise and cognitive intervention may be equivalent to lumbar fusion without the potentially high surgical complication rates. (Ivar Brox-Spine, 2003) (Keller-Spine, 2004) (Fairbank-BMJ, 2005) (Brox, 2006) In acute spinal cord injury (SCI), if the spine is unstable following injury, surgical fusion and bracing may be necessary. (Bagnall-Cochrane, 2004) (Siebenga, 2006) A study on improving quality through identifying inappropriate care found that use of guideline-based Utilization Review (UR) protocols resulted in a denial rate for lumbar fusion 59 times as high as denial rates using non-guideline based UR. (Wickizer, 2004) The profit motive and market medicine have had a significant impact on clinical practice and research in the field of spine surgery. (Weiner-Spine, 2004) (Shah-Spine, 2005) (Abelson, 2006) Data on geographic variations in medical procedure rates suggest that there is significant variability in spine fusion rates, which may be interpreted to suggest a poor professional consensus on the appropriate indications for performing spinal fusion. (Deyo-Spine, 2005) (Weinstein, 2006) Outcomes from complicated surgical fusion techniques (with internal fixation) may be no better than the traditional posterolateral fusion. (van Tulder, 2006) (Maghout-Juratli, 2006) Despite the new technologies, reoperation rates after lumbar fusion have become higher. (Martin, 2007) According to the recent Medicare Coverage Advisory Committee Technology Assessment, the evidence for lumbar spinal fusion does not conclusively demonstrate short-term or long-term benefits compared with nonsurgical treatment for elderly patients. (CMS, 2006) When lumbar fusion surgery is performed, either with lateral fusion alone or with interbody fusion, unlike cervical fusion, there is no absolute contraindication to patients returning even to contact sports after complete recovery from surgery. Like patients with a thoracic injury, those with a lumbar injury should be pain free, have no disabling neurological deficit, and exhibit evidence of bone fusion on x-ray films before returning. (Burnett, 2006) A recent randomized controlled trial comparing decompression with decompression and instrumented fusion in patients with foraminal stenosis and single-level degenerative disease found that patients universally improved with surgery, and this improvement was maintained at 5 years. However, no obvious additional benefit was noted by combining decompression with an instrumented fusion. (Hallett, 2007) Lumbar spinal fusion surgeries use bone grafts, and are sometimes combined with metal devices, to produce a rigid connection between two or more adjacent vertebrae. The therapeutic objective of spinal fusion surgery for patients with low back problems is to prevent any movement in the intervertebral spaces between the fused vertebrae, thereby reducing

pain and any neurological deficits.

Lumbar fusion in workers' comp patients: In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. Until further research is conducted there remains insufficient evidence to recommend fusion for chronic low back pain in the absence of stenosis and spondylolisthesis, and this treatment for this condition remains "under study." It appears that workers' compensation populations require particular scrutiny when being considered for fusion for chronic low back pain, as there is evidence of poorer outcomes in subgroups of patients who were receiving compensation or involved in litigation. ([Fritzell-Spine, 2001](#)) ([Harris-JAMA, 2005](#)) ([Maghout-Juratli, 2006](#)) ([Atlas, 2006](#)) Despite poorer outcomes in workers' compensation patients, utilization is much higher in this population than in group health. ([Texas, 2001](#)) ([NCCI, 2006](#)) Presurgical biopsychosocial variables predict patient outcomes from lumbar fusion, which may help improve patient selection. Workers' compensation status, smoking, depression, and litigation were the most consistent presurgical predictors of poorer patient outcomes. Other predictors of poor results were number of prior low back operations, low household income, and older age. ([DeBerard-Spine, 2001](#)) ([DeBerard, 2003](#)) ([Deyo, 2005](#)) ([LaCaille, 2005](#)) ([Trief-Spine, 2006](#)) Obesity and litigation in workers' compensation cases predict high costs associated with interbody cage lumbar fusion. ([LaCaille, 2007](#))

Lumbar fusion for spondylolisthesis: Recommended as an option for spondylolisthesis. Patients with increased instability of the spine after surgical decompression at the level of degenerative spondylolisthesis are candidates for fusion. ([Eckman, 2005](#)) This study found only a 27% success from spinal fusion in patients with low back pain and a positive single-level low-pressure provocative discogram, versus a 72% success in patients having a well-accepted single-level lumbar pathology of unstable spondylolisthesis. ([Carragee, 2006](#)) Unilateral instrumentation used for the treatment of degenerative lumbar spondylolisthesis is as effective as bilateral instrumentation. ([Fernandez-Fairen, 2007](#)) Patients with degenerative spondylolisthesis and spinal stenosis who undergo standard decompressive laminectomy (with or without fusion) showed substantially greater improvement in pain and function during a period of 2 years than patients treated nonsurgically, according to the recent results from the Spine Patient Outcomes Research Trial (SPORT). ([Weinstein-spondylolisthesis, 2007](#)) ([Deyo-NEJM, 2007](#)) For degenerative lumbar spondylolisthesis, spinal fusion may lead to a better clinical outcome than decompression alone. No conclusion about the clinical benefit of instrumenting a spinal fusion can be made, but there is moderate evidence that the use of instrumentation improves the chance of achieving solid fusion. ([Martin, 2007](#))

Patient Selection Criteria for Lumbar Spinal Fusion:

For chronic low back problems, fusion should not be considered within the first 6 months of symptoms, except for fracture, dislocation or progressive neurologic loss. Indications for spinal fusion may include: (1) Neural Arch Defect - Spondylolytic spondylolisthesis, congenital unilateral neural arch hypoplasia. (2) Segmental Instability (objectively demonstrable) - Excessive motion, as in degenerative spondylolisthesis, surgically induced segmental instability and mechanical intervertebral collapse of the motion segment and advanced degenerative changes after surgical disectomy. For spinal instability criteria, see AMA Guides, 5th Edition, page 379 (lumbar inter-segmental movement of more than 4.5 mm). ([Andersson, 2000](#)) (3) Primary Mechanical Back Pain (i.e., pain aggravated by physical activity)/Functional Spinal Unit Failure/Instability, including one or two

	<p>level segmental failure with progressive degenerative changes, loss of height, disc loading capability. In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. (4) Revision Surgery for failed previous operation(s) if significant functional gains are anticipated. Revision surgery for purposes of pain relief must be approached with extreme caution due to the less than 50% success rate reported in medical literature. (5) Infection, Tumor, or Deformity of the lumbosacral spine that cause intractable pain, neurological deficit and/or functional disability.</p> <p>Pre-Operative Surgical Indications Recommended: Pre-operative clinical surgical indications for spinal fusion should include all of the following: (1) All pain generators are identified and treated; & (2) All physical medicine and manual therapy interventions are completed; & (3) X-rays demonstrating spinal instability and/or myelogram, CT-myelogram, or discography (see discography criteria) & MRI demonstrating disc pathology; & (4) Spine pathology limited to two levels; & (5) Psychosocial screen with confounding issues addressed. (6) For any potential fusion surgery, it is recommended that the injured worker refrain from smoking for at least six weeks prior to surgery and during the period of fusion healing. (Colorado, 2001) (BlueCross BlueShield, 2002)</p>
--	--

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)