

C-IRO, Inc.
An Independent Review Organization
7301 Ranch Rd. 620 N, Suite 155-199
Austin, TX 78726

Notice of Independent Review Decision

DATE OF REVIEW: AUGUST 29, 2008

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

Medical necessity of C6-7 ACDF anterior cervical fusion instrumentation and removal C5-6 instrumentation with three day length of stay, 63075, 22554, 22845, 20931, 22851, 20936, 22855.

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION

MD, Board Certified Orthopedic Surgeon

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

Provide a description of the review outcome that clearly states whether or not medical necessity exists for each of the health care services in dispute.

The reviewer finds that medical necessity exists for C6-7 ACDF anterior cervical fusion instrumentation and removal C5-6 instrumentation with three day length of stay, 63075, 22554, 22845, 20931, 22851, 20936, 22855.

INFORMATION PROVIDED TO THE IRO FOR REVIEW

MRI cervical spine, 12/27/07
Office notes, Dr. , 04/01/08
Letters, Dr. , 04/09/08, 05/02/08, 07/07/08, 07/10/08, 07/16/08, 07/31/08, 08/12/08
Peer review, I, 07/03/08, 07/31/08
Peer review, undated
ODG Guidelines and Treatment Guidelines

PATIENT CLINICAL HISTORY [SUMMARY]:

This is a xx year-old female. Dr. has been following the claimant. The MRI of the cervical spine dated 12/27/07 showed an extruded disc posteriorly and to the left at the C6-7 level causing some impingement upon the neural foramen on the left at this level as well as some impingement upon the cervical cord. A previous anterior cervical fusion with a plate and screws at the C5-6 vertebral levels was reported. Dr. felt that the claimant had weakness to the left triceps and absent triceps reflexes bilaterally.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION

Evidence based Official Disability Guidelines suggest that cervical discectomy and fusion is indicated for individuals who have failed traditional forms of conservative treatment, have symptoms, exam findings, and imaging studies all of which are consistent. In this particular case, the claimant has previously undergone a fusion at C5-6, but reportedly has imaging studies which show a herniated disc at C6-7 consistent with her subjective complaints of left arm pain and triceps weakness. Records reflect months of conservative care in the form of physical therapy, activity modifications and medical management.

The information provided does satisfy the traditional criteria established within the evidence based literature, and as such the request for anterior cervical discectomy and fusion in this particular case would appear to be reasonable and medically necessary.

The reviewer finds that medical necessity exists for C6-7 ACDF anterior cervical fusion instrumentation and removal C5-6 instrumentation with three day length of stay, 63075, 22554, 22845, 20931, 22851, 20936, 22855.

Official Disability Guidelines Treatment in Workers' Comp 2008 Updates, neck and upper back-fusion anterior

Recommended as an option in combination with anterior cervical discectomy for approved indications, although current evidence is conflicting about the benefit of fusion in general. (See Discectomy/laminectomy/laminoplasty.) Evidence is also conflicting as to whether autograft or allograft is preferable and/or what specific benefits are provided with fixation devices. Many patients have been found to have excellent outcomes while undergoing simple discectomy alone (for one- to two-level procedures), and have also been found to go on to develop spontaneous fusion after an anterior discectomy. (Bertalanffy, 1988) (Savolainen, 1998) (Donaldson, 2002) (Rosenorn, 1983) Cervical fusion for degenerative disease resulting in axial neck pain and no radiculopathy remains controversial and conservative therapy remains the choice if there is no evidence of instability. (Bambakidis, 2005) Conservative anterior cervical fusion techniques appear to be equally effective compared to techniques using allografts, plates or cages. (Savolainen, 1998) (Dowd, 1999) (Colorado, 2001) (Fouyas-Cochrane, 2002) (Goffin, 2003) Cervical fusion may demonstrate good results in appropriately chosen patients with cervical spondylosis and axial neck pain. (Wieser, 2007) This evidence was substantiated in a recent Cochrane review that stated that hard evidence for the need for a fusion procedure after discectomy was lacking, as outlined below:

(1) *Anterior cervical discectomy compared to anterior cervical discectomy with interbody fusion with a bone graft or substitute:* Three of the six randomized controlled studies discussed in the 2004 Cochrane review found no difference between the two techniques and/or that fusion was not necessary. The Cochrane review felt there was conflicting evidence of the relative effectiveness of either procedure. Overall it was noted that patients with discectomy only had shorter hospital stays, and shorter length of operation. There was moderate evidence that pain relief after five to six weeks was higher for the patients who had discectomy with fusion. Return to work was higher early on (five weeks) in the patients

with discectomy with fusion, but there was no significant difference at ten weeks. (Jacobs-Cochrane, 2004) (Abd-Alrahman, 1999) (Dowd, 1999) (Martins, 1976) (van den Bent, 1996) (Savolainen, 1998) One disadvantage of fusion appears to be abnormal kinematic strain on adjacent spinal levels. (Ragab, 2006) (Eck, 2002) (Matsunaga, 1999) (Katsuura, 2001) The advantage of fusion appears to be a decreased rate of kyphosis in the operated segments. (Yamamoto, 1991) (Abd-Alrahman, 1999)

(2) *Fusion with autograft versus allograft:* The Cochrane review found limited evidence that the use of autograft provided better pain reduction than animal allograft. It also found that there was no difference between biocompatible osteoconductive polymer or autograft (limited evidence). (Jacobs-Cochrane, 2004) (McConnell, 2003) A problem with autograft is morbidity as related to the donor site including infection, prolonged drainage, hematomas, persistent pain and sensory loss. (Younger, 1989) (Sawin, 1998) (Sasso, 2005) Autograft is thought to increase fusion rates with less graft collapse. (Deutsch, 2007). See Decompression, myelopathy.

(3) *Fusion with autograft with plate fixation versus allograft with plate fixation, Single level:* A recent retrospective review of patients who received allograft with plate fixation versus autograft with plate fixation at a single level found fusion rates in 100% versus 90.3% respectively. This was not statistically significant. Satisfactory outcomes were noted in all non-union patients. (Samartzis, 2005)

(4) *Fusion with different types of autograft:* The Cochrane review did not find evidence that a vertebral body graft was superior to an iliac crest graft. (McGuire, 1994)

(5) *Fusion with autograft versus fusion with autograft and additional instrumentation:*

Plate Fixation: In single-level surgery there is limited evidence that there is any difference between the use of plates and fusion with autograft in terms of union rates. For two-level surgery, there was moderate evidence that there was more improvement in arm pain for patients treated with a plate than for those without a plate. Fusion rate is improved with plating in multi-level surgery. (Wright, 2007) See Plate fixation, cervical spine surgery.

Cage: Donor site pain may be decreased with the use of a cage rather than a plate, but donor site pain was not presented in a standardized manner. At two years pseudoarthrosis rate has been found to be lower in the fusion group (15%) versus the cage group (44%). A six-year follow-up of the same study group revealed no significant difference in outcome variables between the two treatment groups (both groups had pain relief). In the subgroup of patients with the cage who attained fusion, the overall outcome was better than with fusion alone. Patients treated with cage instrumentation have less segmental kyphosis and better-preserved disc height. This only appears to affect outcome in a positive way in cage patients that achieve fusion (versus cage patients with pseudoarthrosis). (Poelsson, 2007) (Varuch, 2002) (Hacker 2000) See also Adjacent segment disease/degeneration (fusion).

(6) *Fusion with allograft alone versus with allograft and additional instrumentation:*

Plate Fixation: Retrospective studies indicate high levels of pseudoarthrosis rates (as high as 20% for one-level and 50% for two-level procedures) using allograft alone. In a recent comparative retrospective study examining fusion rate with plating, successful fusion was achieved in 96% of single-level cases and 91% of two-level procedures. This could be compared to a previous retrospective study by the same authors of non-plated cases that achieved successful fusion in 90% of single-level procedures and 72% of two-level procedures. (Kaiser, 2002) (Martin, 1999) See Plate fixation, cervical spine surgery.

Complications:

Collapse of the grafted bone and loss of cervical lordosis: collapse of grafted bone has been found to be less likely in plated groups for patients with multiple-level fusion. Plating has been found to maintain cervical lordosis in both multi-level and one-level procedures. (Troyanovich, 2002) (Herrmann, 2004) (Katsuura, 1996) The significance on outcome of kyphosis or loss of cervical lordosis in terms of prediction of clinical outcome remains under investigation. (Peolsson, 2004) (Haden, 2005) (Poelsson, 2007) (Hwang, 2007)

Pseudoarthrosis: This is recognized as an etiology of continued cervical pain and unsatisfactory outcome. Treatment options include a revision anterior approach vs. a posterior approach. Regardless of approach, there is a high rate of continued moderate to severe pain even after solid fusion is achieved. (Kuhns, 2005) (Mummaneni, 2004) (Coric, 1997)

Anterior versus posterior fusion: In a study based on 932,009 hospital discharges associated with cervical spine surgery, anterior fusions were shown to have a much lower rate of complications compared to posterior fusions, with the overall percent of cases with complications being 2.40% for anterior decompression, 3.44% for anterior fusion, and 10.49% for posterior fusion. (Wang, 2007)

Predictors of outcome of ACDF: Predictors of good outcome include non-smoking, a pre-operative lower pain level, soft disc disease, disease in one level, greater segmental kyphosis pre-operatively, radicular pain without additional neck or lumbar pain, short duration of symptoms, younger age, no use of analgesics, and normal ratings on biopsychosocial tests such as the Distress and Risk Assessment Method (DRAM). Predictors of poor outcomes include non-specific neck pain, psychological distress, psychosomatic problems and poor general health. (Peolsson, 2006) (Peolsson, 2003) See Plate fixation, cervical spine surgery. See also Adjacent segment disease/degeneration (fusion) & Iliac crest donor-site pain treatment.

Milliman Care Guidelines, Inpatient Surgery, 12th Edition

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)