



IRO#
5068 West Plano Parkway Suite 122
Plano, Texas 75093
Phone: (972) 931-5100

DATE OF REVIEW: 04/29/2009

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Lumbar laminectomy, discectomy, foraminotomy and partial facetectomy at L5-S1.

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

This case was reviewed by a Texas licensed MD, specializing in Orthopedic Trauma, Orthopedic Surgery. The physician advisor has the following additional qualifications, if applicable:

ABMS Orthopaedic Surgery

REVIEW OUTCOME:

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld

Health Care Service(s) in Dispute	CPT Codes	Date of Service(s)	Outcome of Independent Review
Lumbar laminectomy, discectomy, foraminotomy and partial facetectomy at L5-S1.			Upheld

INFORMATION PROVIDED TO THE IRO FOR REVIEW:

PATIENT CLINICAL HISTORY [SUMMARY]:

The patient is a male with a history of repetitive lifting lumbar straining injury. The DOI is given as. On that specific date, the patient was lifting a heavy box when the load shifted and he prevented the box from falling by applying additional force to control the weight. He felt severe pain in the lumbar spine. He was initially evaluated at facility. He was diagnosed with "lumbar strain" and was treated with activity modification, medication and physical therapy. The patient has been evaluated by a number of physicians and surgeons. He has consistently complained of severe low back pain and left leg pain interfering with performance of activities of daily living. Physical findings have been limited and not directly attributable to specific nerve root compressive compromise. MRI on 10/07/08 revealed findings compatible with degenerative disc disease at L2-L3 and L5-S1. Epidural steroid injections have not been effective in alleviating symptoms. EMG/NCV study 10/08/09 failed to demonstrate changes compatible with radiculopathy. Designated Doctor Examinations on 11/14/08 and 03/04/09 failed to conclude the patient was at a point of maximum medical improvement. The diagnosis was lumbar strain and herniated nucleus pulposus L5-S1. A surgical recommendation has been made to perform "lumbar laminectomy, discectomy, foraminotomy, and partial facetectomy L5-S1". Request for pre authorization was denied; reconsideration was submitted and denied.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION:

The below cited passage from the ODG, 2009, low back chapter includes a number of criteria that have to be met to justify pre authorization for the laminectomy, discectomy, foraminotomy and partial facetectomy. Many of these criteria have not been met. Prior denials of the preauthorization request have been appropriate and should be upheld. For example, there are no objective physical findings to suggest specific nerve root compressive compromise. The EMG/NCV study is normal. The MRI fails to demonstrate herniated fragment of nucleus pulposus impinging on a specific nerve root. The findings are more consistent with degenerative disc disease. In the absence of compressive nerve root compromise, the likelihood of a good or excellent result after the surgery recommended is limited.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

ODG:

Discectomy/ laminectomy	Recommended for indications below. Surgical discectomy for carefully selected patients with radiculopathy due to lumbar disc prolapse provides faster relief from the acute attack than conservative management, although any positive or negative effects on the lifetime natural history of the underlying disc disease are still unclear. Unequivocal objective findings are required based on neurological examination and testing. (Gibson-Cochrane, 2000) (Malter, 1996) (Stevens, 1997) (Stevenson, 1995) (BlueCross BlueShield, 2002) (Buttermann, 2004) Standard discectomy and microdiscectomy are of similar efficacy in treatment of herniated disc. (Bigos, 1999)
----------------------------	--

While there is evidence in favor of discectomy for prolonged symptoms of lumbar disc herniation, in patients with a shorter period of symptoms but no absolute indication for surgery, there are only modest short-term benefits, although discectomy seemed to be associated with a more rapid initial recovery, and discectomy was superior to conservative treatment when the herniation was at L4-L5. (Osterman, 2006) The SPORT studies concluded that both lumbar discectomy and nonoperative treatment resulted in substantial improvement after 2 years, but those who chose discectomy reported somewhat greater improvements than patients who elected nonoperative care. (Weinstein, 2006) (Weinstein2, 2006) A recent RCT compared decompressive surgery with nonoperative measures in the treatment of patients with lumbar spinal stenosis, and concluded that, although patients improved over the 2-year follow-up regardless of initial treatment, those undergoing decompressive surgery reported greater improvement regarding leg pain, back pain, and overall disability, but the relative benefit of initial surgical treatment diminished over time while still remaining somewhat favorable at 2 years. (Malmivaara, 2007) Patients undergoing lumbar discectomy are generally satisfied with the surgery, but only half are satisfied with preoperative patient information. (Ronnberg, 2007) If patients are pain free, there appears to be no contraindication to their returning to any type of work after lumbar discectomy. A regimen of stretching and strengthening the abdominal and back muscles is a crucial aspect of the recovery process. (Burnett, 2006) According to a major recent trial, early surgery (microdiscectomy) in patients with 6-12 weeks of severe sciatica caused by herniated disks is associated with better short-term outcomes, but at 1 year, disability outcomes of early surgery vs conservative treatment with eventual surgery if needed are similar. The median time to recovery was 4.0 weeks for early surgery and 12.1 weeks for prolonged conservative treatment. The authors concluded, "Patients whose pain is controlled in a manner that is acceptable to them may decide to postpone surgery in the hope that it will not be needed, without reducing their chances for complete recovery at 12 months. Although both strategies have similar outcomes after 1 year, early surgery remains a valid treatment option for well-informed patients." (Peul-NEJM, 2007) (Devo-NEJM, 2007) A recent randomized controlled trial comparing decompression with decompression and instrumented fusion in patients with foraminal stenosis and single-level degenerative disease found that patients universally improved with surgery, and this improvement was maintained at 5 years. However, no obvious additional benefit was noted by combining decompression with an instrumented fusion. (Hallett, 2007) A recent British study found that lumbar discectomy improved patients' self-reported overall physical health more than other elective surgeries. (Guilfoyle, 2007) Microscopic sequestrectomy may be an alternative to standard microdiscectomy. In this RCT, both groups showed dramatic improvement. (Barth, 2008) There is consistent evidence that for patients with a herniated disk, discectomy is associated with better short-term outcomes than continued conservative management, although outcomes begin to look similar after 3 to 6 months. This is a decision to be made with the patients, discussing the likelihood that they are going to improve either way but will improve faster with surgery. Similar evidence supports the use of surgery for spinal stenosis, although the outcomes look better with surgery out to about 2 years. (Chou, 2008) Standard open discectomy is moderately cost-effective compared with nonsurgical treatment, a new Spine Patient Outcomes Research Trial (SPORT) study shows. The costs per quality-adjusted life-year gained with surgery compared with nonoperative treatment, including work-related productivity costs, ranges from \$34,355 to \$69,403, depending on the cost of surgery. It is wise and proper to wait before initiating surgery, but if the patient continues to experience pain and is missing work, then the higher-cost option such as surgery may be worthwhile. (Tosteson, 2008) Note: Surgical decompression of a lumbar nerve root or roots may include the following procedures: discectomy or microdiscectomy (partial removal of the disc) and laminectomy, hemilaminectomy, laminotomy, or foraminotomy (providing access by partial or total removal of various parts of vertebral bone). Discectomy is the surgical removal of herniated disc material that presses on a nerve root or the spinal cord. A laminectomy is often involved to permit access to the intervertebral disc in a traditional discectomy.

Patient Selection: Microdiscectomy for symptomatic lumbar disc herniations in

patients with a preponderance of leg pain who have failed nonoperative treatment demonstrated a high success rate based on validated outcome measures (80% decrease in VAS leg pain score of greater than 2 points), patient satisfaction (85%), and return to work (84%). Patients should be encouraged to return to their preinjury activities as soon as possible with no restrictions at 6 weeks. Overall, patients with sequestered lumbar disc herniations fared better than those with extruded herniations, although both groups consistently had better outcomes than patients with contained herniations. Patients with herniations at the L5-S1 level had significantly better outcomes than did those at the L4-L5 level. Lumbar disc herniation level and type should be considered in preoperative outcomes counseling. Smokers had a significantly lower return to work rate. In the carefully screened patient, lumbar microdiscectomy for symptomatic disc herniation results in an overall high success rate, patient satisfaction, and return to physically demanding activities. ([Dewing, 2008](#)) Workers' comp back surgery patients are at greater risk for poor lumbar discectomy outcomes than noncompensation patients. ([DeBerard, 2008](#))

Spinal Stenosis: For patients with lumbar spinal stenosis, standard posterior decompressive laminectomy alone (without discectomy) offers a significant advantage over nonsurgical treatment. Discectomy should be reserved for those conditions of disc herniation causing radiculopathy. (See Indications below.) Laminectomy may be used for spinal stenosis secondary to degenerative processes exhibiting ligamentum hypertrophy, facet hypertrophy, and disc protrusion, in addition to anatomical derangements of the spinal column such as tumor, trauma, etc. ([Weinstein, 2008](#)) ([Katz, 2008](#)) See also [Laminectomy](#).

Recent Research: Four-year results for the Dartmouth Spine Patient Outcomes Research Trial (SPORT, n= 1244) indicated that patients who underwent standard open discectomy for a lumbar disc herniation achieved significantly greater improvement than nonoperatively treated patients (using recommended treatments - active physical therapy, home exercise instruction, and NSAIDs) in all primary and secondary outcomes except work status (78.4% for the surgery group compared with 84.4%). Although patients receiving surgery did better generally, all patients in the study improved. Consequently, for patients who don't want an operation no matter how bad their pain is, this study suggests that they will improve and they will not have complications (e.g., paralysis) from nonoperative treatment, but those patients whose leg pain is severe and is limiting their function, who meet the ODG criteria for discectomy, can do better with surgery than without surgery, and the risks are extremely low. ([Weinstein2, 2008](#)) In most patients with low back pain, symptoms resolve without surgical intervention. ([Madigan, 2009](#)) This study showed that surgery for disc herniation was not as successful as total hip replacement but was comparable to total knee replacement in success. Pain was reduced to within 60% of normal levels, function improved to 65% normal, and quality of life was improved by about 50%. The study compared the gains in quality of life achieved by total hip replacement, total knee replacement, surgery for spinal stenosis, disc excision for lumbar disc herniation, and arthrodesis for chronic low back pain. ([Hansson, 2008](#))

ODG Indications for Surgery™ -- Discectomy/laminectomy --

Required symptoms/findings; imaging studies; & conservative treatments below:

I. Symptoms/Findings which confirm presence of radiculopathy. Objective findings on examination need to be present. For unequivocal evidence of radiculopathy, see AMA Guides, 5th Edition, page 382-383. ([Andersson, 2000](#)) Straight leg raising test, crossed straight leg raising and reflex exams should correlate with symptoms and imaging.

Findings require ONE of the following:

A. L3 nerve root compression, requiring ONE of the following:

1. Severe unilateral quadriceps weakness/mild atrophy
2. Mild-to-moderate unilateral quadriceps weakness
3. Unilateral hip/thigh/knee pain

B. L4 nerve root compression, requiring ONE of the following:

1. Severe unilateral quadriceps/anterior tibialis weakness/mild atrophy
2. Mild-to-moderate unilateral quadriceps/anterior tibialis weakness
3. Unilateral hip/thigh/knee/medial pain

C. L5 nerve root compression, requiring ONE of the following:

1. Severe unilateral foot/toe/dorsiflexor weakness/mild atrophy
2. Mild-to-moderate foot/toe/dorsiflexor weakness
3. Unilateral hip/lateral thigh/knee pain

D. S1 nerve root compression, requiring ONE of the following:

1. Severe unilateral foot/toe/plantar flexor/hamstring weakness/atrophy
2. Moderate unilateral foot/toe/plantar flexor/hamstring weakness
3. Unilateral buttock/posterior thigh/calf pain

(EMGs are optional to obtain unequivocal evidence of radiculopathy but not necessary if radiculopathy is already clinically obvious.)

II. Imaging Studies, requiring ONE of the following, for concordance between radicular findings on radiologic evaluation and physical exam findings:

A. Nerve root compression (L3, L4, L5, or S1)

B. Lateral disc rupture

C. Lateral recess stenosis

Diagnostic imaging modalities, requiring ONE of the following:

1. MR imaging
2. CT scanning

	<ol style="list-style-type: none">3. Myelography4. CT myelography & X-Ray <p>III. <u>Conservative Treatments</u>, requiring ALL of the following:</p> <ol style="list-style-type: none">A. Activity modification (not bed rest) after patient education (>= 2 months)B. Drug therapy, requiring at least ONE of the following:<ol style="list-style-type: none">1. NSAID drug therapy2. Other analgesic therapy3. Muscle relaxants4. Epidural Steroid Injection (ESI)C. Support provider referral, requiring at least ONE of the following (in order of priority):<ol style="list-style-type: none">1. Physical therapy (teach home exercise/stretching)2. Manual therapy (chiropractor or massage therapist)3. Psychological screening that could affect surgical outcome <ol style="list-style-type: none">4. Back school (Fisher, 2004)
--	---

TEXAS DEPARTMENT OF INSURANCE COMPLAINT PROCESS: The Texas Department of Insurance requires Independent Review Organizations to be licensed to perform Independent Review in Texas. To contact the Texas Department of Insurance regarding any complaint, you may call or write the Texas Department of Insurance. The telephone number is 1-800-578-4677 or in writing at: Texas Department of Insurance, PO Box 149104 Austin TX, 78714. In accordance with Rule 102.4(h), a copy of this Independent Review Organization (IRO) Decision was sent to the carrier, the requestor and claimant via facsimile or U.S. Postal Service from the office of the IRO on 04/29/2009.