

Clear Resolutions Inc.

An Independent Review Organization
7301 RANCH RD 620 N, STE 155-199A
Austin, TX 78726
Phone: (512) 772-4390
Fax: (512) 519-7316
Email: resolutions.manager@cri-iro.com

NOTICE OF INDEPENDENT REVIEW DECISION

DATE OF REVIEW:

Jul/30/2009

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

SI joint injection, under fluoroguide.

DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

MD, Board Certified in Physical Medicine and Rehabilitation
Subspecialty Board Certified in Pain Management
Subspecialty Board Certified in Electrodiagnostic Medicine
Residency Training PMR and Orthopedic Surgery

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Overturned (Disagree)

Partially Overturned (Agree in part/Disagree in part)

INFORMATION PROVIDED TO THE IRO FOR REVIEW

ODG Guidelines and Treatment Guidelines
Adverse Determination Letters, 6/17/09, 5/8/09
Pain Consultant Associates, 4/27/09
MRI of the Lumbar Spine, 9/11/08

PATIENT CLINICAL HISTORY SUMMARY

This is a man involved in an MVA in xx/xxxx. He had a pelvic fracture and tibia fibula fracture that required surgery. He describes chronic low back pain to the left buttock. There is no radicular component of the pain. Dr. 's examination showed paravertebral tenderness and left sided SI joint pain and a positive bilateral Patrick sign. The lumbar MRI showed minimal disc bulge at L5/S1. He advised physical therapy and then possible SI injections when seen in April 2009.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION

The ODG requires conservative care. It has been more than a year since the injury. Dr. advised physical therapy. The records are unclear as to whether the PT was attempted or what the outcome was. The ODG requires 3 positive signs on physical examination. Only a positive Patrick sign was provided. The request does not meet the criteria. The reviewer finds that medical necessity does not exist for SI joint injection, under fluoroguide.

Sacroiliac joint injections (SJI)

Recommended as an option if failed at least 4-6 weeks of aggressive conservative therapy. See the Hip & Pelvis Chapter for more information, references, and ODG Criteria for the use of sacroiliac blocks

Sacroiliac joint block

Recommended as an option if failed at least 4-6 weeks of aggressive conservative therapy as indicated below. Sacroiliac dysfunction is poorly defined and the diagnosis is often difficult to make due to the presence of other low back pathology (including spinal stenosis and facet arthropathy). The diagnosis is also difficult to make as pain symptoms may depend on the region of the SI joint that is involved (anterior, posterior, and/or extra-articular ligaments). Pain may radiate into the buttock, groin and entire ipsilateral lower limb, although if pain is present above L5, it is not thought to be from the SI joint.

Innervation: The anterior portion is thought to be innervated by the posterior rami of the L1-S2 roots and the posterior portion by the posterior rami of L4-S3. although the actual innervation remains unclear. Anterior innervation may also be supplied by the obturator nerve, superior gluteal nerve and/or lumbosacral trunk. (Vallejo, 2006) Other research supports innervation by the S1 and S2 sacral dorsal rami

Etiology: includes degenerative joint disease, joint laxity, and trauma (such as a fall to the buttock). The main cause is SI joint disruption from significant pelvic trauma.

Diagnosis: Specific tests for motion palpation and pain provocation have been described for SI joint dysfunction: Cranial Shear Test; Extension Test; Flamingo Test; Fortin Finger Test; Gaenslen's Test; Gillet's Test (One Legged-Stork Test); Patrick's Test (FABER); Pelvic Compression Test; Pelvic Distraction Test; Pelvic Rock Test; Resisted Abduction Test (REAB); Sacroiliac Shear Test; Standing Flexion Test; Seated Flexion Test; Thigh Thrust Test (POSH). Imaging studies are not helpful. It has been questioned as to whether SI joint blocks are the "diagnostic gold standard." The block is felt to show low sensitivity, and discordance has been noted between two consecutive blocks (questioning validity). (Schwarzer, 1995) There is also concern that pain relief from diagnostic blocks may be confounded by infiltration of extra-articular ligaments, adjacent muscles, or sheaths of the nerve roots themselves. Sacral lateral branch injections have demonstrated a lack of diagnostic power and area not endorsed for this purpose. (Yin, 2003)

Treatment: There is limited research suggesting therapeutic blocks offer long-term effect. There should be evidence of a trial of aggressive conservative treatment (at least six weeks of a comprehensive exercise program, local icing, mobilization/manipulation and anti-inflammatories) as well as evidence of a clinical picture that is suggestive of sacroiliac injury and/or disease prior to a first SI joint block. If helpful, the blocks may be repeated; however, the frequency of these injections should be limited with attention placed on the comprehensive exercise program. (Forst, 2006) (Berthelot, 2006) (van der Wurff, 2006) (Laslett, 2005) (Zelle, 2005) (McKenzie-Brown 2005) (Pekkafehli, 2003) (Manchikanti, 2003) (Slipman, 2001) (Nelemans-Cochrane, 2000) See also Intra-articular steroid hip injection; & Sacroiliac joint radiofrequency neurotomy

Criteria for the use of sacroiliac blocks

1. The history and physical should suggest the diagnosis (with documentation of at least 3

positive exam findings as listed above)

2. Diagnostic evaluation must first address any other possible pain generators
3. The patient has had and failed at least 4-6 weeks of aggressive conservative therapy including PT, home exercise and medication management
4. Blocks are performed under fluoroscopy. (Hansen, 2003)
5. A positive diagnostic response is recorded as 80% for the duration of the local anesthetic. If the first block is not positive, a second diagnostic block is not performed
6. If steroids are injected during the initial injection, the duration of pain relief should be at least 6 weeks with at least > 70% pain relief recorded for this period
7. In the treatment or therapeutic phase (after the stabilization is completed), the suggested frequency for repeat blocks is 2 months or longer between each injection, provided that at least >70% pain relief is obtained for 6 weeks
8. The block is not to be performed on the same day as a lumbar epidural steroid injection (ESI), transforaminal ESI, facet joint injection or medial branch block
9. In the treatment or therapeutic phase, the interventional procedures should be repeated only as necessary judging by the medical necessity criteria, and these should be limited to a maximum of 4 times for local anesthetic and steroid blocks over a period of 1 year.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION

ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE

AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES

DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES

EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN

INTERQUAL CRITERIA

MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

MERCY CENTER CONSENSUS CONFERENCE GUIDELINES

MILLIMAN CARE GUIDELINES

ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR

TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS

TEXAS TACADA GUIDELINES

TMF SCREENING CRITERIA MANUAL

PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)

OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES
(PROVIDE A DESCRIPTION)