

SENT VIA EMAIL OR FAX ON  
Jul/02/2009

## **P-IRO Inc.**

An Independent Review Organization  
835 E. Lamar Blvd. #394  
Arlington, TX 76011  
Phone: (817) 349-6420  
Fax: (866) 328-3894  
Email: resolutions.manager@p-iro.com

**DATE OF REVIEW:**

Jun/26/2009

**IRO CASE #:**

**DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:**

Lumbar ESI X 2 under Fluroscopic Guidance and X-Ray

**DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:**

Board Certified in Physical Medicine and Rehabilitation  
Subspecialty Board Certified in Pain Management  
Subspecialty Board Certified in Electrodiagnostic Medicine  
Residency Training PMR and ORTHOPAEDIC SURGERY

**REVIEW OUTCOME:**

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Overturned (Disagree)

Partially Overturned (Agree in part/Disagree in part)

**INFORMATION PROVIDED TO THE IRO FOR REVIEW**

**PATIENT CLINICAL HISTORY SUMMARY**

This is a man who was injured when he fell on his buttocks on xx-xx-xx. An MRI on 6/20/08 showed multiple level degenerative changes at L3/L4, L4/5 and L5/S1. There were disc bulges, facet arthropathy, and some L5/S1 foraminal stenosis, but no disc herniations and no report of any nerve root compromise. Subsequent xrays done in 1/09 did not show any significant change. Dr. described pain in the right S1 distribution, but found no atrophy or neurological abnormalities. He wants to proceed with ESI injections.

**ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION**

There are both diagnostic and therapeutic criteria. The diagnostic block is indicted when the radiological findings and physical findings are inconsistent. This man has no specific physical findings of a radiculopathy as defined in the AMA Guides. The radiological studies did not

show any probable source of a pain generator. The absence of a documented radiculopathy per the criteria provided, even in the presence of symptoms, does not justify the epidural injection.

## The AMA Guides

“...For reflex abnormalities to be considered valid, the involved and normal limb(s) should show marked asymmetry...”

### “Weakness and Loss of Sensatio

“To be valid, the sensory findings must be in a strict anatomic distribution, i.e follow dermatomal patterns...Motor findings should be consistent with the affected nerve structures(s). Significant, long standing weakness is usually accompanied by atrophy.”

### “Radiculopath

Radiculopathy for the purposes of the Guides is defined as significant alteration in the function of a nerve root or nerve roots and is usually caused by pressure on one or several nerve roots. The diagnosis requires a dermatomal distribution of pain, numbness, and/or paresthesias in a dermatomal distribution. The diagnosis of herniated disc must be substantiated by an appropriate finding on the imaging study. The presence of findings on a imaging study in and of itself does not make the diagnosis of radiculopathy. There must also be evidence as described above. “

### “Atroph

Atrophy is measured with a tape measure at identical levels on both limbs. For reasons or reproducibility, the difference in circumference should be 2cm or greater in the thigh and 1cm or greater in the arm, forearm, or leg...”

### “Electrodiagnostic verification of Radiculopathy

Unequivocal electrodiagnostic evidence of acute nerve root pathology includes the presence of multiple positive sharp waves or fibrillation potentials in muscles innervated by one nerve root. However the quality of the person performing and interpreting the study is critical. Electromyography should be performed only by a licensed physician qualified by reason of education, training and experience in these procedures. Electromyography does not detect all compressive radiculopathies and cannot determine the cause of the nerve root pathology. On the other hand, electromyography can detect noncompressive radiculopathies, which are not identified by imaging studies. “

Page 382-382. AMA Guides to the Evaluation of Permanent Impairment. 5th edition

## Epidural steroid injections (ESIs), therapeuti

Recommended as a possible option for short-term treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy) with use in conjunction with active rehab efforts. See specific criteria for use below. Radiculopathy symptoms are generally due to herniated nucleus pulposus or spinal stenosis, although ESIs have not been found to be as beneficial a treatment for the latter condition....

## Criteria for the use of Epidural steroid injections

Note: The purpose of ESI is to reduce pain and inflammation, thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit

(1) Radiculopathy must be documented. Objective findings on examination need to be present. For unequivocal evidence of radiculopathy, see AMA Guides, 5th Edition, page 382-383. (Andersson, 2000)

(2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants)

(3) Injections should be performed using fluoroscopy (live x-ray) and injection of contrast for guidance

(4) Diagnostic Phase: At the time of initial use of an ESI (formally referred to as the “diagnostic phase” as initial injections indicate whether success will be obtained with this treatment intervention), a maximum of one to two injections should be performed. A repeat block is not recommended if there is inadequate response to the first block (< 30% is a standard placebo response). A second block is also not indicated if the first block is accurately placed unless: (a) there is a question of the pain generator; (b) there was possibility of inaccurate placement; or (c) there is evidence of multilevel pathology. In these cases a different level or approach might be proposed. There should be an interval of at least one to two weeks between injections

(5) No more than two nerve root levels should be injected using transforaminal blocks

(6) No more than one interlaminar level should be injected at one session

(7) Therapeutic phase: If after the initial block/blocks are given (see “Diagnostic Phase” above) and found to produce pain relief of at least 50-70% pain relief for at least 6-8 weeks, additional blocks may be required. This is generally referred to as the “therapeutic phase.” Indications for repeat blocks include acute exacerbation of pain, or new onset of symptoms. The general consensus recommendation is for no more than 4 blocks per region per year. (CMS, 2004) (Boswell, 2007)

(8) Repeat injections should be based on continued objective documented pain relief, decreased need for pain medications, and functional response

(9) Current research does not support a routine use of a “series-of-three” injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections for the initial phase and rarely more than 2 for therapeutic treatment

(10) It is currently not recommended to perform epidural blocks on the same day of treatment as facet blocks or sacroiliac blocks or lumbar sympathetic blocks or trigger point injections as this may lead to improper diagnosis or unnecessary treatment

(11) Cervical and lumbar epidural steroid injection should not be performed on the same day. (Doing both injections on the same day could result in an excessive dose of steroids, which can be dangerous, and not worth the risk for a treatment that has no long-term benefit.)

#### **A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION**

[ ] ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE

[ ] AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES

[ ] DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES

- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER ERVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)