mcmc
managing care. managing claims.
88 Black Falcon Avenue, Suite 353 Boston, MA 02210 (T) 800-227-1464 (F) 617-375-7777

Notice of Independent Review Decision

DATE OF REVIEW:
10/09/2009

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE
Anterior and posterior lumbar decompression with fusion with length of stay five days

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION
Board Certified Orthopaedic Surgeon

REVIEW OUTCOME
Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be: **Upheld**

Provide a description of the review outcome that clearly states whether or not medical necessity exists for each of the health care services in dispute.
The requested procedure (anterior and posterior lumbar decompression with fusion with length of stay five days) is not medically necessary, reasonable, related or supported by the evidence-based Official Disability Guidelines (ODG).

INFORMATION PROVIDED TO THE IRO FOR REVIEW
•
•

PATIENT CLINICAL HISTORY [SUMMARY]:
The injured individual is a male who was reported to have sustained a work-related injury on xx/xx/xx. The mechanism of injury was a lifting injury. The first medical available was a letter dated 09/28/04 denying a request for a posterior decompression and fusion. The next note is an operative report for an endoscopic lumbar decompression with annuloplasty on the right at L4-L5 performed by M.D. There is a gap in documentation until 01/11/09 when a MRI report revealed multi-level degenerative disc disease including an old compression fracture at L1-L2 with retropulsion. An electromyogram done on 09/24/08 was normal with no evidence of radiculopathy. There is no information regarding how the injured individual did following the previous procedure. Dr. evaluated the injured individual on 02/24/09. He recommended a surgical procedure but it was unclear at what levels. Straight leg raise was negative. He reported a mild decrease extensor hallices longus (EHL) and anterior tibialis strength. He also noted grade I spondylothesis of L4 on L5 with instability of 4mm. The procedure was noted to be denied on 03/27/09. The injured individual was then followed by a Physician Assistant (PA) in Dr. 's office. The PA noted in his notes that an IRO was pending. The requested procedure was denied on 08/14/09 on initial review by a neurosurgeon. He noted that the injured

individual had a previous surgery at L4- L5. He felt that the request did not meet ODG criteria. An Orthopaedist denied the request on reconsideration/appeal on 08/27/09. He felt that the injured individual may require surgery but the exact levels were unclear. Both reviewers attempted to speak to Dr. without success.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

The injured individual is a male who has documented multi-level degenerative disc disease. He has undergone a previous surgical procedure with unclear outcome. Electromyogram (EMG) study done in the past has been normal.

ODG Indications for Surgery: -- Discectomy/laminectomy --
Required symptoms/findings; imaging studies; & conservative treatments below:
I. Symptoms/Findings which confirm presence of radiculopathy. Objective findings on examination need to be present. For unequivocal evidence of radiculopathy, see AMA Guides, 5th Edition, page 382-383. (Andersson, 2000) Straight leg raising test, crossed straight leg raising and reflex exams should correlate with symptoms and imaging.
Findings require ONE of the following:

A. L3 nerve root compression, requiring ONE of the following:
   1. Severe unilateral quadriceps weakness/mild atrophy
   2. Mild-to-moderate unilateral quadriceps weakness
   3. Unilateral hip/thigh/knee pain
B. L4 nerve root compression, requiring ONE of the following:
   1. Severe unilateral quadriceps/anterior tibialis weakness/mild atrophy
   2. Mild-to-moderate unilateral quadriceps/anterior tibialis weakness
   3. Unilateral hip/thigh/knee/medial pain
C. L5 nerve root compression, requiring ONE of the following:
   1. Severe unilateral foot/toe/dorsiflexor weakness/mild atrophy
   2. Mild-to-moderate foot/toe/dorsiflexor weakness
   3. Unilateral hip/lateral thigh/knee pain
D. S1 nerve root compression, requiring ONE of the following:
   1. Severe unilateral foot/toe/plantar flexor/hamstring weakness/atrophy
   2. Moderate unilateral foot/toe/plantar flexor/hamstring weakness
   3. Unilateral buttock/posterior thigh/calf pain

(EMGs are optional to obtain unequivocal evidence of radiculopathy but not necessary if radiculopathy is already clinically obvious.)

II. Imaging Studies, requiring ONE of the following, for concordance between radicular findings on radiologic evaluation and physical exam findings:

A. Nerve root compression (L3, L4, L5, or S1)
B. Lateral disc rupture
C. Lateral recess stenosis

Diagnostic imaging modalities, requiring ONE of the following:
   1. MR imaging
   2. CT scanning
   3. Myelography
   4. CT myelography & X-Ray

III. Conservative Treatments, requiring ALL of the following:

A. Activity modification (not bed rest) after patient education (>= 2 months)
B. Drug therapy, requiring at least ONE of the following:
   1. NSAID drug therapy
   2. Other analgesic therapy
   3. Muscle relaxants
   4. Epidural Steroid Injection (ESI)
C. Support provider referral, requiring at least ONE of the following (in order of priority):
   1. Physical therapy (teach home exercise/stretching)
   2. Manual therapy (chiropractor or massage therapist)
   3. Psychological screening that could affect surgical outcome
   4. Back school (Fisher, 2004)

There is no information regarding an adequate trial of conservative treatment to include active physical therapy. There is no objective evidence of radiculopathy on EMG.

Discectomy/ laminectomy: Recommended for indications below. Surgical discectomy for carefully selected patients with radiculopathy due to lumbar disc prolapse provides faster relief from the acute attack than conservative management, although any positive or negative effects on the lifetime natural history of the underlying disc disease are still unclear. Unequivocal objective findings are required based on neurological examination and testing. (Gibson-Cochrane, 2000) (Malter, 1996) (Stevens, 1997) (Stevenson, 1995) (BlueCross BlueShield, 2002) (Buttermann, 2004) Standard discectomy and microdiscectomy are of similar efficacy in treatment of herniated disc. (Bigos, 1999) While there is evidence in favor of discectomy for prolonged symptoms of lumbar disc herniation, in patients with a shorter period of symptoms but no absolute indication for surgery, there are only modest short-term benefits, although discectomy seemed to be associated with a more rapid initial recovery, and discectomy was superior to conservative treatment when the herniation was at L4-L5. (Osterman, 2006) The SPORT studies concluded that both lumbar discectomy and nonoperative treatment resulted in substantial improvement after 2 years, but those who chose discectomy reported somewhat greater improvements than patients who elected nonoperative care. (Weinstein, 2006) (Weinstein2, 2006) A recent RCT compared decompressive surgery with nonoperative measures in the treatment of patients with lumbar spinal stenosis, and concluded that, although patients improved over the 2-year follow-up regardless of initial treatment, those undergoing decompressive surgery reported greater improvement regarding leg pain, back pain, and overall disability, but the relative benefit of initial surgical treatment diminished over time while still remaining somewhat favorable at 2 years. (Malmivaara, 2007) Patients undergoing lumbar discectomy are generally satisfied with the surgery, but only half are satisfied with preoperative patient information. (Ronnberg, 2007) If patients are pain free, there appears to be no contraindication to their returning to any type of work after lumbar discectomy. A regimen of stretching and strengthening the abdominal and back muscles is a crucial aspect of the recovery process. (Burnett, 2006) According to a major recent trial, early surgery (microdiscectomy) in patients with 6-12 weeks of severe sciatica caused by herniated disks is associated with better short-term outcomes, but at 1 year, disability outcomes of early surgery vs conservative treatment with eventual surgery if needed are similar. The median time to recovery was 4.0 weeks for early surgery and 12.1 weeks for prolonged conservative treatment. The authors concluded, "Patients whose pain is controlled in a manner that is acceptable to them may decide to postpone surgery in the hope that it will not be needed, without reducing their chances for complete recovery at 12 months. Although both strategies have similar outcomes after 1 year, early surgery remains a valid treatment option for well-informed patients." (Peul-NEJM, 2007) (Deyo-NEJM, 2007) A recent randomized controlled trial comparing decompression with decompression and

instrumented fusion in patients with foraminal stenosis and single-level degenerative disease found that patients universally improved with surgery, and this improvement was maintained at 5 years. However, no obvious additional benefit was noted by combining decompression with an instrumented fusion. (Hallett, 2007) A recent British study found that lumbar discectomy improved patients' self-reported overall physical health more than other elective surgeries. (Guilfoyle, 2007) Microscopic sequestrectomy may be an alternative to standard microdiscectomy. In this RCT, both groups showed dramatic improvement. (Barth, 2008) There is consistent evidence that for patients with a herniated disk, discectomy is associated with better short-term outcomes than continued conservative management, although outcomes begin to look similar after 3 to 6 months. This is a decision to be made with the patients, discussing the likelihood that they are going to improve either way but will improve faster with surgery. Similar evidence supports the use of surgery for spinal stenosis, although the outcomes look better with surgery out to about 2 years. (Chou, 2008) Standard open discectomy is moderately cost-effective compared with nonsurgical treatment, a new Spine Patient Outcomes Research Trial (SPORT) study shows. The costs per quality-adjusted life-year gained with surgery compared with nonoperative treatment, including work-related productivity costs, ranges from $34,355 to $69,403, depending on the cost of surgery. It is wise and proper to wait before initiating surgery, but if the patient continues to experience pain and is missing work, then the higher-cost option such as surgery may be worthwhile. (Tosteson, 2008) Note: Surgical decompression of a lumbar nerve root or roots may include the following procedures: discectomy or microdiscectomy (partial removal of the disc) and laminectomy, hemilaminectomy, laminotomy, or foraminotomy (providing access by partial or total removal of various parts of vertebral bone). Discectomy is the surgical removal of herniated disc material that presses on a nerve root or the spinal cord. A laminectomy is often involved to permit access to the intervertebral disc in a traditional discectomy.
Patient Selection: Microdiscectomy for symptomatic lumbar disc herniations in patients with a preponderance of leg pain who have failed nonoperative treatment demonstrated a high success rate based on validated outcome measures (80% decrease in VAS leg pain score of greater than 2 points), patient satisfaction (85%), and return to work (84%). Patients should be encouraged to return to their preinjury activities as soon as possible with no restrictions at 6 weeks. Overall, patients with sequestered lumbar disc herniations fared better than those with extruded herniations, although both groups consistently had better outcomes than patients with contained herniations. Patients with herniations at the L5-S1 level had significantly better outcomes than did those at the L4-L5 level. Lumbar disc herniation level and type should be considered in preoperative outcomes counseling. Smokers had a significantly lower return to work rate. In the carefully screened patient, lumbar microdiscectomy for symptomatic disc herniation results in an overall high success rate, patient satisfaction, and return to physically demanding activities. (Dewing, 2008) Workers' comp back surgery patients are at greater risk for poor lumbar discectomy outcomes than noncompensation patients. (DeBerard, 2008)
Spinal Stenosis: For patients with lumbar spinal stenosis, standard posterior decompressive laminectomy alone (without discectomy) offers a significant advantage over nonsurgical treatment. Discectomy should be reserved for those conditions of disc herniation causing radiculopathy. (See Indications below.) Laminectomy may be used for spinal stenosis secondary to degenerative processes exhibiting ligamental hypertrophy, facet hypertrophy, and disc protrusion, in addition to anatomical derangements of the spinal column such as tumor, trauma, etc. (Weinstein, 2008) (Katz, 2008) See also Laminectomy.
Recent Research: Four-year results for the Dartmouth Spine Patient Outcomes Research Trial (SPORT, n= 1244) indicated that patients who underwent standard open discectomy for a lumbar disc herniation achieved significantly greater improvement than nonoperatively treated patients (using recommended treatments - active physical therapy, home exercise instruction, and NSAIDs) in all

primary and secondary outcomes except work status (78.4% for the surgery group compared with 84.4%). Although patients receiving surgery did better generally, all patients in the study improved. Consequently, for patients who don't want an operation no matter how bad their pain is, this study suggests that they will improve and they will not have complications (e.g., paralysis) from nonoperative treatment, but those patients whose leg pain is severe and is limiting their function, who meet the ODG criteria for discectomy, can do better with surgery than without surgery, and the risks are extremely low. (Weinstein2, 2008) In most patients with low back pain, symptoms resolve without surgical intervention. (Madigan, 2009) This study showed that surgery for disc herniation was not as successful as total hip replacement but was comparable to total knee replacement in success. Pain was reduced to within 60% of normal levels, function improved to 65% normal, and quality of life was improved by about 50%. The study compared the gains in quality of life achieved by total hip replacement, total knee replacement, surgery for spinal stenosis, disc excision for lumbar disc herniation, and arthrodesis for chronic low back pain. (Hansson, 2008) For radiculopathy with herniated lumbar disc, there is good evidence that standard open discectomy and microdiscectomy are moderately superior to nonsurgical therapy for improvement in pain and function through 2 to 3 months, but patients on average experience improvement either with or without surgery, and benefits associated with surgery decrease with long-term follow-up. (Chou, 2009

Patient Selection Criteria for Lumbar Spinal Fusion:

For chronic low back problems, fusion should not be considered within the first 6 months of symptoms, except for fracture, dislocation or progressive neurologic loss. Indications for spinal fusion may include: (1) Neural Arch Defect - Spondylolytic spondylolisthesis, congenital neural arch hypoplasia. (2) Segmental Instability (objectively demonstrable) - Excessive motion, as in degenerative spondylolisthesis, surgically induced segmental instability and mechanical intervertebral collapse of the motion segment and advanced degenerative changes after surgical discectomy. [For excessive motion criteria, see AMA Guides, 5th Edition, page 384 (relative angular motion greater than 20 degrees). (Andersson, 2000) (Luers, 2007)] (3) Primary Mechanical Back Pain (i.e., pain aggravated by physical activity)/Functional Spinal Unit Failure/Instability, including one or two level segmental failure with progressive degenerative changes, loss of height, disc loading capability. In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. There is a lack of support for fusion for mechanical low back pain for subjects with failure to participate effectively in active rehab pre-op, total disability over 6 months, active psych diagnosis, and narcotic dependence. [For spinal instability criteria, see AMA Guides, 5th Edition, page 379 (lumbar inter-segmental movement of more than 4.5 mm). (Andersson, 2000)] (4) Revision Surgery for failed previous operation(s) if significant functional gains are anticipated. Revision surgery for purposes of pain relief must be approached with extreme caution due to the less than 50% success rate reported in medical literature. (5) Infection, Tumor, or Deformity of the lumbosacral spine that cause intractable pain, neurological deficit and/or functional disability. (6) After failure of two discectomies on the same disc, fusion may be an option at the time of the third discectomy, which should also meet the ODG criteria. (See ODG Indications for Surgery -- Discectomy.)

Pre-Operative Surgical Indications Recommended: Pre-operative clinical surgical indications for spinal fusion should include all of the following: (1) All pain generators are identified and treated; & (2) All physical medicine and manual therapy interventions are completed; & (3) X-rays demonstrating spinal instability and/or myelogram, CT-myelogram, or discography (see discography criteria) & MRI demonstrating disc pathology; & (4) Spine pathology limited to two levels; & (5) Psychosocial screen with confounding issues addressed. (6) For any potential fusion surgery, it is recommended that the injured worker refrain from smoking for at least six weeks prior to surgery and during the period of fusion healing. (xxxxx, xxxxx) (xxxxx, xxxxx)

88 Black Falcon Avenue, Suite 353 Boston, MA 02210 (T) 800-227-1464 (F) 617-375-7777

The patient has multi-level disease as documented in the medical record. It is unclear exactly what the pain generators are in this setting. There is no evidence of any attempt to address any psychosocial issues.

Fusion (spinal): Not recommended for patients who have less than six months of failed recommended conservative care unless there is objectively demonstrated severe structural instability and/or acute or progressive neurologic dysfunction, but recommended as an option for spinal fracture, dislocation, spondylolisthesis or frank neurogenic compromise, subject to the selection criteria outlined in the section below entitled, “Patient Selection Criteria for Lumbar Spinal Fusion,” after 6 months of conservative care. For workers’ comp populations, see also the heading, “Lumbar fusion in workers' comp patients.” After screening for psychosocial variables, outcomes are improved and fusion may be recommended for degenerative disc disease with spinal segment collapse with or without neurologic compromise after 6 months of compliance with recommended conservative therapy. [For spinal instability criteria, see AMA Guides (Andersson, 2000)] For complete references, see separate document with all studies focusing on Fusion (spinal). There is limited scientific evidence about the long-term effectiveness of fusion for degenerative disc disease compared with natural history, placebo, or conservative treatment. Studies conducted in order to compare different surgical techniques have shown success for fusion in carefully selected patients. (Gibson-Cochrane, 2000) (Savolainen, 1998) (Wetzel, 2001) (Molinari, 2001) (Bigos, 1999) (Washington, 1995) (DeBarard-Spine, 2001) (Fritzell-Spine, 2001) (Fritzell-Spine, 2002) (Deyo-NEJM, 2004) (Gibson-Cochrane/Spine, 2005) (Soegaard, 2005) (Glassman, 2006) (Atlas, 2006) According to the recently released AANS/NASS Guidelines, lumbar fusion is recommended as a treatment for carefully selected patients with disabling low back pain due to one- or two-level degenerative disc disease after failure of an appropriate period of conservative care. This recommendation was based on one study that contained numerous flaws, including a lack of standardization of conservative care in the control group. At the time of the 2-year follow up it appeared that pain had significantly increased in the surgical group from year 1 to 2. Follow-up post study is still pending publication. In addition, there remains no direction regarding how to define the “carefully selected patient.” (Resnick, 2005) (Fritzell, 2004) A recently published well respected international guideline, the “European Guidelines,” concluded that fusion surgery for nonspecific chronic LBP cannot be recommended unless 2 years of all other recommended conservative treatments – including multidisciplinary approaches with combined programs of cognitive intervention and exercises – have failed, or such combined programs are not available, and only then in carefully selected patients with maximum 2-level degenerative disc disease. (Airaksinen, 2006) For chronic LBP, exercise and cognitive intervention may be equivalent to lumbar fusion without the potentially high surgical complication rates. (Ivar Brox-Spine, 2003) (Keller-Spine, 2004) (Fairbank-BMJ, 2005) (Brox, 2006) In acute spinal cord injury (SCI), if the spine is unstable following injury, surgical fusion and bracing may be necessary. (Bagnall-Cochrane, 2004) (Siebenga, 2006) A study on improving quality through identifying inappropriate care found that use of guideline-based Utilization Review (UR) protocols resulted in a denial rate for lumbar fusion 59 times as high as denial rates using non-guideline based UR. (Wickizer, 2004) The profit motive and market medicine have had a significant impact on clinical practice and research in the field of spine surgery. (Weiner-Spine, 2004) (Shah-Spine, 2005) (Abelson, 2006) Data on geographic variations in medical procedure rates suggest that there is significant variability in spine fusion rates, which may be interpreted to suggest a poor professional consensus on the appropriate indications for performing spinal fusion. (Deyo-Spine, 2005) (Weinstein, 2006) Outcomes from complicated surgical fusion techniques (with internal fixation) may be no better than the traditional posterolateral fusion. (van Tulder, 2006) (Maghout-Juratli, 2006) Despite the new technologies, reoperation rates after lumbar

fusion have become higher. (Martin, 2007) According to the recent Medicare Coverage Advisory Committee Technology Assessment, the evidence for lumbar spinal fusion does not conclusively demonstrate short-term or long-term benefits compared with nonsurgical treatment for elderly patients. (CMS, 2006) When lumbar fusion surgery is performed, either with lateral fusion alone or with interbody fusion, unlike cervical fusion, there is no absolute contraindication to patients returning even to contact sports after complete recovery from surgery. Like patients with a thoracic injury, those with a lumbar injury should be pain free, have no disabling neurological deficit, and exhibit evidence of bone fusion on x-ray films before returning. (Burnett, 2006) A recent randomized controlled trial comparing decompression with decompression and instrumented fusion in patients with foraminal stenosis and single-level degenerative disease found that patients universally improved with surgery, and this improvement was maintained at 5 years. However, no obvious additional benefit was noted by combining decompression with an instrumented fusion. (Hallett, 2007) Discography may be supported if the decision has already been made to do a spinal fusion, and a negative discogram could rule out the need for fusion on that disc (but a positive discogram in itself would not justify fusion). Discography may help distinguish asymptomatic discs among morphologically abnormal discs in patients without psychosocial issues. Precise prospective categorization of discographic diagnoses may predict outcomes from treatment, surgical or otherwise. (Derby, 2005) (Derby2, 2005) (Derby, 1999) New research shows that healthcare expenditures for back and neck problems have increased substantially over time, but with little improvement in healthcare outcomes such as functional disability and work limitations. Rates of imaging, injections, opiate use, and spinal surgery have increased substantially over the past decade, but it is unclear what impact, if any, this has had on health outcomes. (Martin, 2008) The efficacy of surgery for nonspecific back pain is uncertain. There may be some patients for whom surgery, fusion specifically, might be helpful, but it is important for doctors to discuss the fact that surgery doesn't tend to lead to huge improvements on average, about a 10- to 20-point improvement in function on a 100-point scale, and a significant proportion of patients still need to take pain medication and don't return to full function. (Chou, 2008) This study showed that fusion for chronic lower back pain was the least successful common orthopaedic surgery. The study compared the gains in quality of life achieved by total hip replacement, total knee replacement, surgery for spinal stenosis, disc excision for lumbar disc herniation, and arthrodesis for chronic low back pain. For chronic lower back pain, improvements were statistically significant but clinically negligible. Although pain was reduced and function improved slightly, outcomes remained in the moderately affected range, quality of life was not improved and rendered worse, on average. While surgery for spinal stenosis and for disc herniation compare well with archetypical orthopaedic operations, the outcomes of surgery for chronic lower back pain do not even approach those of other orthopaedic procedures, and the data show that patients with back pain are rendered worse off by surgery. (Hansson, 2008) Recent studies document a 220% increase in lumbar spinal fusion surgery rates, without demonstrated improvements in patient outcomes or disability rates. (Deyo, 2009) In a study of 2,378 Washington State workers' compensation claimants who underwent fusion to assess the frequency, timing, and causes of death, the 3-year cumulative mortality rate post-fusion was 1.93% and analgesic-related deaths were responsible for 21% of all deaths and 31.4% of all potential life lost. (Juratli, 2009) A study to compare the surgical experience, clinical outcomes, and effect on body weight between obese and morbidly obese patients undergoing lumbar spine fusion surgery concluded that clinical outcomes were independent of the BMI of the patient, but the incidence of postoperative complications was significant in 45% of morbidly obese and 44% of obese patients. The authors proposed that morbidly obese patients should undergo bariatric surgery before spine surgery in nonemergent situations. (Vaidya, 2009) For nonradicular low back pain with common degenerative changes, there is fair evidence that fusion is no better than intensive rehabilitation with a cognitive-behavioral emphasis for improvement in pain or function, and less than half of patients

experience optimal outcomes (defined as no more than sporadic pain, slight restriction of function, and occasional analgesics) following fusion. (Chou, 2009) Posterolateral bone-grafting fusion is not necessary when a Denis type-B thoracolumbar burst fracture associated with a load-sharing score of <or=6 is treated with short-segment pedicle screw fixation. (Dai, 2009) Lumbar spinal fusion surgeries use bone grafts, and are sometimes combined with metal devices, to produce a rigid connection between two or more adjacent vertebrae. The therapeutic objective of spinal fusion surgery for patients with low back problems is to prevent any movement in the intervertebral spaces between the fused vertebrae, thereby reducing pain and any neurological deficits. See also Adjacent segment disease/degeneration (fusion) & Iliac crest donor-site pain treatment.
Lumbar fusion in workers' comp patients: In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. Until further research is conducted there remains insufficient evidence to recommend fusion for chronic low back pain in the absence of stenosis and spondylolisthesis, and this treatment for this condition remains “under study.” It appears that workers’ compensation populations require particular scrutiny when being considered for fusion for chronic low back pain, as there is evidence of poorer outcomes in subgroups of patients who were receiving compensation or involved in litigation. (Fritzell-Spine, 2001) (Harris-JAMA, 2005) (Maghout-Juratli, 2006) (Atlas, 2006) Despite poorer outcomes in workers’ compensation patients, utilization is much higher in this population than in group health. (Texas, 2001) (NCCI, 2006) Presurgical biopsychosocial variables predict patient outcomes from lumbar fusion, which may help improve patient selection. Workers' compensation status, smoking, depression, and litigation were the most consistent presurgical predictors of poorer patient outcomes. Other predictors of poor results were number of prior low back operations, low household income, and older age. (DeBerard-Spine, 2001) (DeBerard, 2003) (Deyo, 2005) (LaCaille, 2005) (Trief-Spine, 2006) Obesity and litigation in workers' compensation cases predict high costs associated with interbody cage lumbar fusion. (LaCaille, 2007) A recent study of 725 workers' comp patients in Ohio who had lumbar fusion found only 6% were able to go back to work a year later, 27% needed another operation, and over 90% were in enough pain that they were still taking narcotics at follow-up. (Nguyen, 2007)

The criteria as outlined above have not been met.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:
ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES