

US Decisions Inc.

An Independent Review Organization
71 Court Street
Belfast, ME 04915
Phone: (512) 782-4560
Fax: (207) 470-1085
Email: manager@us-decisions.com

NOTICE OF INDEPENDENT REVIEW DECISION

DATE OF REVIEW:

Sep/04/2009

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Left Upper Extremity EMG/NCV study (CPT code 95860)

DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

MD, Board Certified in Physical Medicine and Rehabilitation
Board Certified in Pain Management
Board Certified in Electrodiagnostic Medicine

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Overturned (Disagree)

Partially Overturned (Agree in part/Disagree in part)

INFORMATION PROVIDED TO THE IRO FOR REVIEW

PATIENT CLINICAL HISTORY SUMMARY

This is a xx year old man with an injury date on xx/xx/xx. He had left shoulder and neck pain and elbow pain. Dr. evaluation in 6/09 reported neck and shoulder pain with dysesthesias. He found local tenderness in the cervical region and reduced cervical flexion and extension. He felt there was shoulder impingement. The cervical MRI in March 2008 reportedly showed C3/4 osteophytes, but no nerve root compression. The shoulder MRI showed some osteophytes with possible impingement. There is a 6/08 exam by Dr. describing pain in the radial epicondyle with elbow pain with forearm (wrist) pronation. Dr. wrote 6/24/09 that "It is medically necessary to conduct this EMG/NCV to rule out impingement of the shoulder and to rule out that the patient is not suffering from pain due to cervical degeneration."

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION

This request is for Left Upper Extremity EMG/NCV study (CPT code 95860). Only the EMG CPT code is provided.

Dr. has written that the request is necessary to differentiate a neurological problem vs.

cervical radiculopathy. The ODG recommends EMG studies when used to differentiate a radiculopathy from a nerve compression injury. The EMG for a cervical radiculopathy has a low sensitivity. There is only one physical examination provided of the elbow. This was the 6/08 exam. The provider's records indicate that the pain at the radial head could be explained by radial nerve compression in the proximal forearm (radial tunnel), which would necessitate both the EMG and NCV. Based on the records in this case, clinical judgment, and the ODG, the reviewer finds that medical necessity exists for Left Upper Extremity EMG/NCV study (CPT code 95860).

Electromyography (EMG)

Recommended (needle, not surface) as an option in selected cases. The American Association of Electrodiagnostic Medicine conducted a review on electrodiagnosis in relation to cervical radiculopathy and concluded that the test was moderately sensitive (50%-71%) and highly specific (65%-85%). (AAEM, 1999) EMG findings may not be predictive of surgical outcome in cervical surgery, and patients may still benefit from surgery even in the absence of EMG findings of nerve root impingement. This is in stark contrast to the lumbar spine where EMG findings have been shown to be highly correlative with symptoms

Positive diagnosis of radiculopathy: Requires the identification of neurogenic abnormalities in two or more muscles that share the same nerve root innervation but differ in their peripheral nerve supply.

Timing: Timing is important as nerve root compression will reflect as positive if active changes are occurring. Changes of denervation develop within the first to third week after compression (fibrillations and positive sharp waves develop first in the paraspinals at 7-10 days and in the limb muscles at 2-3 weeks), and reinnervation is found at about 3-6 months

Acute findings: Identification of fibrillation potentials in denervated muscles with normal motor unit action potentials (usually within 6 months of symptoms: may disappear within 6 weeks in the paraspinals and persist for up to 1-2 years in distal limbs)

Chronic findings: Findings of motor unit action potentials with increased duration and phases that represent reinnervation. With time these become broad, large and polyphasic and may persist for years

Anatomy: The test primarily evaluates ventral (anterior) root function (motor) and may be negative if there is dorsal root compression (sensory) only. Only C4-8 and T1 in the neck region have limb representation that can be tested electrodiagnostically. The anatomic basis for this lies in the fact that the cervical nerve roots have a motor and a sensory component. It is possible to impinge the sensory component with a herniated disc or bone spur and not affect the motor component. As a result, the patient may report radicular pain that correlates to the MRI without having EMG evidence of motor loss.

Paraspinal fibrillation potentials: May be seen in normal individuals and are nonspecific for etiology. The presence of these alone is insufficient to make a diagnosis of radiculopathy and they may be absent when there is a diagnosis of radiculopathy secondary to sampling error, timing, or because they were spared. They may support a diagnosis of radiculopathy when corresponding abnormalities are present in the limb muscles

Indications when particularly helpful: EMG may be helpful for patients with double crush phenomenon, in particular, when there is evidence of possible metabolic pathology such as neuropathy secondary to diabetes or thyroid disease, or evidence of peripheral compression such as carpal tunnel syndrome.

H-reflex: Technically difficult to perform in the upper extremity but can be derived from the median nerve. The test is not specific for etiology and may be difficult to obtain in obese patients or those older than 60 years of age.

(Negrin, 1991) (Alrawi, 2006) (Ashkan, 2002) (Nardin, 1999) (Tsao, 2007) See Discectomy-laminectomy-laminoplasty. (Surface EMG and F-wave tests are not very specific and therefore are not recommended. For more information on surface EMG, see the Low Back

Chapter.)

Nerve conduction studies (NCS)

Not recommended. There is minimal justification for performing nerve conduction studies when a patient is presumed to have symptoms on the basis of radiculopathy. (Utah, 2006) See also the Carpal Tunnel Syndrome Chapter for more details on NCS. Studies have not shown portable nerve conduction devices to be effective.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION

ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE

AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES

DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES

EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN

INTERQUAL CRITERIA

MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

MERCY CENTER CONSENSUS CONFERENCE GUIDELINES

MILLIMAN CARE GUIDELINES

ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR

TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS

TEXAS TACADA GUIDELINES

TMF SCREENING CRITERIA MANUAL

PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)

OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES
(PROVIDE A DESCRIPTION)