

US Decisions Inc.

An Independent Review Organization
71 Court Street
Belfast, ME 04915
Phone: (512) 782-4560
Fax: (207) 470-1085
Email: manager@us-decisions.com

NOTICE OF INDEPENDENT REVIEW DECISION

DATE OF REVIEW:

Sep/12/2009

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Chronic Pain Management Program x 10 Sessions

DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

MD, Board Certified in Physical Medicine and Rehabilitation
Board Certified in Pain Management

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Overturned (Disagree)

Partially Overturned (Agree in part/Disagree in part)

INFORMATION PROVIDED TO THE IRO FOR REVIEW

ODG Guidelines and Treatment Guidelines
Adverse Determination Letters, 7/20/09, 8/17/09
Patient Referral Form, 4/28/09
Preauth Request, 7/13/09, 8/10/09
, 7/13/09, 8/10/09
Interdisciplinary Pain Treatment Components
CPMP Design
Environmental Intervention, 7/17/09, 7/15/09
, 6/1/09
, DO, 6/30/09
, MD, 7/6/09
BME, 4/30/09
Addendum, 4/30/09
Cervical Spine Films, 9/14/06
MRI of Cervical Spine, 6/23/08
Letter from Patient, 12/8/08
Response, 8/24/09
Summary
, DO, 4/22/09

PATIENT CLINICAL HISTORY SUMMARY

This is a xx year old man injured on xx/xx/xx. He had a cervical fusion in December 1992. He did well and continued to work until about 3 years ago when he began having more neck pain. The EMG in 8/06 showed chronic left C6/7 and right C7/8 radiculopathy. The MRI in 2006 showed the intervertebral fusion between C6/7 and spondylosis. He apparently had no relief with cervical ESIs. He is on Norco 1 qid. He is on Cymbalta and Robaxin.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION

This patient had been doing well until his functional level dropped with increased pain in the past 3 years. His goal is increasing his work and improving the quality of his life. He functions but with more pain and taking more pain medications. Records indicate he is not abusing the pain medications. The ODG addresses concerns about programs more than 2 years post injury. The patient's letter of 12/8/08 demonstrated his motivation and plea for relief. His symptoms are in fact 3 years old and one should not use the 1992 date as the baseline. There are psychological and physical issues present. He has a good work history before and since his injury. The patient meets all the requirements for a CPMP, with only one negative predictor, that being the interval since the injury. However, his symptoms are 3 years old, not 17 years old. Most pain program treatment goals are to improve the probability of returning to work. He is working, but at a reduced level in the past 3 years. One study cited outcomes after delayed treatment that did not use return to work as the end point. It cited some improved response, but no other changes with and without treatment. Mr. and Dr. have addressed the criteria and psychological issues involved. Criteria 2 and 7 are specifically significant, although not all of the criteria discussed in section 7 are pertinent. Criteria 9 addresses not only the return to work, but also the need for the post treatment care. The reviewer finds that a 10 session trial would be reasonable in this atypical case. The reviewer finds that medical necessity exists for Chronic Pain Management Program x 10 Sessions.

Chronic pain programs (functional restoration programs)

Recommended where there is access to programs with proven successful outcomes (i.e., decreased pain and medication use, improved function and return to work, decreased utilization of the health care system), for patients with conditions that have resulted in "Delayed recovery." There should be evidence that a complete diagnostic assessment has been made, with a detailed treatment plan of how to address physiologic, psychological and sociologic components that are considered components of the patient's pain. Patients should show evidence of motivation to improve and return to work, and meet the patient selection criteria outlined below. While these programs are recommended (see criteria below), the research remains ongoing as to (1) what is considered the "gold-standard" content for treatment; (2) the group of patients that benefit most from this treatment; (3) the ideal timing of when to initiate treatment; (4) the intensity necessary for effective treatment; and (5) cost-effectiveness. It has been suggested that interdisciplinary/multidisciplinary care models for treatment of chronic pain may be the most effective way to treat this condition. (Flor, 1992) (Gallagher, 1999) (Guzman, 2001) (Gross, 2005) (Sullivan, 2005) (Dysvik, 2005) (Airaksinen, 2006) (Schonstein, 2003) (Sanders, 2005) (Patrick, 2004) (Buchner, 2006) These treatment modalities are based on the biopsychosocial model, one that views pain and disability in terms of the interaction between physiological, psychological and social factors. (Gatchel, 2005) See Biopsychosocial model of chronic pain....

Predictors of success and failure: As noted, one of the criticisms of interdisciplinary/multidisciplinary rehabilitation programs is the lack of an appropriate screening tool to help to determine who will most benefit from this treatment. Retrospective research has examined decreased rates of completion of functional restoration programs, and there is ongoing research to evaluate screening tools prior to entry. (Gatchel, 2006) There is need for research in terms of necessity and/or effectiveness of counseling for patients considered to be "at-risk" for post-discharge problems. (Proctor, 2004) The following variables have been found to be negative predictors of efficacy of treatment with the

programs as well as negative predictors of completion of the programs: (1) a negative relationship with the employer/supervisor; (2) poor work adjustment and satisfaction; (3) a negative outlook about future employment; (4) high levels of psychosocial distress (higher pretreatment levels of depression, pain and disability); (5) involvement in financial disability disputes; (6) greater rates of smoking; (7) increased duration of pre-referral disability time; (8) higher prevalence of opioid use; and (9) elevated pre-treatment levels of pain. (Linton, 2001) (Bendix, 1998) (McGeary, 2006) (McGeary, 2004) (Gatchel2, 2005) (Dersh, 2007)

Role of duration of disability: There is little research as to the success of return to work with functional restoration programs in long-term disabled patients (> 24 months).

Studies supporting programs for patients with long-term disability: Long-term disabled patients (at least 18 months) vs. short-term disabled (4 to 8 months) were evaluated using Pride data (1990-1993). No control was given for patients that did not undergo a program. During the time studied program dropouts averaged 8% to 12%. (It does appear that at the time of this study, participants in the program were detoxified from opioids prior to beginning.) The long-term disabled group was more likely to have undergone spinal surgery, with this likelihood increasing with time. Return to work was statistically different between the short-term disabled (93%) and the long-term disabled-18 months (80%). The long-term disabled-24 months group had a 75% return to work. Long-term disabled-18 month patients were statistically more likely to visit new health providers than short-term disabled patients (34% and 25% respectively). Work retention at one year in groups up to 24 months duration of disability was 80%. This dropped to 66% in the group that had been disabled for > 24 months. The percentage of recurrent lost time injury claims increased from around 1% in the groups disabled for < 35 months to 8.3% in the groups disabled for > 36 months. A main criterion for success appeared to be the decision of the patient to actively participate in the program rehabilitation goals. (Jordan, 1998)

Studies suggesting limited results in patients with long-term disability: While early studies have suggested that time out-of-work is a predictor of success for occupational outcomes, these studies have flaws when an attempt is made to apply them to chronic pain programs. (Gallagher, 1989) (Beals, 1972) (Krause, 1994) Washington State studied the role of duration of work injury on outcome using a statistical model that allowed for a comparison of patients that participated in a multidisciplinary pain program (using data from 1991-1993) vs. those that were evaluated and not treated. This was not an actual study of time of disability, but of duration of injury (mean years from injury to evaluation of 2.6 years for the treated group and 4.0 years for the evaluated only group). The original statistical analysis allowed for a patient to be included in a "treated group" for those individuals that both completed and did not complete the program. Data was collected from 10 sites. Each of the centers was CARF approved and included Pysch/behavioral treatment, vocation counseling and physical therapy. A sub-study evaluated a comparison of patients that were treatment completers vs. those that did not participate (78.6%, N=963). No information was given in terms of surgical procedures or medications. The primary outcome was time loss status of subjects 2 years after they had undergone the index pain center evaluation.

In the 2001 study, if chronicity of duration of injury was controlled for, there was no significant benefit produced in terms of patients that were receiving time-loss benefits at 2-years post treatment between the two groups. Approximately 60% of both groups were not receiving benefits at the two-year period. As noted, the "treated patient" was only guaranteed to have started a program. A repeat analysis of only the patients who completed the study did not significantly change the results of the study. In a 2004 survey follow-up no significant difference was found between treated and untreated groups, although the treated group had better response. The survey response was 50%, and the treatment responders were more likely to be disabled at the time of the survey. The authors suggest that the results indicated early intervention was a key to response of the programs, and that modest goals (improvement, not cure) be introduced. (Robinson, 2004) (Robinson, 2001) [The authors also concluded that there was no evidence that pain center treatment affects either disability status or clinical status of injured workers.]...

Criteria for the general use of multidisciplinary pain management programs

Outpatient pain rehabilitation programs may be considered medically necessary in the following circumstances

(1) The patient has a chronic pain syndrome, with evidence of loss of function that persists beyond three months and has evidence of three or more of the following: (a) Excessive dependence on health-care providers, spouse, or family; (b) Secondary physical deconditioning due to disuse and/or fear-avoidance of physical activity due to pain; (c) Withdrawal from social activities or normal contact with others, including work, recreation, or other social contacts; (d) Failure to restore preinjury function after a period of disability such that the physical capacity is insufficient to pursue work, family, or recreational needs; (e) Development of psychosocial sequelae that limits function or recovery after the initial incident, including anxiety, fear-avoidance, depression, sleep disorders, or nonorganic illness behaviors (with a reasonable probability to respond to treatment intervention); (f) The diagnosis is not primarily a personality disorder or psychological condition without a physical component; (g) There is evidence of continued use of prescription pain medications (particularly those that may result in tolerance, dependence or abuse) without evidence of improvement in pain or function

(2) Previous methods of treating chronic pain have been unsuccessful and there is an absence of other options likely to result in significant clinical improvement

(3) An adequate and thorough multidisciplinary evaluation has been made. This should include pertinent validated diagnostic testing that addresses the following: (a) A physical exam that rules out conditions that require treatment prior to initiating the program. All diagnostic procedures necessary to rule out treatable pathology, including imaging studies and invasive injections (used for diagnosis), should be completed prior to considering a patient a candidate for a program. The exception is diagnostic procedures that were repeatedly requested and not authorized. Although the primary emphasis is on the work-related injury, underlying non-work related pathology that contributes to pain and decreased function may need to be addressed and treated by a primary care physician prior to or coincident to starting treatment; (b) Evidence of a screening evaluation should be provided when addiction is present or strongly suspected; (c) Psychological testing using a validated instrument to identify pertinent areas that need to be addressed in the program (including but not limited to mood disorder, sleep disorder, relationship dysfunction, distorted beliefs about pain and disability, coping skills and/or locus of control regarding pain and medical care) or diagnoses that would better be addressed using other treatment should be performed; (d) An evaluation of social and vocational issues that require assessment

(4) If a goal of treatment is to prevent or avoid controversial or optional surgery, a trial of 10 visits (80 hours) may be implemented to assess whether surgery may be avoided.

(5) If a primary reason for treatment in the program is addressing possible substance use issues, an evaluation with an addiction clinician may be indicated upon entering the program to establish the most appropriate treatment approach (pain program vs. substance dependence program). This must address evaluation of drug abuse or diversion (and prescribing drugs in a non-therapeutic manner). In this particular case, once drug abuse or diversion issues are addressed, a 10-day trial may help to establish a diagnosis, and determine if the patient is not better suited for treatment in a substance dependence program. Addiction consultation can be incorporated into a pain program. If there is indication that substance dependence may be a problem, there should be evidence that the program has the capability to address this type of pathology prior to approval.

(6) Once the evaluation is completed, a treatment plan should be presented with specifics for treatment of identified problems, and outcomes that will be followed

(7) There should be documentation that the patient has motivation to change, and is willing to change their medication regimen (including decreasing or actually weaning substances

known for dependence). There should also be some documentation that the patient is aware that successful treatment may change compensation and/or other secondary gains. In questionable cases, an opportunity for a brief treatment trial may improve assessment of patient motivation and/or willingness to decrease habituating medications.

(8) Negative predictors of success (as outlined above) should be identified, and if present, the pre-program goals should indicate how these will be addressed

(9) If a program is planned for a patient that has been continuously disabled for greater than 24 months, the outcomes for the necessity of use should be clearly identified, as there is conflicting evidence that chronic pain programs provide return-to-work beyond this period. These other desirable types of outcomes include decreasing post-treatment care including medications, injections and surgery

(10) Treatment is not suggested for longer than 2 weeks without evidence of compliance and significant demonstrated efficacy as documented by subjective and objective gains. (Note: Patients may get worse before they get better. For example, objective gains may be moving joints that are stiff from lack of use, resulting in increased subjective pain.) However, it is also not suggested that a continuous course of treatment be interrupted at two weeks solely to document these gains, if there are preliminary indications that they are being made on a concurrent basis.

(11) Integrative summary reports that include treatment goals, compliance, progress assessment with objective measures and stage of treatment, must be made available upon request at least on a bi-weekly basis during the course of the treatment program

(12) Total treatment duration should generally not exceed 20 full-day (160 hours) sessions (or the equivalent in part-day sessions if required by part-time work, transportation, childcare, or comorbidities). (Sanders, 2005) Treatment duration in excess of 160 hours requires a clear rationale for the specified extension and reasonable goals to be achieved. Longer durations require individualized care plans explaining why improvements cannot be achieved without an extension as well as evidence of documented improved outcomes from the facility (particularly in terms of the specific outcomes that are to be addressed)

(13) At the conclusion and subsequently, neither re-enrollment in repetition of the same or similar rehabilitation program (e.g. work hardening, work conditioning, out-patient medical rehabilitation) is medically warranted for the same condition or injury (with possible exception for a medically necessary organized detox program). Prior to entry into a program the evaluation should clearly indicate the necessity for the type of program required, and providers should determine upfront which program their patients would benefit more from. A chronic pain program should not be considered a "stepping stone" after less intensive programs, but prior participation in a work conditioning or work hardening program does not preclude an opportunity for entering a chronic pain program if otherwise indicated

(14) Suggestions for treatment post-program should be well documented and provided to the referral physician. The patient may require time-limited, less intensive post-treatment with the program itself. Defined goals for these interventions and planned duration should be specified

(15) Post-treatment medication management is particularly important. Patients that have been identified as having substance abuse issues generally require some sort of continued addiction follow-up to avoid relapse.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION

- ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)