

# Wren Systems

An Independent Review Organization  
71 Court Street  
Belfast, ME 04915  
Phone: (512) 553-0533  
Fax: (207) 470-1064  
Email: manager@wrensystems.com

## NOTICE OF INDEPENDENT REVIEW DECISION

**DATE OF REVIEW:**

Sep/09/2009

**IRO CASE #:**

**DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:**

12 Sessions of Additional Lumbar Physical Therapy at 3 times a week for 4 weeks

**DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:**

MD, Board Certified in Physical Medicine and Rehabilitation  
Board Certified in Pain Management

**REVIEW OUTCOME:**

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Overturned (Disagree)

Partially Overturned (Agree in part/Disagree in part)

**INFORMATION PROVIDED TO THE IRO FOR REVIEW**

ODG Guidelines and Treatment Guidelines  
Adverse Determination Letters, 7/7/09, 7/21/09  
Letter to IRO from Pain & Recovery Clinic, 8/24/09  
Dr. Request for Reconsideration, 7/13/09  
PT Progress Note, 7/2/09  
Orthopedic Report, 7/3/09  
MD, 7/7/09  
DO, 4/13/09

**PATIENT CLINICAL HISTORY SUMMARY**

This is a male who developed low back pain going to the left lower extremity after a work related injury on xx/xx/xx. Apparently this was initially felt to be a lumbar strain. He had an MRI, but the only report is the description provided by Dr. in his DD exam. It reported facet hypertrophy and disc bulges at L3/4 and a protrusion at L4/5 with encroachment of the right L4 nerve root. His symptoms are on the left side. He had an ESI in June 2009 and, per Dr. had some symptom improvement. The prior reviewer noted there were 18 sessions of PT. Dr. requested an additional 12 sessions as the man reportedly needed to sit, stand, bend, squat, lift, push and pull. Dr. described reduced left ankle jerks, a positive left SLR and some sensory changes in the left L5/S1 dermatomes.

**ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS**

## **AND CONCLUSIONS USED TO SUPPORT THE DECISION**

The man's symptoms are on the left side, but the MRI suggested right sided L4 radicular compression. Dr. noted in June 2009 some signs of a left radiculopathy. The ODG recognizes the need for an aggressive therapy program targeting a home or self directed program. The ODG advises up to 10 therapy sessions for lumbar strains/sprains and radiculopathy. It will permit up to 2 additional sessions for instructions in a self directed program. The request for an additional 12 sessions in addition to the prior 18 exceeds the recommended treatment programs. Dr. did not provide a medical reason to vary from the ODG recommendations. Therefore, I can not justify the additional therapies. The reviewer finds that medical necessity does not exist for 12 Sessions of Additional Lumbar Physical Therapy at 3 times a week for 4 weeks.

### Physical therapy (PT)

Recommended. There is strong evidence that physical methods, including exercise and return to normal activities, have the best long-term outcome in employees with low back pain. See also Exercise. Direction from physical and occupational therapy providers can play a role in this, with the evidence supporting active therapy and not extensive use of passive modalities. The most effective strategy may be delivering individually designed exercise programs in a supervised format (for example, home exercises with regular therapist follow-up), encouraging adherence to achieve high dosage, and stretching and muscle-strengthening exercises seem to be the most effective types of exercises for treating chronic low back pain. (Hayden, 2005) Studies also suggest benefit from early use of aggressive physical therapy ("sports medicine model"), training in exercises for home use, and a functional restoration program, including intensive physical training, occupational therapy, and psychological support. (Zigenfus, 2000) (Linz, 2002) (Cherkin-NEJM, 1998) (Rainville, 2002) Successful outcomes depend on a functional restoration program, including intensive physical training, versus extensive use of passive modalities. (Mannion, 2001) (Jousset, 2004) (Rainville, 2004) (Airaksinen, 2006) One clinical trial found both effective, but chiropractic was slightly more favorable for acute back pain and physical therapy for chronic cases. (Skargren, 1998) A spinal stabilization program is more effective than standard physical therapy sessions, in which no exercises are prescribed. With regard to manual therapy, this approach may be the most common physical therapy modality for chronic low back disorder, and it may be appropriate as a pain reducing modality, but it should not be used as an isolated modality because it does not concomitantly reduce disability, handicap, or improve quality of life. (Goldby-Spine, 2006) Better symptom relief is achieved with directional preference exercise. (Long, 2004) As compared with no therapy, physical therapy (up to 20 sessions over 12 weeks) following disc herniation surgery was effective. Because of the limited benefits of physical therapy relative to "sham" therapy (massage), it is open to question whether this treatment acts primarily physiologically, but psychological factors may contribute substantially to the benefits observed. (Erdogmus, 2007) See also specific physical therapy modalities, as well as Exercise; Work conditioning; Lumbar extension exercise equipment; McKenzie method; Stretching; & Aquatic therapy. [Physical therapy is the treatment of a disease or injury by the use of therapeutic exercise and other interventions that focus on improving posture, locomotion, strength, endurance, balance, coordination, joint mobility, flexibility, activities of daily living and alleviating pain. (BlueCross BlueShield, 2005) As for visits with any medical provider, physical therapy treatment does not preclude an employee from being at work when not visiting the medical provider, although time off may be required for the visit.

Active Treatment versus Passive Modalities: The use of active treatment modalities instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with acute low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) The most commonly used active treatment modality is Therapeutic exercises (97110), but other active therapies may be recommended as well, including Neuromuscular reeducation (97112), Manual therapy (97140), and Therapeutic activities/exercises (97530).

A recent RCT comparing active spinal stabilization exercises (using the GDS or Godelive Denys-Struyf method) with passive electrotherapy using TENS plus microwave treatment (considered conventional physical therapy in Spanish primary care), concluded that treatment of nonspecific LBP using the GDS method provides greater improvements in the midterm (6 months) in terms of pain, functional ability, and quality of life. (Arribas, 2009)

Patient Selection Criteria: Multiple studies have shown that patients with a high level of fear-avoidance do much better in a supervised physical therapy exercise program, and patients with low fear-avoidance do better following a self-directed exercise program. When using the Fear-Avoidance Beliefs Questionnaire (FABQ), scores greater than 34 predicted success with PT supervised care. (Fritz, 2001) (Fritz, 2002) (George, 2003) (Kluber, 2004) (Riipinen, 2005) (Hicks, 2005) Without proper patient selection, routine physical therapy may be no more effective than one session of assessment and advice from a physical therapist. (Frost, 2004) Patients exhibiting the centralization phenomenon during lumbar range of motion testing should be treated with the specific exercises (flexion or extension) that promote centralization of symptoms. When findings from the patient's history or physical examination are associated with clinical instability, they should be treated with a trunk strengthening and stabilization exercise program. (Fritz-Spine, 2003) Practitioners must be cautious when implementing the wait-and-see approach for LBP, and once medical clearance has been obtained, patients should be advised to keep as active as possible. Patients presenting with high fear avoidance characteristics should have these concerns addressed aggressively to prevent long-term disability, and they should be encouraged to promote the resumption of physical activity. (Hanney, 2009)

Post Epidural Steroid Injections: ESIs are currently recommended as a possible option for short-term treatment of radicular pain (sciatica), defined as pain in dermatomal distribution with corroborative findings of radiculopathy. The general goal of physical therapy during the acute/subacute phase of injury is to decrease guarding, maintain motion, and decrease pain and inflammation. Progression of rehabilitation to a more advanced program of stabilization occurs in the maintenance phase once pain is controlled. There is little evidence-based research that addresses the use of physical therapy post ESIs, but it appears that most randomized controlled trials have utilized an ongoing, home directed program post injection. Based on current literature, the only need for further physical therapy treatment post ESI would be to emphasize the home exercise program, and this requirement would generally be included in the currently suggested maximum visits for the underlying condition, or at least not require more than 2 additional visits to reinforce the home exercise program. ESIs have been found to have limited effectiveness for treatment of chronic pain. The claimant should continue to follow a home exercise program post injection. (Luijsterburg, 2007) (Luijsterburg2, 2007) (Price, 2005) (Vad, 2002) (Smeal, 2004)

#### ODG Physical Therapy Guidelines –

Allow for fading of treatment frequency (from up to 3 or more visits per week to 1 or less), plus active self-directed home PT. Also see other general guidelines that apply to all conditions under Physical Therapy in the ODG Preface, including assessment after a "six-visit clinical trial"

Lumbar sprains and strains (ICD9 847.2)

10 visits over 8 weeks

Sprains and strains of unspecified parts of back (ICD9 847)

10 visits over 5 weeks

Sprains and strains of sacroiliac region (ICD9 846)

Medical treatment: 10 visits over 8 weeks

Lumbago; Backache, unspecified (ICD9 724.2; 724.5)

9 visits over 8 weeks

Intervertebral disc disorders without myelopathy (ICD9 722.1; 722.2; 722.5; 722.6; 722.8)

Medical treatment: 10 visits over 8 weeks

Post-injection treatment: 1-2 visits over 1 week

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION**

ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE

AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES

DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES

EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN

INTERQUAL CRITERIA

MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

MERCY CENTER CONSENSUS CONFERENCE GUIDELINES

MILLIMAN CARE GUIDELINES

ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR

TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS

TEXAS TACADA GUIDELINES

TMF SCREENING CRITERIA MANUAL

PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)

OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)