



5068 West Plano Parkway Suite 122  
 Plano, Texas 75093  
 Phone: (972) 931-5100

**DATE OF REVIEW:** 01/27/2010

**IRO CASE #:**

**DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:**

IRO - L5-S1 Transforaminal Epidural Steroid Injection, Fluoroscopically guided, IV Conscious Sedation

**A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:**

This case was reviewed by a Texas licensed MD, specializing in Pain Management, Anesthesiology. The physician advisor has the following additional qualifications, if applicable:

ABMS Anesthesiology: Pain Medicine, Anesthesiology

**REVIEW OUTCOME:**

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld

Health Care Service(s) in Dispute	CPT Codes	Date of Service(s)	Outcome of Independent Review
IRO - L5-S1 Transforaminal Epidural Steroid Injection, Fluoroscopically guided, IV Conscious Sedation		-	Upheld

**INFORMATION PROVIDED TO THE IRO FOR REVIEW:**

No	Document Type	Provider or Sender	Page Count	Service Start Date	Service End Date
1	IRO Request	Texas Department of Insurance	15	01/08/2010	01/08/2010
2	IRO Carrier/URA Records	HDI; Health Direct Inc.	15	03/18/2009	01/08/2010

**PATIENT CLINICAL HISTORY [SUMMARY]:**

The patient is a male with a date of injury of xx/xx/xx. The mechanism of injury was due to lifting a heavy object. There was a diagnosis of low back pain. An MRI was performed on 03-15-09, which revealed an L3-

4 and L4-5 anular tear. There was an L5-S1 spondylolisthesis causing moderate foraminal stenosis. Physical therapy and medications were utilized without reported benefit. The patient was seen by Dr. and there was noted to be a positive straight leg raise on physical exam. There was no documentation of a neurologic exam.

**ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION:**

The ODG Guides will support the use of epidural steroid injections for the treatment of radiculopathy. Radiculopathy is defined as significant alteration in the function of a nerve root or nerve roots and is usually caused by pressure on one or several nerve roots. The diagnosis requires a dermatomal distribution of pain, numbness, and/or paresthesias in a dermatomal distribution. A root tension sign is usually positive. The diagnosis of herniated disk must be substantiated by an appropriate finding on an imaging study. The presence of findings on an imaging study in and of itself does not make the diagnosis of radiculopathy. In this patient, there was no documentation of a dermatomal distribution of the pain. There was also no neurologic exam indicating levels of abnormal findings. Based on this, the request is not supported by the ODG guides and is not considered medically necessary.

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:**

ODG; Low Back Chapter

Recommended as a possible option for short-term treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy) with use in conjunction with active rehab efforts. See specific criteria for use below. Radiculopathy symptoms are generally due to herniated nucleus pulposus or spinal stenosis, although ESIs have not been found to be as beneficial a treatment for the latter condition.

*Short-term symptoms:* The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months. ([Armon, 2007](#)) Epidural steroid injection can offer short-term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. There is little information on improved function or return to work. There is no high-level evidence to support the use of epidural injections of steroids, local anesthetics, and/or opioids as a treatment for acute low back pain without radiculopathy. ([Benzon, 1986](#)) ([ISIS, 1999](#)) ([DePalma, 2005](#)) ([Molloy, 2005](#)) ([Wilson-MacDonald, 2005](#)) This recent RCT concluded that both ESIs and PT seem to be effective for lumbar spinal stenosis for up to 6 months. Both ESI and PT groups demonstrated significant improvement in pain and functional parameters compared to control and no significant difference was noted between the 2 treatment groups at 6 months, but the ESI group was significantly more improved at the 2nd week. ([Koc, 2009](#))

*Use for chronic pain:* Chronic duration of symptoms (> 6 months) has also been found to decrease success rates with a threefold decrease found in patients with symptom duration > 24 months. The ideal time of either when to initiate treatment or when treatment is no longer thought to be effective has not been determined. ([Hopwood, 1993](#)) ([Cyteval, 2006](#)) Indications for repeating ESIs in patients with chronic pain at a level previously injected (> 24 months) include a symptom-free interval or indication of a new clinical presentation at the level.

*Transforaminal approach:* Some groups suggest that there may be a preference for a transforaminal approach as the technique allows for delivery of medication at the target tissue site, and an advantage for transforaminal injections in herniated nucleus pulposus over translaminar or caudal injections has been suggested in the best available studies. ([Riew, 2000](#)) ([Vad, 2002](#)) ([Young, 2007](#)) This approach may be

particularly helpful in patients with large disc herniations, foraminal stenosis, and lateral disc herniations. ([Colorado, 2001](#)) ([ICSI, 2004](#)) ([McLain, 2005](#)) ([Wilson-MacDonald, 2005](#))

*Fluoroscopic guidance:* Fluoroscopic guidance with use of contrast is recommended for all approaches as needle misplacement may be a cause of treatment failure. ([Manchikanti, 1999](#)) ([Colorado, 2001](#)) ([ICSI, 2004](#)) ([Molloy, 2005](#)) ([Young, 2007](#))

*Factors that decrease success:* Decreased success rates have been found in patients who are unemployed due to pain, who smoke, have had previous back surgery, have pain that is not decreased by medication, and/or evidence of substance abuse, disability or litigation. ([Jamison, 1991](#)) ([Abram, 1999](#)) Research reporting effectiveness of ESIs in the past has been contradictory, but these discrepancies are felt to have been, in part, secondary to numerous methodological flaws in the early studies, including the lack of imaging and contrast administration. Success rates also may depend on the technical skill of the interventionalist. ([Carette, 1997](#)) ([Bigos, 1999](#)) ([Rozenberg, 1999](#)) ([Botwin, 2002](#)) ([Manchikanti, 2003](#)) ([CMS, 2004](#)) ([Delpont, 2004](#)) ([Khot, 2004](#)) ([Buttermann, 2004](#)) ([Buttermann2, 2004](#)) ([Samanta, 2004](#)) ([Cigna, 2004](#)) ([Benzon, 2005](#)) ([Dashfield, 2005](#)) ([Arden, 2005](#)) ([Price, 2005](#)) ([Resnick, 2005](#)) ([Abdi, 2007](#)) ([Boswell, 2007](#)) ([Buenaventura, 2009](#)) Also see [Epidural steroid injections, "series of three"](#) and [Epidural steroid injections, diagnostic](#). ESIs may be helpful with radicular symptoms not responsive to 2 to 6 weeks of conservative therapy. ([Kinkade, 2007](#)) Epidural steroid injections are an option for short-term pain relief of persistent radiculopathy, although not for nonspecific low back pain or spinal stenosis. ([Chou, 2008](#)) As noted above, injections are recommended if they can facilitate a return to functionality (via activity & exercise). If post-injection physical therapy visits are required for instruction in these active self-performed exercise programs, these visits should be included within the overall recommendations under [Physical therapy](#), or at least not require more than 2 additional visits to reinforce the home exercise program.

*With discectomy:* Epidural steroid administration during lumbar discectomy may reduce early neurologic impairment, pain, and convalescence and enhance recovery without increasing risks of complications. ([Rasmussen, 2008](#))

An updated Cochrane review of injection therapies (ESIs, facets, trigger points) for low back pain concluded that there is no strong evidence for or against the use of any type of injection therapy, but it cannot be ruled out that specific subgroups of patients may respond to a specific type of injection therapy. ([Staal-Cochrane, 2009](#)) Recent studies document a 629% increase in expenditures for ESIs, without demonstrated improvements in patient outcomes or disability rates. ([Devo, 2009](#)) There is fair evidence that epidural steroid injection is moderately effective for short-term (but not long-term) symptom relief. ([Chou3, 2009](#))

#### **Criteria for the use of Epidural steroid injections:**

*Note: The purpose of ESI is to reduce pain and inflammation, thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit.*

(1) Radiculopathy must be documented. Objective findings on examination need to be present. For unequivocal evidence of radiculopathy, see AMA Guides, 5th Edition, page 382-383. ([Andersson, 2000](#))

(2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants).

(3) Injections should be performed using fluoroscopy (live x-ray) and injection of contrast for guidance.

(4) *Diagnostic Phase:* At the time of initial use of an ESI (formally referred to as the "diagnostic phase" as initial injections indicate whether success will be obtained with this treatment intervention), a maximum of one to two injections should be performed. A repeat block is not recommended if there is inadequate response to the first block (< 30% is a standard placebo response). A second block is also not indicated if the first block is accurately placed unless: (a) there is a question of the pain generator; (b) there was possibility of inaccurate placement; or (c) there is evidence of multilevel pathology. In these cases a different level or approach might be proposed. There should be an interval of at least one to two weeks between injections.

(5) No more than two nerve root levels should be injected using transforaminal blocks.

(6) No more than one interlaminar level should be injected at one session.

(7) *Therapeutic phase:* If after the initial block/blocks are given (see “Diagnostic Phase” above) and found to produce pain relief of at least 50-70% pain relief for at least 6-8 weeks, additional blocks may be required. This is generally referred to as the “therapeutic phase.” Indications for repeat blocks include acute exacerbation of pain, or new onset of symptoms. The general consensus recommendation is for no more than 4 blocks per region per year. ([CMS, 2004](#)) ([Boswell, 2007](#))

(8) Repeat injections should be based on continued objective documented pain relief, decreased need for pain medications, and functional response.

(9) Current research does not support a routine use of a “series-of-three” injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections for the initial phase and rarely more than 2 for therapeutic treatment.

(10) It is currently not recommended to perform epidural blocks on the same day of treatment as facet blocks or sacroiliac blocks or lumbar sympathetic blocks or trigger point injections as this may lead to improper diagnosis or unnecessary treatment.

(11) Cervical and lumbar epidural steroid injection should not be performed on the same day. (Doing both injections on the same day could result in an excessive dose of steroids, which can be dangerous, and not worth the risk for a treatment that has no long-term benefit.)

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE**
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES**
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES**
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN**
- INTERQUAL CRITERIA**
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS**
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES**
- MILLIMAN CARE GUIDELINES**
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES**
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR**
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS**
- TEXAS TACADA GUIDELINES**
- TMF SCREENING CRITERIA MANUAL**
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)**
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)**

**TEXAS DEPARTMENT OF INSURANCE COMPLAINT PROCESS:** The Texas Department of Insurance requires Independent Review Organizations to be licensed to perform Independent Review in Texas. To contact the Texas Department of Insurance regarding any complaint, you may call or write the Texas Department of Insurance. The telephone number is 1-800-578-4677 or in writing at: Texas Department of Insurance, PO Box 149104 Austin TX, 78714. In accordance with Rule 102.4(h), a copy of this Independent Review Organization (IRO) Decision was sent to the carrier, the requestor and claimant via facsimile or U.S. Postal Service from the office of the IRO on 01/27/2010.