

# Wren Systems

An Independent Review Organization  
3112 Windsor Road #A Suite 376  
Austin, TX 78703  
Phone: (512) 553-0533  
Fax: (207) 470-1064  
Email: manager@wrensystems.com

## NOTICE OF INDEPENDENT REVIEW DECISION

**DATE OF REVIEW:** Feb/09/2010

**IRO CASE #:**

**DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:**

Lumbar ESI at L5-S1 under fluoroscopy

**DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:**

MD, Board Certified in Physical Medicine and Rehabilitation  
Board Certified in Pain Management  
Board Certified in Electrodiagnostic Medicine

**REVIEW OUTCOME:**

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)  
 Overturned (Disagree)  
 Partially Overturned (Agree in part/Disagree in part)

**INFORMATION PROVIDED TO THE IRO FOR REVIEW**

Adverse Determination Letters, 1/13/10, 12/30/09  
Surgery Center 12/14/09  
Surgeons Associates 10/22/09, 10/28/09, 10/14/09, 11/9/09,  
11/5/09  
Masters 12/16/09  
ODG Guidelines and Treatment Guidelines, Low Back

**PATIENT CLINICAL HISTORY SUMMARY**

This woman was injured in a fall from a stepladder when she landed on her buttocks. The date of injury is xx/xx/xx. She has back pain going to the buttocks. She had an MRI that showed a disc protrusion at L5/S1 compromising the left S1 nerve root. She failed to improve with PT. An ESI at L5/S1 was performed on 12/14/09 and reportedly provided 50% relief. A second was requested. Dr. wrote her neurological exam was intact. Dr. did not provide a neurological examination.

**ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION**

The records presented did not clearly identify a specific dermatome as required in the ODG and in the AMA Guides. Further, there was no described neurological loss. The presence of radiological findings alone do not substantiate a radiculopathy per the ODG and AMA Guides. There is no justification for repeating a second diagnostic block. A repeat (second)

therapeutic block is permissible if there is 50% relief for 6-8 weeks. The records presented did not contain evidence after the 12/14/09 injection to confirm the effectiveness. For these reasons, the request for a repeat injection does not meet the ODG criteria at this time. The reviewer finds that medical necessity does not exist at this time for Lumbar ESI at L5-S1 under fluoroscopy.

## Epidural steroid injections (ESIs), therapeutic

Recommended as a possible option for short-term treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy) with use in conjunction with active rehab efforts. See specific criteria for use below. Radiculopathy symptoms are generally due to herniated nucleus pulposus or spinal stenosis, although ESIs have not been found to be as beneficial a treatment for the latter condition

Short-term symptoms: The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months. (Armon, 2007) Epidural steroid injection can offer short-term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. There is little information on improved function or return to work. There is no high-level evidence to support the use of epidural injections of steroids, local anesthetics, and/or opioids as a treatment for acute low back pain without radiculopathy. (Benzon, 1986) (ISIS, 1999) (DePalma, 2005) (Molloy, 2005) (Wilson-MacDonald, 2005) This recent RCT concluded that both ESIs and PT seem to be effective for lumbar spinal stenosis for up to 6 months. Both ESI and PT groups demonstrated significant improvement in pain and functional parameters compared to control and no significant difference was noted between the 2 treatment groups at 6 months, but the ESI group was significantly more improved at the 2nd week. (Koc, 2009)...

ESIs may be helpful with radicular symptoms not responsive to 2 to 6 weeks of conservative therapy. (Kinkade, 2007) Epidural steroid injections are an option for short-term pain relief of persistent radiculopathy, although not for nonspecific low back pain or spinal stenosis. (Chou, 2008) As noted above, injections are recommended if they can facilitate a return to functionality (via activity & exercise). If post-injection physical therapy visits are required for instruction in these active self-performed exercise programs, these visits should be included within the overall recommendations under Physical therapy, or at least not require more than 2 additional visits to reinforce the home exercise program....

### Criteria for the use of Epidural steroid injections

Note: The purpose of ESI is to reduce pain and inflammation, thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit

(1) Radiculopathy must be documented. Objective findings on examination need to be present. For unequivocal evidence of radiculopathy, see AMA Guides, 5th Edition, page 382-383. (Andersson, 2000)

(2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants)

(3) Injections should be performed using fluoroscopy (live x-ray) and injection of contrast for guidance

(4) Diagnostic Phase: At the time of initial use of an ESI (formally referred to as the “diagnostic phase” as initial injections indicate whether success will be obtained with this treatment intervention), a maximum of one to two injections should be performed. A repeat block is not recommended if there is inadequate response to the first block (< 30% is a standard placebo response). A second block is also not indicated if the first block is accurately placed unless: (a) there is a question of the pain generator; (b) there was possibility of inaccurate placement; or (c) there is evidence of multilevel pathology. In these cases a different level or approach might be proposed. There should be an interval of at least one to two weeks between injections

(5) No more than two nerve root levels should be injected using transforaminal blocks

(6) No more than one interlaminar level should be injected at one session

(7) Therapeutic phase: If after the initial block/blocks are given (see “Diagnostic Phase” above) and found to produce pain relief of at least 50-70% pain relief for at least 6-8 weeks, additional blocks may be required. This is generally referred to as the “therapeutic phase.” Indications for repeat blocks include acute exacerbation of pain, or new onset of symptoms. The general consensus recommendation is for no more than 4 blocks per region per year. (CMS, 2004) (Boswell, 2007)

(8) Repeat injections should be based on continued objective documented pain relief, decreased need for pain medications, and functional response

(9) Current research does not support a routine use of a “series-of-three” injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections for the initial phase and rarely more than 2 for therapeutic treatment

(10) It is currently not recommended to perform epidural blocks on the same day of treatment as facet blocks or sacroiliac blocks or lumbar sympathetic blocks or trigger point injections as this may lead to improper diagnosis or unnecessary treatment

(11) Cervical and lumbar epidural steroid injection should not be performed on the same day. (Doing both injections on the same day could result in an excessive dose of steroids, which can be dangerous, and not worth the risk for a treatment that has no long-term benefit.)

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION**

ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE

AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES

DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES

EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN

INTERQUAL CRITERIA

MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

MERCY CENTER CONSENSUS CONFERENCE GUIDELINES

MILLIMAN CARE GUIDELINES

ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR

TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS

TEXAS TACADA GUIDELINES

TMF SCREENING CRITERIA MANUAL

PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)

OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)

AMA Guides to the Evaluation of Permanent Impairment, 5th edition