

C-IRO Inc.

An Independent Review Organization
7301 RANCH RD 620 N, STE 155-199A
Austin, TX 78726
Phone: (512) 772-4390
Fax: (512) 519-7098
Email: resolutions.manager@ciro-site.com

NOTICE OF INDEPENDENT REVIEW DECISION

DATE OF REVIEW:

Jan/16/2010

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

6 sessions of physical therapy for the lumbar spine and cervical spine

DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

MD, Board Certified in Physical Medicine and Rehabilitation

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Overturned (Disagree)

Partially Overturned (Agree in part/Disagree in part)

INFORMATION PROVIDED TO THE IRO FOR REVIEW

ODG Guidelines and Treatment Guidelines
Adverse Determination Letters, 12/7/09, 12/22/09
Inc., 12/11/09, 11/25/09
M.D. 11/12/09
HDI, 12/7/09
Back and Neck Clinic, 7/27/06, 6/30/06
Dr. D.C. 6/30/06

PATIENT CLINICAL HISTORY SUMMARY

This man reportedly was injured on x/xx/xx when boxes fell on his head and neck. He developed neck and low back pain. He had 8 sessions of PT and chiropractic care in 2006 with Dr. He subsequently was incarcerated and was not treated for three years. He currently describes pain in his neck and low back going to the shoulders and legs, plus numbness in his hands. Dr. noted on his examination the presence of local cervical and lumbar tenderness, normal reflexes, sensation and strength. He has positive bilateral SLR.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDING CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION

This patient's problem is now more than 3 years old. No radiological reports were provided for this review. The ODG permits up to 9-10 therapy sessions for neck pain and lumbago (low back pain) attributed to no other cause. The claimant already had this treatment 3 years ago. There was nothing provided in the records to establish why a new program should be instituted at this time. The request does not conform to the guidelines. The reviewer finds that medical necessity does not exist for 6 sessions of physical therapy for the lumbar spine and cervical spine.

Physical therapy (PT) (cervical)

Recommended. Low stress aerobic activities and stretching exercises can be initiated at home and supported by a physical therapy provider, to avoid debilitation and further restriction of motion. (Rosenfeld, 2000) (Bigos, 1999) For mechanical disorders for the neck, therapeutic exercises have demonstrated clinically significant benefits in terms of pain, functional restoration, and patient global assessment scales. (Philadelphia, 2001) (Colorado, 2001) (Kjellman, 1999) (Seferiadis, 2004) Physical therapy seems to be more effective than general practitioner care on cervical range of motion at short-term follow-up. (Scholten-Peeters, 2006) In a recent high quality study, mobilization appears to be one of the most effective non-invasive interventions for the treatment of both pain and cervical range of motion in the acutely injured WAD patient. (Conlin, 2005) A recent high quality study found little difference among conservative whiplash therapies, with some advantage to an active mobilization program with physical therapy twice weekly for 3 weeks. (Kongsted, 2007) See also specific physical therapy modalities, as well as Exercise

ODG Physical Therapy Guidelines –

Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home PT. Also see other general guidelines that apply to all conditions under Physical Therapy in the ODG Preface, including assessment after a "six-visit clinical trial"

Cervicalgia (neck pain); Cervical spondylosis (ICD9 723.1; 721.0)

9 visits over 8 weeks

Sprains and strains of neck (ICD9 847.0)

10 visits over 8 weeks

Physical therapy (PT) (lumbar)

Recommended. There is strong evidence that physical methods, including exercise and return to normal activities, have the best long-term outcome in employees with low back pain. See also Exercise. Direction from physical and occupational therapy providers can play a role in this, with the evidence supporting active therapy and not extensive use of passive modalities. The most effective strategy may be delivering individually designed exercise programs in a supervised format (for example, home exercises with regular therapist follow-up), encouraging adherence to achieve high dosage, and stretching and muscle-strengthening exercises seem to be the most effective types of exercises for treating chronic low back pain. (Hayden, 2005) Studies also suggest benefit from early use of aggressive physical therapy ("sports medicine model"), training in exercises for home use, and a functional restoration program, including intensive physical training, occupational therapy, and psychological support. (Zigenfus, 2000) (Linz, 2002) (Cherkin-NEJM, 1998) (Rainville, 2002) Successful outcomes depend on a functional restoration program, including intensive physical training, versus extensive use of passive modalities. (Mannion, 2001) (Jousset, 2004) (Rainville, 2004) (Airaksinen, 2006) One clinical trial found both effective, but chiropractic was slightly more favorable for acute back pain and physical therapy for chronic cases. (Skargren, 1998)

A spinal stabilization program is more effective than standard physical therapy sessions, in which no exercises are prescribed. With regard to manual therapy, this approach may be the most common physical therapy modality for chronic low back disorder, and it may be appropriate as a pain reducing modality, but it should not be used as an isolated modality because it does not concomitantly reduce disability, handicap, or improve quality of life. (Goldby-Spine, 2006) Better symptom relief is achieved with directional preference exercise. (Long, 2004) As compared with no therapy, physical therapy (up to 20 sessions over 12 weeks) following disc herniation surgery was effective. Because of the limited benefits of physical therapy relative to "sham" therapy (massage), it is open to question whether this

treatment acts primarily physiologically, but psychological factors may contribute substantially to the benefits observed. (Erdogmus, 2007) See also specific physical therapy modalities, as well as Exercise; Work conditioning; Lumbar extension exercise equipment; McKenzie method; Stretching; & Aquatic therapy. [Physical therapy is the treatment of a disease or injury by the use of therapeutic exercise and other interventions that focus on improving posture, locomotion, strength, endurance, balance, coordination, joint mobility, flexibility, activities of daily living and alleviating pain. (BlueCross BlueShield, 2005) As for visits with any medical provider, physical therapy treatment does not preclude an employee from being at work when not visiting the medical provider, although time off may be required for the visit.

Active Treatment versus Passive Modalities: The use of active treatment modalities instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with acute low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) The most commonly used active treatment modality is Therapeutic exercises (97110), but other active therapies may be recommended as well, including Neuromuscular reeducation (97112), Manual therapy (97140), and Therapeutic activities/exercises (97530). A recent RCT comparing active spinal stabilization exercises (using the GDS or Godelive Denys-Struyf method) with passive electrotherapy using TENS plus microwave treatment (considered conventional physical therapy in Spanish primary care), concluded that treatment of nonspecific LBP using the GDS method provides greater improvements in the midterm (6 months) in terms of pain, functional ability, and quality of life. (Arribas, 2009)

Patient Selection Criteria: Multiple studies have shown that patients with a high level of fear-avoidance do much better in a supervised physical therapy exercise program, and patients with low fear-avoidance do better following a self-directed exercise program. When using the Fear-Avoidance Beliefs Questionnaire (FABQ), scores greater than 34 predicted success with PT supervised care. (Fritz, 2001) (Fritz, 2002) (George, 2003) (Klaber, 2004) (Riipinen, 2005) (Hicks, 2005) Without proper patient selection, routine physical therapy may be no more effective than one session of assessment and advice from a physical therapist. (Frost, 2004) Patients exhibiting the centralization phenomenon during lumbar range of motion testing should be treated with the specific exercises (flexion or extension) that promote centralization of symptoms. When findings from the patient's history or physical examination are associated with clinical instability, they should be treated with a trunk strengthening and stabilization exercise program. (Fritz-Spine, 2003) Practitioners must be cautious when implementing the wait-and-see approach for LBP, and once medical clearance has been obtained, patients should be advised to keep as active as possible. Patients presenting with high fear avoidance characteristics should have these concerns addressed aggressively to prevent long-term disability, and they should be encouraged to promote the resumption of physical activity. (Hanney, 2009)

Post Epidural Steroid Injections: ESIs are currently recommended as a possible option for short-term treatment of radicular pain (sciatica), defined as pain in dermatomal distribution with corroborative findings of radiculopathy. The general goal of physical therapy during the acute/subacute phase of injury is to decrease guarding, maintain motion, and decrease pain and inflammation. Progression of rehabilitation to a more advanced program of stabilization occurs in the maintenance phase once pain is controlled. There is little evidence-based research that addresses the use of physical therapy post ESIs, but it appears that most randomized controlled trials have utilized an ongoing, home directed program post injection. Based on current literature, the only need for further physical therapy treatment post ESI would be to emphasize the home exercise program, and this requirement would generally be included in the currently suggested maximum visits for the underlying condition, or at least not require more than 2 additional visits to reinforce the home exercise program. ESIs have been found to have limited effectiveness for treatment of chronic pain. The claimant should continue to follow a home exercise program post injection. (Luijsterburg, 2007) (Luijsterburg2, 2007) (Price, 2005) (Vad, 2002) (Smeal, 2004)

Post-surgical (discectomy) rehab: A recent Cochrane review concluded that exercise programs starting 4-6 weeks post-surgery seem to lead to a faster decrease in pain and disability than no treatment; high intensity exercise programs seem to lead to a faster decrease in pain and disability than low intensity programs; home exercises are as good as supervised exercises; and active programs do not increase the re-operation rate. Although it is not harmful to return to activity after lumbar disc surgery, it is still unclear what exact components should be included in rehabilitation programs. High intensity programs seem to be more effective but they could also be more expensive. Another question is whether all patients should be treated post-surgery or is a minimal intervention with the message return to an active lifestyle sufficient, with only patients that still have symptoms 4 to 6 weeks post-surgery requiring rehabilitation programs. (Ostelo, 2009)

ODG Physical Therapy Guidelines –

Allow for fading of treatment frequency (from up to 3 or more visits per week to 1 or less), plus active self-directed home PT. Also see other general guidelines that apply to all conditions under Physical Therapy in the ODG Preface, including assessment after a "six-visit clinical trial"

Lumbar sprains and strains (ICD9 847.2)

10 visits over 8 week

Sprains and strains of unspecified parts of back (ICD9 847)

10 visits over 5 week

Lumbago; Backache, unspecified (ICD9 724.2; 724.5)

9 visits over 8 weeks

Physical medicine treatment (PAIN)

Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of acute pain treatment or acute exacerbations of chronic pain and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007)

ODG Physical Therapy Guidelines –

Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active

self-directed home PT. Also see other general guidelines that apply to all conditions under Physical Therapy in the ODG Preface

Myalgia and myositis, unspecified (ICD9 729.1)

9-10 visits over 8 weeks

Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2)

8-10 visits over 4 weeks

Reflex sympathetic dystrophy (CRPS) (ICD9 337.2)

26 visits over 16 weeks

Arthritis (ICD9 715)

Medical treatment: 9 visits over 8 weeks

Post-injection treatment: 1-2 visits over 1 week

Post-surgical treatment (see body-part chapters): 18 visits over 12 weeks

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION

ACOEM-AMERICA COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGBASE

AHCPR-AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES

DWC-DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES

EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN

INTERQUAL CRITERIA

MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

MERCY CENTER CONSENSUS CONFERENCE GUIDELINES

MILLIMAN CARE GUIDELINES

ODG-OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR

TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS

TEXAS TACADA GUIDELINES

TMF SCREENING CRITERIA MANUAL

PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)

OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)