

Notice of Independent Review Decision

**February 27, 2013**

**IRO CASE #:**

**DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:**

Lumbar Discogram with CT Scan

**A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:**

The physician performing this review is Board Certified, American Board of Orthopedic Surgery. The physician has been in practice since 1998 and is licensed in Texas, Oklahoma, Minnesota and South Dakota.

**REVIEW OUTCOME:**

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)  
 Overturned (Disagree)  
 Partially Overturned (Agree in part/Disagree in part)

Provide a description of the review outcome that clearly states whether medical necessity exists for each of the health care services in dispute.

*Upon independent review, the reviewer finds that the previous adverse determination should be **Upheld***

**INFORMATION PROVIDED TO THE IRO FOR REVIEW:**

**PATIENT CLINICAL HISTORY [SUMMARY]:**

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

Xx is a female who was injured on xx/xx/xx lifting a heavy patient while working for xx. She had an onset of low back pain and bilateral hip and leg pain. She has had physical therapy, interventional pain management, and medication management and continues to have similar symptoms. Her imaging studies were relatively unremarkable with the exception of what would appear to be mild broad-based disk bulge at each of the three lower lumbar levels.

## **ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS, AND CONCLUSIONS USED TO SUPPORT THE DECISION:**

ODG guidelines are very clear in the use of discography. While it is allowed under very strict criteria, this patient does not appear to meet those with the exception of the length of her disability and symptoms and her failure to achieve pain relief with nonoperative measures. This does not, however, indicate she is an operative candidate.

ODG guidelines for discography can allow the procedure if the patient has had satisfactory results from a detailed psychosocial assessment. This does not appear to have been performed at this time. The criteria for discography also include that the patient meets all criteria for spinal fusion. Again, the ODG guidelines for a lumbar spinal fusion include psychosocial screening examination, which again appears not to have been undertaken in this patient's particular case.

## **ODG -TWC**

*ODG Treatment*

*Integrated Treatment/Disability Duration Guidelines*

### **Low Back - Lumbar & Thoracic (Acute & Chronic)**

Discography	Not recommended. In the past, discography has been used as part of the pre-operative evaluation of patients for consideration of surgical intervention for lower back pain. However, the conclusions of recent, high quality studies on discography have significantly questioned the use of discography results as a preoperative indication for either IDET or spinal fusion. These studies have suggested that reproduction of the patient's specific back complaints on injection of one or more discs (concordance of symptoms) is of limited diagnostic value. (Pain production was found to be common in non-back pain patients, pain reproduction was found to be inaccurate in many patients with chronic back pain and abnormal psychosocial testing, and in this latter patient type, the test itself was sometimes found to produce significant symptoms in non-back pain controls more than a year after testing.) Also, the findings of discography have not been shown to consistently correlate well
-------------	--

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

with the finding of a High Intensity Zone (HIZ) on MRI. Discography may be justified if the decision has already been made to do a spinal fusion, and a negative discogram could rule out the need for fusion on that disc (but a positive discogram in itself would not allow fusion). ([Carragee-Spine, 2000](#)) ([Carragee2-Spine, 2000](#)) ([Carragee3-Spine, 2000](#)) ([Carragee4-Spine, 2000](#)) ([Bigos, 1999](#)) ([ACR, 2000](#)) ([Resnick, 2002](#)) ([Madan, 2002](#)) ([Carragee-Spine, 2004](#)) ([Carragee2, 2004](#)) ([Maghout-Juratli, 2006](#)) ([Pneumaticos, 2006](#)) ([Airaksinen, 2006](#)) ([Manchikanti, 2009](#)) Discography may help distinguish asymptomatic discs among morphologically abnormal discs in patients without psychosocial issues. Precise prospective categorization of discographic diagnoses may predict outcomes from treatment, surgical or otherwise. ([Derby, 2005](#)) ([Derby2, 2005](#)) ([Derby, 1999](#)) Positive discography was not highly predictive in identifying outcomes from spinal fusion. A recent study found only a 27% success from spinal fusion in patients with low back pain and a positive single-level low-pressure provocative discogram, versus a 72% success in patients having a well-accepted single-level lumbar pathology of unstable spondylolisthesis. ([Carragee, 2006](#)) The prevalence of positive discogram may be increased in subjects with chronic low back pain who have had prior surgery at the level tested for lumbar disc herniation. ([Heggeness, 1997](#)) Invasive diagnostics such as provocative discography have not been proven to be accurate for diagnosing various spinal conditions, and their ability to effectively guide therapeutic choices and improve ultimate patient outcomes is uncertain. ([Chou, 2008](#)) Although discography, especially combined with CT scanning, may be more accurate than other radiologic studies in detecting degenerative disc disease, its ability to improve surgical outcomes has yet to be proven. It is routinely used before IDET, yet only occasionally used before spinal fusion. ([Cohen, 2005](#)) Provocative discography is not recommended because its diagnostic accuracy remains uncertain, false-positives can occur in persons without low back pain, and its use has not been shown to improve clinical outcomes. ([Chou2, 2009](#)) This recent RCT concluded that, compared with discography, injection of a small amount of bupivacaine into the painful disc was a better tool for the diagnosis of discogenic LBP. ([Ohtori, 2009](#)) Discography may cause disc degeneration. Even modern discography techniques using small gauge needle and limited pressurization resulted in accelerated disc degeneration (35% in the discography group compared to 14% in the control group), disc herniation, loss of disc height and signal and the development of reactive endplate changes compared to match-controls. These findings are of concern for several reasons. Discography as a diagnostic test is controversial and in view of these findings the utility of this test should be reviewed. Furthermore, discography in current practice will often include injecting discs with a low probability of being symptomatic in an effort to validate other disc injections, a so-called control disc. Although this strategy has never been confirmed to increase test validity or utility, injecting normal discs even with small gauge needles appears to increase the rate of degeneration in these discs over time. The phenomenon of accelerated adjacent segment degeneration adjacent to fusion levels may be, in part, explained by previous disc puncture if discography was used in segments adjacent to the fusion. Similarly, intradiscal therapeutic strategies (injecting steroids, sclerosing agents, growth factors, etc.) have been proposed as a method to treat, arrest or prevent symptomatic disc disease. This study suggests that the injection procedure itself is not completely innocuous and a recalculation of these demonstrated risks versus hypothetical benefits should be considered. ([Carragee, 2009](#)) More in vitro evidence that discography may cause disc degeneration. ([Gruber, 2012](#)) Discography involves the injection of a water-soluble imaging

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

material directly into the nucleus pulposus of the disc. Information is then recorded about the pressure in the disc at the initiation and completion of injection, about the amount of dye accepted, about the configuration and distribution of the dye in the disc, about the quality and intensity of the patient's pain experience and about the pressure at which that pain experience is produced. Both routine x-ray imaging during the injection and post-injection CT examination of the injected discs are usually performed as part of the study. There are two diagnostic objectives: (1) to evaluate radiographically the extent of disc damage on discogram and (2) to characterize the pain response (if any) on disc injection to see if it compares with the typical pain symptoms the patient has been experiencing. Criteria exist to grade the degree of disc degeneration from none (normal disc) to severe. A symptomatic degenerative disc is considered one that disperses injected contrast in an abnormal, degenerative pattern, extending to the outer margins of the annulus and at the same time reproduces the patient's lower back complaints (concordance) at a low injection pressure. Discography is not a sensitive test for radiculopathy and has no role in its confirmation. It is, rather, a confirmatory test in the workup of axial back pain and its validity is intimately tied to its indications and performance. As stated, it is the end of a diagnostic workup in a patient who has failed all reasonable conservative care and remains highly symptomatic. Its validity is enhanced (and only achieves potential meaningfulness) in the context of an MRI showing both dark discs and bright, normal discs -- both of which need testing as an internal validity measure. And the discogram needs to be performed according to contemporary diagnostic criteria -- namely, a positive response should be low pressure, concordant at equal to or greater than a VAS of 7/10 and demonstrate degenerative changes (dark disc) on MRI and the discogram with negative findings of at least one normal disc on MRI and discogram. See also [Functional anesthetic discography \(FAD\)](#).

## **Discography is Not Recommended in ODG.**

### **Patient selection criteria for Discography if provider & payor agree to perform anyway:**

- o Back pain of at least 3 months duration
- o Failure of recommended conservative treatment including active physical therapy
- o An MRI demonstrating one or more degenerated discs as well as one or more normal appearing discs to allow for an internal control injection (injection of a normal disc to validate the procedure by a lack of a pain response to that injection)
- o Satisfactory results from detailed psychosocial assessment (discography in subjects with emotional and chronic pain problems has been linked to reports of significant back pain for prolonged periods after injection, and therefore should be avoided)
- o Intended as screening tool to assist surgical decision making, i.e., the surgeon feels that lumbar spine fusion is appropriate but is looking for this to determine if it is not indicated (although discography is not highly predictive) ([Carragee, 2006](#)) NOTE: In a situation where the selection criteria and other surgical indications for fusion are conditionally met, discography can be considered in preparation for the surgical

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

	<p>procedure. However, all of the qualifying conditions must be met prior to proceeding to discography as discography should be viewed as a non-diagnostic but confirmatory study for selecting operative levels for the proposed surgical procedure. Discography should not be ordered for a patient who does not meet surgical criteria.</p> <ul style="list-style-type: none"> <li>o Briefed on potential risks and benefits from discography and surgery</li> <li>o Single level testing (with control) (<a href="#">Colorado, 2001</a>)</li> <li>o Due to high rates of positive discogram after surgery for lumbar disc herniation, this should be potential reason for non-certification</li> </ul>
--	--

Fusion (spinal)	<p>Not recommended for patients who have less than six months of failed recommended conservative care unless there is objectively demonstrated severe structural instability and/or acute or progressive neurologic dysfunction, but recommended as an option for spinal fracture, dislocation, spondylolisthesis or frank neurogenic compromise, subject to the selection criteria outlined in the section below entitled, "<a href="#">Patient Selection Criteria for Lumbar Spinal Fusion</a>," after 6 months of conservative care. For workers' comp populations, see also the heading, "<a href="#">Lumbar fusion in workers' comp patients</a>." After screening for psychosocial variables, outcomes are improved and fusion may be recommended for degenerative disc disease with spinal segment collapse with or without neurologic compromise after 6 months of compliance with recommended <a href="#">conservative therapy</a>. [For spinal instability criteria, see AMA Guides (<a href="#">Andersson, 2000</a>)] For complete references, see separate document with all studies focusing on <a href="#">Fusion (spinal)</a>. There is limited scientific evidence about the long-term effectiveness of fusion for degenerative disc disease compared with natural history, placebo, or conservative treatment. Studies conducted in order to compare different surgical techniques have shown success for fusion in carefully selected patients. (<a href="#">Gibson-Cochrane, 2000</a>) (<a href="#">Savolainen, 1998</a>) (<a href="#">Wetzel, 2001</a>) (<a href="#">Molinari, 2001</a>) (<a href="#">Bigos, 1999</a>) (<a href="#">Washington, 1995</a>) (<a href="#">DeBarard-Spine, 2001</a>) (<a href="#">Fritzell-Spine, 2001</a>) (<a href="#">Fritzell-Spine, 2002</a>) (<a href="#">Deyo-NEJM, 2004</a>) (<a href="#">Gibson-Cochrane/Spine, 2005</a>) (<a href="#">Soegaard, 2005</a>) (<a href="#">Glassman, 2006</a>) (<a href="#">Atlas, 2006</a>) According to the recently released AANS/NASS Guidelines, lumbar fusion is recommended as a treatment for carefully selected patients with disabling low back pain due to one- or two-level degenerative disc disease after failure of an appropriate period of conservative care. This recommendation was based on one study that contained numerous flaws, including a lack of standardization of conservative care in the control group. At the time of the 2-year follow up it appeared that pain had significantly increased in the surgical group from year 1 to 2. Follow-up post study is still pending publication. In addition, there remains no direction regarding how to define the "carefully selected patient." (<a href="#">Resnick, 2005</a>) (<a href="#">Fritzell, 2004</a>) A recently published well respected international guideline, the "European Guidelines," concluded that fusion surgery for nonspecific chronic LBP cannot be recommended unless 2 years of all other recommended conservative treatments – including multidisciplinary approaches with combined programs of cognitive intervention and exercises – have failed, or such combined programs are not available, and only then in carefully selected patients with maximum 2-level degenerative disc disease. (<a href="#">Airaksinen, 2006</a>) For chronic LBP, exercise and cognitive intervention may be equivalent to lumbar fusion without the potentially</p>
-----------------	---



# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

high surgical complication rates. ([Ivar Brox-Spine, 2003](#)) ([Keller-Spine, 2004](#)) ([Fairbank-BMJ, 2005](#)) ([Brox, 2006](#)) In acute spinal cord injury (SCI), if the spine is unstable following injury, surgical fusion and bracing may be necessary. ([Bagnall-Cochrane, 2004](#)) ([Siebenga, 2006](#)) A study on improving quality through identifying inappropriate care found that use of guideline-based Utilization Review (UR) protocols resulted in a denial rate for lumbar fusion 59 times as high as denial rates using non-guideline based UR. ([Wickizer, 2004](#)) The profit motive and market medicine have had a significant impact on clinical practice and research in the field of spine surgery. ([Weiner-Spine, 2004](#)) ([Shah-Spine, 2005](#)) ([Abelson, 2006](#)) Data on geographic variations in medical procedure rates suggest that there is significant variability in spine fusion rates, which may be interpreted to suggest a poor professional consensus on the appropriate indications for performing spinal fusion. ([Deyo-Spine, 2005](#)) ([Weinstein, 2006](#)) Outcomes from complicated surgical fusion techniques (with internal fixation) may be no better than the traditional posterolateral fusion. ([van Tulder, 2006](#)) ([Maghout-Juratli, 2006](#)) Despite the new technologies, reoperation rates after lumbar fusion have become higher. ([Martin, 2007](#)) According to the recent Medicare Coverage Advisory Committee Technology Assessment, the evidence for lumbar spinal fusion does not conclusively demonstrate short-term or long-term benefits compared with nonsurgical treatment for elderly patients. ([CMS, 2006](#)) When lumbar fusion surgery is performed, either with lateral fusion alone or with interbody fusion, unlike cervical fusion, there is no absolute contraindication to patients returning even to contact sports after complete recovery from surgery. Like patients with a thoracic injury, those with a lumbar injury should be pain free, have no disabling neurological deficit, and exhibit evidence of bone fusion on x-ray films before returning. ([Burnett, 2006](#)) A recent randomized controlled trial comparing decompression with decompression and instrumented fusion in patients with foraminal stenosis and single-level degenerative disease found that patients universally improved with surgery, and this improvement was maintained at 5 years. However, no obvious additional benefit was noted by combining decompression with an instrumented fusion. ([Hallett, 2007](#)) Discography may be supported if the decision has already been made to do a spinal fusion, and a negative discogram could rule out the need for fusion on that disc (but a positive discogram in itself would not justify fusion). Discography may help distinguish asymptomatic discs among morphologically abnormal discs in patients without psychosocial issues. Precise prospective categorization of discographic diagnoses may predict outcomes from treatment, surgical or otherwise. ([Derby, 2005](#)) ([Derby2, 2005](#)) ([Derby, 1999](#)) New research shows that healthcare expenditures for back and neck problems have increased substantially over time, but with little improvement in healthcare outcomes such as functional disability and work limitations. Rates of imaging, injections, opiate use, and spinal surgery have increased substantially over the past decade, but it is unclear what impact, if any, this has had on health outcomes. ([Martin, 2008](#)) The efficacy of surgery for nonspecific back pain is uncertain. There may be some patients for whom surgery, fusion specifically, might be helpful, but it is important for doctors to discuss the fact that surgery doesn't tend to lead to huge improvements on average, about a 10- to 20-point improvement in function on a 100-point scale, and a significant proportion of patients still need to take pain medication and don't return to full function. ([Chou, 2008](#)) This study showed that fusion for chronic lower back pain was the least successful common orthopaedic surgery. The study compared the gains in quality of life achieved by total hip replacement, total knee replacement, surgery for spinal stenosis, disc excision for lumbar disc herniation, and arthrodesis for chronic low back pain. For

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

chronic lower back pain, improvements were statistically significant but clinically negligible. Although pain was reduced and function improved slightly, outcomes remained in the moderately affected range, quality of life was not improved and rendered worse, on average. While surgery for spinal stenosis and for disc herniation compare well with archetypical orthopaedic operations, the outcomes of surgery for chronic lower back pain do not even approach those of other orthopaedic procedures, and the data show that patients with back pain are rendered worse off by surgery. (Hansson, 2008) Recent studies document a 220% increase in lumbar spinal fusion surgery rates, without demonstrated improvements in patient outcomes or disability rates. (Deyo, 2009) In a study of 2,378 Washington State workers' compensation claimants who underwent fusion to assess the frequency, timing, and causes of death, the 3-year cumulative mortality rate post-fusion was 1.93% and analgesic-related deaths were responsible for 21% of all deaths and 31.4% of all potential life lost. (Juratli, 2009) A study to compare the surgical experience, clinical outcomes, and effect on body weight between obese and morbidly obese patients undergoing lumbar spine fusion surgery concluded that clinical outcomes were independent of the BMI of the patient, but the incidence of postoperative complications was significant in 45% of morbidly obese and 44% of obese patients. The authors proposed that morbidly obese patients should undergo bariatric surgery before spine surgery in nonemergent situations. (Vaidya, 2009) For nonradicular low back pain with common degenerative changes, there is fair evidence that fusion is no better than intensive rehabilitation with a cognitive-behavioral emphasis for improvement in pain or function, and less than half of patients experience optimal outcomes (defined as no more than sporadic pain, slight restriction of function, and occasional analgesics) following fusion. (Chou, 2009) Posterolateral bone-grafting fusion is not necessary when a Denis type-B thoracolumbar burst fracture associated with a load-sharing score of  $\leq 6$  is treated with short-segment pedicle screw fixation. (Dai, 2009) Discography (and not merely the fusion) may actually be the cause of adjacent segment disc degeneration. This study suggested that the phenomenon of accelerated adjacent segment degeneration adjacent to fusion levels may be, in part, explained by previous disc puncture if discography was used in segments adjacent to the fusion. (Carragee, 2009) Among Medicare recipients, the frequency of complex fusion procedures for spinal stenosis increased 15-fold in just 6 years. The introduction and marketing of new surgical devices and financial incentives may stimulate more invasive surgery. (Deyo-JAMA, 2010) Results of this study suggest that postmenopausal female patients who underwent lumbar spinal instrumentation fusion were susceptible to subsequent vertebral fractures within 2 years after surgery (in 24% of patients). (Toyone, 2010) A four-year follow-up of an RCT of instrumented transpedicular fusion versus cognitive intervention and exercises for disc degeneration with chronic low back pain concluded that this invasive and high-cost procedure does not afford better outcomes compared with the conservative treatment approach to low back pain, and this study should give doctors pause when recommending lumbar fusion surgery without compelling indications, particularly when strong back rehabilitation programs are available. (Brox, 2010) The ECRI health technology assessment concluded that the evidence is insufficient to support lumbar fusion being more effective (to a clinically meaningful degree) than nonsurgical treatments (intensive exercise and rehabilitation plus cognitive behavioral therapy) in patients with and without prior surgery. (ECRI, 2007) There is a high rate of complications (56.4%) in spinal fusion procedures, especially related to instrumentation. (Campbell, 2011) The draft AHRQ Comparative Effectiveness Research concluded that limited data suggests

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

that fusion leads to greater improvement in back pain relief and function than physical therapy at 2-year followup, but whether the difference is clinically significant is unclear, and serious adverse events occurred in the fusion group but not the noninvasive-intervention group. ([Clancy, 2012](#)) Lumbar spinal fusion surgeries use bone grafts, and are sometimes combined with metal devices, to produce a rigid connection between two or more adjacent vertebrae. The therapeutic objective of spinal fusion surgery for patients with low back problems is to prevent any movement in the intervertebral spaces between the fused vertebrae, thereby reducing pain and any neurological deficits. See also [Adjacent segment disease/degeneration](#) (fusion) & [Iliac crest donor-site pain treatment](#).

*Lumbar fusion in workers' comp patients:* In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. Until further research is conducted there remains insufficient evidence to recommend fusion for chronic low back pain in the absence of stenosis and spondylolisthesis, and this treatment for this condition remains "under study." It appears that workers' compensation populations require particular scrutiny when being considered for fusion for chronic low back pain, as there is evidence of poorer outcomes in subgroups of patients who were receiving compensation or involved in litigation. ([Fritzell-Spine, 2001](#)) ([Harris-JAMA, 2005](#)) ([Maghout-Juratli, 2006](#)) ([Atlas, 2006](#)) Despite poorer outcomes in workers' compensation patients, utilization is much higher in this population than in group health. ([Texas, 2001](#)) ([NCCI, 2006](#)) Presurgical biopsychosocial variables predict patient outcomes from lumbar fusion, which may help improve patient selection. Workers' compensation status, smoking, depression, and litigation were the most consistent presurgical predictors of poorer patient outcomes. Other predictors of poor results were number of prior low back operations, low household income, and older age. ([DeBerard-Spine, 2001](#)) ([DeBerard, 2003](#)) ([Deyo, 2005](#)) ([LaCaille, 2005](#)) ([Trief-Spine, 2006](#)) Obesity and litigation in workers' compensation cases predict high costs associated with interbody cage lumbar fusion. ([LaCaille, 2007](#)) A recent study of 725 workers' comp patients in Ohio who had lumbar fusion found only 6% were able to go back to work a year later, 27% needed another operation, and over 90% were in enough pain that they were still taking narcotics at follow-up. ([Nguyen, 2007](#)) A recent case-control study of lumbar fusion outcomes in worker's compensation (WC) patients concluded that only 9% of patients receiving WC achieved substantial clinical benefit compared to 33% of those not receiving WC. ([Carreon, 2009](#)) This large historical cohort study suggests that lumbar fusion may not be an effective operation in workers' compensation patients with disc degeneration, disc herniation, and/or radiculopathy, and it is associated with significant increase in disability, opiate use, prolonged work loss, and poor RTW status. ([Nguyen, 2011](#)) After controlling for covariates known to affect lumbar fusion outcomes, patients on workers' comp have significantly less improvement. ([Carreon, 2010](#)) The presidents of AAOS, NASS, AANS, CNS, and SAS issued a joint statement to BlueCross BlueShield recommending patient selection criteria for lumbar fusion in degenerative disc disease. The criteria included at least one year of physical and cognitive therapy, inflammatory endplate changes (i.e., Modic changes), moderate to severe disc space collapse, absence of significant psychological comorbidities (e.g. depression, somatization disorder), and absence of litigation or compensation issues. The criteria of denying fusion if there are compensation issues may apply to workers' compensation patients. ([Rutka, 2011](#)) On the other hand, a separate policy statement

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

from the International Society for the Advancement of Spine Surgery disagrees that worker's compensation should be a contraindication for lumbar fusion. ([ISASS, 2011](#)) This study demonstrated a significant difference in outcomes after lumbar spinal fusion between workers' comp populations and those on long-term disability insurance. Both populations only achieved marginal improvement, but workers' comp had a clear, negative influence on outcome even when compared to disability patients. ([Gum, 2012](#))

*Lumbar fusion for spondylolisthesis:* Recommended as an option for spondylolisthesis. Patients with increased instability of the spine after surgical decompression at the level of degenerative spondylolisthesis are candidates for fusion. ([Eckman, 2005](#)) This study found only a 27% success from spinal fusion in patients with low back pain and a positive single-level low-pressure provocative discogram, versus a 72% success in patients having a well-accepted single-level lumbar pathology of unstable spondylolisthesis. ([Carragee, 2006](#)) Unilateral instrumentation used for the treatment of degenerative lumbar spondylolisthesis is as effective as bilateral instrumentation. ([Fernandez-Fairen, 2007](#)) Patients with degenerative spondylolisthesis and spinal stenosis who undergo standard decompressive laminectomy (with or without fusion) showed substantially greater improvement in pain and function during a period of 2 years than patients treated nonsurgically, according to the recent results from the Spine Patient Outcomes Research Trial (SPORT). ([Weinstein-spondylolisthesis, 2007](#)) ([Deyo-NEJM, 2007](#)) For degenerative lumbar spondylolisthesis, spinal fusion may lead to a better clinical outcome than decompression alone. No conclusion about the clinical benefit of instrumenting a spinal fusion can be made, but there is moderate evidence that the use of instrumentation improves the chance of achieving solid fusion. ([Martin, 2007](#)) A recent systematic review of randomized trials comparing lumbar fusion surgery to nonsurgical treatment of chronic back pain associated with lumbar disc degeneration, concluded that surgery may be more efficacious than unstructured nonsurgical care but may not be more efficacious than structured cognitive-behavior therapy. Methodological limitations of the randomized trials prevented firm conclusions. ([Mirza, 2007](#)) A comparison of surgical and nonoperative outcomes between degenerative spondylolisthesis and spinal stenosis patients from the SPORT trial found that fusion was most appropriate for spondylolisthesis, with or without listhesis, and decompressive laminectomy alone most appropriate for spinal stenosis. ([Pearson, 2010](#)) The latest SPORT study concluded that leg pain is associated with better surgical fusion outcomes in spondylolisthesis than low back pain. ([Pearson, 2011](#)) Comparative effectiveness evidence from SPORT shows good value for laminectomy and/or bilateral single-level fusion after an imaging-confirmed diagnosis of degenerative spondylolisthesis [as recommended in ODG], compared with nonoperative care over 4 years. ([Tosteson, 2011](#))

*Lumbar fusion for Scheuermann's kyphosis:* Recommended as an option for adult patients with severe deformities (e.g. more than 70 degrees for thoracic kyphosis), neurological symptoms exist, and pain cannot be adequately resolved non-operatively (e.g. physical therapy, back exercises). Good outcomes have been found in a relatively large series of patients undergoing either combined anterior-posterior or posterior only fusion for Scheuermann's kyphosis. ([Lonner, 2007](#))

# The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

## Patient Selection Criteria for Lumbar Spinal Fusion:

For chronic low back problems, fusion should not be considered within the first 6 months of symptoms, except for fracture, dislocation or progressive neurologic loss. Indications for spinal fusion may include: (1) Neural Arch Defect - Spondylytic spondylolisthesis, congenital neural arch hypoplasia. (2) Segmental Instability (objectively demonstrable) - Excessive motion, as in degenerative spondylolisthesis, surgically induced segmental instability and mechanical intervertebral collapse of the motion segment and advanced degenerative changes after surgical discectomy, with relative angular motion greater than 20 degrees. ([Andersson, 2000](#)) ([Luers, 2007](#)) (3) Primary Mechanical Back Pain (i.e., pain aggravated by physical activity)/Functional Spinal Unit Failure/Instability, including one or two level segmental failure with progressive degenerative changes, loss of height, disc loading capability. In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. There is a lack of support for fusion for mechanical low back pain for subjects with failure to participate effectively in active rehab pre-op, total disability over 6 months, active psych diagnosis, and narcotic dependence. Spinal instability criteria includes lumbar inter-segmental movement of more than 4.5 mm. ([Andersson, 2000](#)) (4) Revision Surgery for failed previous operation(s) if significant functional gains are anticipated. Revision surgery for purposes of pain relief must be approached with extreme caution due to the less than 50% success rate reported in medical literature. (5) Infection, Tumor, or Deformity of the lumbosacral spine that cause intractable pain, neurological deficit and/or functional disability. (6) After failure of two discectomies on the same disc, fusion may be an option at the time of the third discectomy, which should also meet the ODG criteria. (See [ODG Indications for Surgery -- Discectomy.](#))

**Pre-Operative Surgical Indications Recommended:** Pre-operative clinical surgical indications for spinal fusion should include all of the following: (1) All pain generators are identified and treated; & (2) All physical medicine and manual therapy interventions are completed; & (3) X-rays demonstrating spinal instability and/or myelogram, CT-myelogram, or discography (see [discography criteria](#)) & MRI demonstrating disc pathology correlated with symptoms and exam findings; & (4) Spine pathology limited to two levels; & (5) [Psychosocial screen](#) with confounding issues addressed. (6) For any potential fusion surgery, it is recommended that the injured worker refrain from smoking for at least six weeks prior to surgery and during the period of fusion healing. ([Colorado, 2001](#)) ([BlueCross BlueShield, 2002](#))

For average hospital LOS after criteria are met, see [Hospital length of stay](#) (LOS).

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:**

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE**
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES**
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES**
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN**
- INTERQUAL CRITERIA**
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE, AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS**
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES**
- MILLIMAN CARE GUIDELINES**
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES**
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR**
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS**
- TEXAS TACADA GUIDELINES**
- TMF SCREENING CRITERIA MANUAL**
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)**
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)**