

Notice of Independent Review Decision

DATE OF REVIEW: August 29, 2013

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

64635, destroy lumb/sac facet JNT, units REQ: 1, 64636, destroy L/S facet JNT ADDL, units REQ: 1, outpatient RFTC left side L1/2 and L2/3

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION

This case was reviewed by a physician board certified in Physical Medicine and Rehabilitation currently licensed and practicing in the state of Texas.

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

INFORMATION PROVIDED TO THE IRO FOR REVIEW

Type of Document Received	Date(s) of Record
MRI of the lumbar spine	03/30/2012
X-rays of the lumbosacral spine	12/10/2012
Office visit	05/24/2013
Office visit	06/27/2013
Office visit	07/26/2013
An adverse determination letter	07/08/2013
A reconsideration adverse determination letter regarding	08/02/2013

EMPLOYEE CLINICAL HISTORY [SUMMARY]:

This is a male who injured his lower back on xx/xx/xx. Subsequently, he had L4-5 and L5-S1 fusion in 2008 but continued to have lower back pain. He has been treated with



**MEDICAL EVALUATORS
OF TEXAS** ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

conservative care including physical therapy and pain medications. He also had facet joint medial branch blocks and facet joint radiofrequency neurotomies at L1-2 and L2-3 levels. MRI dated 03/20/2012 showed interval development of disk degeneration and minimal retrolisthesis at L2-3 without canal or foraminal stenosis and degenerative disk disease at T11-12. X-rays dated 12/10/2012 showed surgical changes at L4-5 and L5-S1 and no evidence of instability. A progress report dated 07/26/2013 indicated that he reported lower back pain level of 2-3/10 as best and 6 without medications. On physical exam, there was tenderness to palpation at L4-S1 spinous interspace. Sensation was intact to light touch, reflexes were 2+, and motor testing was 5/5 throughout lower extremities. He is currently recommended lumbar RFTC at L1-2 and L2-3.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

The claimant underwent a lumbar RFTC at L1-2, L2-3 on 06/05/2012. The claimant reported 70% relief at 4 weeks, 40% relief at 8 weeks, and pain returned back to baseline at 12 weeks. Based on ODG guidelines, "while repeat neurotomies may be required, they should not occur at an interval of less than 6 months from the first procedure. A neurotomy should not be repeated unless duration of relief from the first procedure is documented for at least 12 weeks at $\geq 50\%$ relief. The current literature does not support that the procedure is successful without sustained pain relief (generally of at least 6 months duration). No more than 3 procedures should be performed in a year's period."

As stated above, the relief from initial lumbar RFTC at L1-2, L2-3 on 06/05/2012 lasted between 8-12 weeks which does not meet ODG guidelines of required sustained relief of at least 6 months after a RFTC procedure to justify a repeat procedure. As such, the request is non-certified.

ODG CRITERIA FOR USE OF FACET JOINT RADIOFREQUENCY NEUROTOMY:

- (1) Treatment requires a diagnosis of facet joint pain using a medial branch block as described above. See Facet joint diagnostic blocks(injections).
- (2) While repeat neurotomies may be required, they should not occur at an interval of less than 6 months from the first procedure. A neurotomy should not be repeated unless duration of relief from the first procedure is documented for at least 12 weeks at $\geq 50\%$ relief. The current literature does not support that the procedure is successful without sustained pain relief (generally of at least 6 months duration). No more than 3 procedures should be performed in a year's period.
- (3) Approval of repeat neurotomies depends on variables such as evidence of adequate diagnostic blocks, documented improvement in VAS score, decreased medications and documented improvement in function.
- (4) No more than two joint levels are to be performed at one time.
- (5) If different regions require neural blockade, these should be performed at intervals of no sooner than one week, and preferably 2 weeks for most blocks.



MEDICAL EVALUATORS OF TEXAS ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

(6) There should be evidence of a formal plan of additional evidence-based conservative care in addition to facet joint therapy.

ODG CRITERIA FOR FACET JOINTS DIAGNOSTIC BLOCKS (INJECTIONS)

Recommend no more than one set of medial branch diagnostic blocks prior to facet neurotomy, if neurotomy is chosen as an option for treatment (a procedure that is still considered “under study”). Diagnostic blocks may be performed with the anticipation that if successful, treatment may proceed to facet neurotomy at the diagnosed levels. Current research indicates that a minimum of one diagnostic block be performed prior to a neurotomy, and that this be a medial branch block (MBB). Although it is suggested that MBBs and intra-articular blocks appear to provide comparable diagnostic information, the results of placebo-controlled trials of neurotomy found better predictive effect with diagnostic MBBs. In addition, the same nerves are tested with the MBB as are treated with the neurotomy. The use of a confirmatory block has been strongly suggested due to the high rate of false positives with single blocks (range of 25% to 40%) but this does not appear to be cost effective or to prevent the incidence of false positive response to the neurotomy procedure itself. (Cohen, 2007) (Bogduk, 2000) (Cohen2, 2007)

(Mancchukonda, 2007) (Dreyfuss, 2000) (Manchikanti2, 2003) (Datta, 2009)

Etiology of false positive blocks: Placebo response (18-32%), use of sedation, liberal use of local anesthetic, and spread of injectate to other pain generators. The concomitant use of sedative during the block can also interfere with an accurate diagnosis. (Cohen, 2007)

MBB procedure: The technique for medial branch blocks in the lumbar region requires a block of 2 medial branch nerves (MBN). The recommendation is the following: (1) L1-L2 (T12 and L1 MBN); (2) L2-L3 (L1 and L2 MBN); (3) L3-L4 (L2 and L3 MBN); (4) L4-L5 (L3 and L4 MBN); (5) L5-S1: the L4 and L5 MBN are blocked, and it is recommended that S1 nerve be blocked at the superior articular process. Blocking two joints such as L3-4 and L4-5 will require blocks of three nerves (L2, L3 and L4). Blocking L4-5 and L5-S1 will require blocks of L3, L4, L5 with the option of blocking S1. (Clemans, 2005) The volume of injectate for diagnostic medial branch blocks must be kept to a minimum (a trace amount of contrast with no more than 0.5 cc of injectate), as increased volume may anesthetize other potential areas of pain generation and confound the ability of the block to accurately diagnose facet pathology. Specifically, the concern is that the lateral and intermediate branches will be blocked; nerves that innervate the paraspinal muscles and fascia, ligaments, sacroiliac joints and skin. (Cohen, 2007) Intraarticular blocks also have limitations due to the fact that they can be technically challenging, and if the joint capsule ruptures, injectate may diffuse to the epidural space, intervertebral foramen, ligamentum flavum and paraspinal musculature. (Cohen, 2007) (Washington, 2005) (Manchikanti , 2003) (Dreyfuss, 2003) (BlueCross BlueShield, 2004) (Pneumatics, 2006) (Boswell, 2007) (Boswell2, 2007) A recent meta-analysis concluded that there is insufficient evidence to evaluate validity or utility of diagnostic selective nerve root block, intra-articular facet joint block, medial branch block, or sacroiliac joint block as diagnostic procedures for low back pain with or without radiculopathy. (Chou2, 2009) This study suggests that proceeding to radiofrequency denervation without a diagnostic block is the



**MEDICAL EVALUATORS
OF TEXAS** ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

most cost-effective treatment paradigm, but does not result in the best pain outcomes. (Cohen, 2010) See also Facet joint pain, signs & symptoms; Facet joint radiofrequency neurotomy; Facet joint medial branch blocks(therapeutic injections); & Facet joint intra-articular injections (therapeutic blocks). Also see Neck Chapter and Pain Chapter.

Criteria for the use of diagnostic blocks for facet “mediated” pain:

Clinical presentation should be consistent with facet joint pain, signs & symptoms.

1. One set of diagnostic medial branch blocks is required with a response of $\geq 70\%$. The pain response should last at least 2 hours for Lidocaine.
2. Limited to patients with low-back pain that is non-radicular and at no more than two levels bilaterally.
3. There is documentation of failure of conservative treatment (including home exercise, PT and NSAIDs) prior to the procedure for at least 4-6 weeks.
4. No more than 2 facet joint levels are injected in one session (see above for medial branch block levels).
5. Recommended volume of no more than 0.5 cc of injectate is given to each joint.
6. No pain medication from home should be taken for at least 4 hours prior to the diagnostic block and for 4 to 6 hours afterward.
7. Opioids should not be given as a “sedative” during the procedure.
8. The use of IV sedation (including other agents such as midazolam) may be grounds to negate the results of a diagnostic block, and should only be given in cases of extreme anxiety.
9. The patient should document pain relief with an instrument such as a VAS scale, emphasizing the importance of recording the maximum pain relief and maximum duration of pain. The patient should also keep medication use and activity logs to support subjective reports of better pain control.
10. Diagnostic facet blocks should not be performed in patients in whom a surgical procedure is anticipated. (Resnick, 2005)
11. Diagnostic facet blocks should not be performed in patients who have had a previous fusion procedure at the planned injection level. [Exclusion Criteria that would require UR physician review: Previous fusion at the targeted level. (Franklin, 2008)]



**MEDICAL EVALUATORS
OF TEXAS** ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER
CLINICAL BASIS USED TO MAKE THE DECISION:**

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES**
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)