

ReviewTex. Inc.
1818 Mountjoy Drive
San Antonio, TX 78232
(phone) 210-598-9381 (fax) 210-598-9382
reviewtex@hotmail.com

Notice of Independent Review Decision

Date notice sent to all parties: February 11, 2014

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Physical Therapy (Active Physical Rehabilitation – 4 sessions)

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

Board Certified Chiropractic Examiner

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

X Upheld (Agree)

Provide a description of the review outcome that clearly states whether medical necessity exists for each of the health care services in dispute.

INFORMATION PROVIDED TO THE IRO FOR REVIEW:

Cover sheet and working documents
Patient re-evaluation dated 12/16/13
Utilization review determination dated 12/23/13, 01/14/14
Letter dated 01/27/14
Notice of reconsideration outcome dated 01/15/14

PATIENT CLINICAL HISTORY [SUMMARY]:

The patient is a female whose date of injury is xx/xx/xx. The mechanism of injury is not described. The earliest record provided is a patient re-evaluation dated 12/16/13. This report states that the patient presents with a flare-up of back pain. She stated that it was improved after doing physical therapy. She has been

working in delivering and still has occasional exacerbation of pain. Her stress level has been improved. The patient underwent some sessions of active care. On physical examination palpation of the cervical spine noted increased tenderness at the base of the cervical spine extending into the mid and upper cervical. Increased tenderness was also noted in the lumbar spine, mostly at the left latissimus dorsi. No myospasms are noted. Spurling's was negative for radiculopathy. Cervical distraction was positive for relief of pain. Max foraminal compression is positive on the left. Straight leg raising is positive on the left side at 65 degrees. Kemp's is positive for increased lumbar pain. Lumbar flexion is 35 degrees, extension at 10 degrees. Muscle testing is +5/5 in all regions. Impression is cervical radiculitis, shoulder sprain/strain, elbow sprain/strain, lumbar radiculitis, wrist contusion and thoracic sprain/strain.

Initial request for physical therapy (active physical rehabilitation 4 sessions) was non-certified on 12/23/13 noting that relevant information is lacking including exact mechanism of injury, objective exam findings, comprehensive treatment to date, and objective response. Additionally, due to the remote nature of the patient's injury, he can be expected to have recovered at this time. The denial was upheld on appeal dated 01/14/14 noting that the patient is stated to have completed therapy although the results of such treatment have not been provided. ODG guidelines recommend a total of 10 visits over 8 weeks for thoracic and lumbosacral neuritis/radiculitis. Additionally, 9-10 visits are recommended for rotator cuff disease and elbow sprains. The total amount of prior therapy completed was not stated. An up-to-date rationale for the requested therapy has not been provided.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS, AND CONCLUSIONS USED TO SUPPORT THE DECISION:

Based on the clinical information provided, the request for physical therapy (active physical rehabilitation – 4 sessions) is not recommended as medically necessary. There is no comprehensive assessment of treatment completed to date or the patient's response thereto submitted for review. It is unclear how many sessions of physical therapy the patient has completed to date. There are no specific, time-limited treatment goals provided. Given the chronicity of the date of injury, the patient should be well-versed in an independent, self-directed home exercise program at this time.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

X MEDICAL JUDGEMENT, CLINICAL EXPERIENCE, AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

X ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

ODG Low Back Chapter

Physical therapy (PT)

Recommended. There is strong evidence that physical methods, including exercise and return to normal activities, have the best long-term outcome in employees with low back pain. See also Exercise. Direction from physical and occupational therapy providers can play a role in this, with the evidence supporting active therapy and not extensive use of passive modalities. The most effective strategy may be delivering individually designed exercise programs in a supervised format (for example, home exercises with regular therapist follow-up), encouraging adherence to achieve high dosage, and stretching and muscle-strengthening exercises seem to be the most effective types of exercises for treating chronic low back pain.

(Hayden, 2005) Studies also suggest benefit from early use of aggressive physical therapy ("sports medicine model"), training in exercises for home use, and a functional restoration program, including intensive physical training, occupational therapy, and psychological support. (Zigenfus, 2000) (Linz, 2002) (Cherkin-NEJM, 1998) (Rainville, 2002) Successful outcomes depend on a functional restoration program, including intensive physical training, versus extensive use of passive modalities. (Mannion, 2001) (Jousset, 2004) (Rainville, 2004) (Airaksinen, 2006) One clinical trial found both effective, but chiropractic was slightly more favorable for acute back pain and physical therapy for chronic cases. (Skargren, 1998) A spinal stabilization program is more effective than standard physical therapy sessions, in which no exercises are prescribed. With regard to manual therapy, this approach may be the most common physical therapy modality for chronic low back disorder, and it may be appropriate as a pain reducing modality, but it should not be used as an isolated modality because it does not concomitantly reduce disability, handicap, or improve quality of life. (Goldby-Spine, 2006) Better symptom relief is achieved with directional preference exercise. (Long, 2004) As compared with no therapy, physical therapy (up to 20 sessions over 12 weeks) following disc herniation surgery was effective. Because of the limited benefits of physical therapy relative to "sham" therapy (massage), it is open to question whether this treatment acts primarily physiologically, but psychological factors may contribute substantially to the benefits observed. (Erdogmus, 2007) In this RCT, exercise and stretching, regardless of whether it is achieved via yoga classes or conventional PT supervision, helps improve low back pain. (Sherman, 2011) See also specific physical therapy modalities, as well as Exercise; Work conditioning; Lumbar extension exercise equipment; McKenzie method; Stretching; & Aquatic therapy. [Physical therapy is the treatment of a disease or injury by the use of therapeutic

exercise and other interventions that focus on improving posture, locomotion, strength, endurance, balance, coordination, joint mobility, flexibility, activities of daily living and alleviating pain. (BlueCross BlueShield, 2005) As for visits with any medical provider, physical therapy treatment does not preclude an employee from being at work when not visiting the medical provider, although time off may be required for the visit.]

Active Treatment versus Passive Modalities: The use of active treatment instead of passive modalities is associated with substantially better clinical outcomes. In a large case series of patients with acute low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) The most commonly used active treatment modality is Therapeutic exercises (97110), but other active therapies may be recommended as well, including Neuromuscular reeducation (97112), Manual therapy (97140), and Therapeutic activities/exercises (97530). A recent RCT comparing active spinal stabilization exercises (using the GDS or Godelive Denys-Struyf method) with passive electrotherapy using TENS plus microwave treatment (considered conventional physical therapy in Spanish primary care), concluded that treatment of nonspecific LBP using the GDS method provides greater improvements in the midterm (6 months) in terms of pain, functional ability, and quality of life. (Arribas, 2009) In this RCT, two active interventions, multidisciplinary rehab (intensive, bio-psychosocial PT) and exercise (exercises targeted at trunk muscles together with stretching and relaxation), reduced the probability of sickness absence, and were more effective for pain than self-care advice at 12 months. (Rantonen, 2012)

Patient Selection Criteria: Multiple studies have shown that patients with a high level of fear-avoidance do much better in a supervised physical therapy exercise program, and patients with low fear-avoidance do better following a self-directed exercise program. When using the Fear-Avoidance Beliefs Questionnaire (FABQ), scores greater than 34 predicted success with PT supervised care. (Fritz, 2001) (Fritz, 2002) (George, 2003) (Klaber, 2004) (Riipinen, 2005) (Hicks, 2005) Without proper patient selection, routine physical therapy may be no more effective than one session of assessment and advice from a physical therapist. (Frost, 2004) Patients exhibiting the centralization phenomenon during lumbar range of motion testing should be treated with the specific exercises (flexion or extension) that promote centralization of symptoms. When findings from the patient's history or physical examination are associated with clinical instability, they should be treated with a trunk strengthening and stabilization exercise program. (Fritz-Spine, 2003) Practitioners must be cautious when implementing the wait-and-see approach for LBP, and once medical clearance has been obtained, patients should be advised to keep as active as possible. Patients presenting with high fear avoidance characteristics should have these concerns addressed aggressively to prevent long-term disability, and they should be encouraged to promote the resumption of physical activity. (Hanney, 2009)

Post Epidural Steroid Injections: ESIs are currently recommended as a possible option for short-term treatment of radicular pain (sciatica), defined as pain in dermatomal distribution with corroborative findings of radiculopathy. The general goal of physical therapy during the acute/subacute phase of injury is to decrease guarding, maintain motion, and decrease pain and inflammation. Progression of rehabilitation to a more advanced program of stabilization occurs in the maintenance phase once pain is controlled. There is little evidence-based research that addresses the use of physical therapy post ESIs, but it appears that most randomized controlled trials have utilized an ongoing, home directed program post injection. Based on current literature, the only need for further physical therapy treatment post ESI would be to emphasize the home exercise program, and this requirement would generally be included in the currently suggested maximum visits for the underlying condition, or at least not require more than 2 additional visits to reinforce the home exercise program. ESIs have been found to have limited effectiveness for treatment of chronic pain. The claimant should continue to follow a home exercise program post injection. (Luijsterburg, 2007) (Luijsterburg2, 2007) (Price, 2005) (Vad, 2002) (Smeal, 2004)

Post-surgical (discectomy) rehab: A recent Cochrane review concluded that exercise programs starting 4-6 weeks post-surgery seem to lead to a faster decrease in pain and disability than no treatment; high intensity exercise programs seem to lead to a faster decrease in pain and disability than low intensity programs; home exercises are as good as supervised exercises; and active programs do not increase the re-operation rate. Although it is not harmful to return to activity after lumbar disc surgery, it is still unclear what exact components should be included in rehabilitation programs. High intensity programs seem to be more effective but they could also be more expensive. Another question is whether all patients should be treated post-surgery or is a minimal intervention with the message return to an active lifestyle sufficient, with only patients that still have symptoms 4 to 6 weeks post-surgery requiring rehabilitation programs. (Ostelo, 2009) There is inconclusive evidence for the effectiveness of outpatient physical therapy after first lumbar discectomy. Although evidence from two trials suggested that intervention might reduce disability short-term, and more intensive intervention may be more beneficial than less intensive therapy, pooled results did not show statistically significant benefit. (Rushton, 2011) A systematic review yielded moderate to low quality evidence for effectiveness of postoperative exercise programs starting 4-6 weeks after lumbar disc surgery. Exercise programs seem to be more beneficial than no treatment, and high intensity exercises may be more effective than low intensity exercises. (Oosterhuis, 2013)

Post-surgical (fusion) rehab: Following lumbar spinal fusion, delayed start of rehabilitation results in better outcomes, and improvements in the group starting at 12-weeks were 4 times better than that in the 6-week group. (Oestergaard, 2012)

Allow for fading of treatment frequency (from up to 3 or more visits per week to 1 or less), plus active self-directed home PT. Also see other general guidelines that apply to all conditions under Physical Therapy in the ODG Preface, including assessment after a "six-visit clinical trial".

Lumbar sprains and strains (ICD9 847.2):

10 visits over 8 weeks

Sprains and strains of unspecified parts of back (ICD9 847):

10 visits over 5 weeks

Sprains and strains of sacroiliac region (ICD9 846):

Medical treatment: 10 visits over 8 weeks

Lumbago; Backache, unspecified (ICD9 724.2; 724.5):

9 visits over 8 weeks

Intervertebral disc disorders without myelopathy (ICD9 722.1; 722.2; 722.5; 722.6; 722.8):

Medical treatment: 10 visits over 8 weeks

Post-injection treatment: 1-2 visits over 1 week

Post-surgical treatment (discectomy/laminectomy): 16 visits over 8 weeks

Post-surgical treatment (arthroplasty): 26 visits over 16 weeks

Post-surgical treatment (fusion, after graft maturity): 34 visits over 16 weeks

Intervertebral disc disorder with myelopathy (ICD9 722.7)

Medical treatment: 10 visits over 8 weeks

Post-surgical treatment: 48 visits over 18 weeks

Spinal stenosis (ICD9 724.0):

10 visits over 8 weeks

See 722.1 for post-surgical visits

Sciatica; Thoracic/lumbosacral neuritis/radiculitis, unspecified (ICD9 724.3; 724.4):

10-12 visits over 8 weeks

See 722.1 for post-surgical visits

Curvature of spine (ICD9 737)

12 visits over 10 weeks

See 722.1 for post-surgical visits

Fracture of vertebral column without spinal cord injury (ICD9 805):

Medical treatment: 8 visits over 10 weeks

Post-surgical treatment: 34 visits over 16 weeks

Fracture of vertebral column with spinal cord injury (ICD9 806):

Medical treatment: 8 visits over 10 weeks

Post-surgical treatment: 48 visits over 18 weeks

Work conditioning (See also Procedure Summary entry):

10 visits over 8 weeks