

The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

Notice of Independent Review Decision

January 6, 2014

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Medical Necessity: 719.06 Effusion of lower leg joint, 717.9 Unspecified Internal Derangement of Knee (Left Knee Arthroscopy w/ Chondroplasty and partial Synovectomy)

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

The physician performing this review is Board Certified, American Board of Orthopedic Surgery. The physician has been in practice since 1998 and is licensed in Texas, Oklahoma, Minnesota and South Dakota.

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

Provide a description of the review outcome that clearly states whether medical necessity exists for each of the health care services in dispute.

Upon independent review, I find the previous adverse determination should be partially overturned.

The requested procedures include arthroscopy for chondroplasty and synovitis, which would be most appropriately considered for osteoarthritis of the knee, and as such, this would not be considered appropriate under ODG guidelines. The ODG indications for chondroplasty require a chondral defect on MRI, which is not specifically mentioned.

However, at this point, clinical signs and symptoms as well as physical examination would be consistent with the potential for meniscal tear. As such, with the patient's having failed an adequate trial of medications and physical therapy and continuing to have pain and functional limitations despite this

The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

conservative care with inconclusive imaging studies, diagnostic arthroscopy certainly would be considered appropriate under ODG guidelines.

PATIENT CLINICAL HISTORY [SUMMARY]:

reportedly injured on xx/xx/xx while on the job. He had immediate onset of a popping sensation, and subsequent to that has continued to have pain in his left knee with swelling, tenderness, and mechanical symptoms. He has undergone a trial of activity modification, physical therapy, anti-inflammatory medications, and an injection. He continues to be symptomatic despite these conservative efforts.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS, AND CONCLUSIONS USED TO SUPPORT THE DECISION:

The requested procedures include arthroscopy for chondroplasty and synovitis, which would be most appropriately considered for osteoarthritis of the knee, and as such, this would not be considered appropriate under ODG guidelines. The ODG indications for chondroplasty require a chondral defect on MRI, which is not specifically mentioned.

However, at this point, clinical signs and symptoms as well as physical examination would be consistent with the potential for meniscal tear. As such, with the patient's having failed an adequate trial of medications and physical therapy and continuing to have pain and functional limitations despite this conservative care with inconclusive imaging studies, diagnostic arthroscopy certainly would be considered appropriate under ODG guidelines.

The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205
Phone: 888-950-4333 Fax: 888-9504-4443

ODG -TWC

ODG Treatment

Integrated Treatment/Disability Duration Guidelines

Knee & Leg (Acute & Chronic)

[Back to ODG - TWC Index](#)

Chondroplasty	<p>Recommended as indicated below. Not recommended as a primary treatment for osteoarthritis, since arthroscopic surgery for knee osteoarthritis offers no added benefit to optimized physical therapy and medical treatment. (Kirkley, 2008) See also Meniscectomy.</p> <p>ODG Indications for Surgery™ -- Chondroplasty: Criteria for chondroplasty (shaving or debridement of an articular surface), requiring ALL of the following:</p> <ol style="list-style-type: none">1. Conservative Care: Medication. OR Physical therapy. PLUS2. Subjective Clinical Findings: Joint pain. AND Swelling. PLUS3. Objective Clinical Findings: Effusion. OR Crepitus. OR Limited range of motion. PLUS4. Imaging Clinical Findings: Chondral defect on MRI <p>(Washington, 2003) (Hunt, 2002) (Janecki, 1998)</p> <p>For average hospital LOS if criteria are met, see Hospital length of stay (LOS).</p>
Diagnostic arthroscopy	<p>Recommended as indicated below. Second look arthroscopy is only recommended in case of complications from OATS or ACI procedures, to assess how the repair is healing, or in individual cases that are ethically defensible for scientific reasons, only after a thorough and full informed consent procedure. (Vanlauwe, 2007) In patients with osteoarthritis, the value of MRI for a precise grading of the cartilage is limited, compared to diagnostic arthroplasty. When the assessment of the cartilage is crucial for a definitive decision regarding therapeutic options in patients with osteoarthritis, arthroscopy should not be generally replaced by MRI. The diagnostic values of MRI grading, using arthroscopy as reference standard, were calculated for each grade of cartilage damage. For grade 1, 2 and 3 lesions, sensitivities were relatively poor, whereas relatively better values were noted for grade 4 disorders. (von Engelhardt, 2010)</p> <p>ODG Indications for Surgery™ -- Diagnostic arthroscopy: Criteria for diagnostic arthroscopy:</p> <ol style="list-style-type: none">1. Conservative Care: Medications. OR Physical therapy. PLUS2. Subjective Clinical Findings: Pain and functional limitations continue despite conservative care. PLUS3. Imaging Clinical Findings: Imaging is inconclusive. <p>(Washington, 2003) (Lee, 2004)</p> <p>For average hospital LOS if criteria are met, see Hospital length of stay (LOS).</p>
Meniscectomy	Recommended as indicated below for symptomatic meniscal tears. Not

The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

recommended for osteoarthritis (OA) in the absence of meniscal findings. ([Kirkley, 2008](#)) Meniscectomy is a surgical procedure associated with a high risk of knee osteoarthritis (OA). One study concludes that the long-term outcome of meniscal injury and surgery appears to be determined largely by the type of meniscal tear, and that a partial meniscectomy may have better long-term results than a subtotal meniscectomy for a degenerative tear. ([Englund, 2001](#)) Another study concludes that partial meniscectomy may allow a slightly enhanced recovery rate as well as a potentially improved overall functional outcome including better knee stability in the long term compared with total meniscectomy. ([Howell-Cochrane, 2002](#)) The following characteristics were associated with a surgeon's judgment that a patient would likely benefit from knee surgery: a history of sports-related trauma, low functional status, limited knee flexion or extension, medial or lateral knee joint line tenderness, a click or pain noted with the McMurray test, and a positive Lachmann or anterior drawer test. ([Solomon, 2004](#)) Our conclusion is that operative treatment with complete repair of all torn structures produces the best overall knee function with better knee stability and patient satisfaction. In patients younger than 35, arthroscopic meniscal repair can preserve meniscal function, although the recovery time is longer compared to partial meniscectomy. Arthroscopy and meniscus surgery will not be as beneficial for older patients who are exhibiting signs of degenerative changes, possibly indicating osteoarthritis, and meniscectomy will not improve the OA. Meniscal repair is much more complicated than meniscal excision (meniscectomy). Some surgeons state in an operative report that they performed a meniscal repair when they may really mean a meniscectomy. A meniscus repair is a surgical procedure done to repair the damaged meniscus. This procedure can restore the normal anatomy of the knee, and has a better long-term prognosis when successful. However, the meniscus repair is a more significant surgery, the recovery is longer, and, because of limited blood supply to the meniscus, it is not always possible. A meniscectomy is a procedure to remove the torn portion of the meniscus. This procedure is far more commonly performed than a meniscus repair. Most meniscus tears cannot be treated by a repair. See also [Meniscal allograft transplantation](#). ([Harner, 2004](#)) ([Graf, 2004](#)) ([Wong, 2004](#)) ([Solomon-JAMA, 2001](#)) ([Chatain, 2003](#)) ([Chatain-Robinson, 2001](#)) ([Englund, 2004](#)) ([Englund, 2003](#)) ([Menetrey, 2002](#)) ([Pearse, 2003](#)) ([Roos, 2000](#)) ([Roos, 2001](#)) Arthroscopic debridement of meniscus tears and knees with low-grade osteoarthritis may have some utility, but it should not be used as a routine treatment for all patients with knee osteoarthritis. ([Siparsky, 2007](#)) Asymptomatic meniscal tears are common in older adults, based on studying MRI scans of the right knee of 991 randomly selected, ambulatory subjects. Incidental meniscal findings on MRI of the knee are common in the general population and increase with increasing age. Identifying a tear in a person with knee pain does not mean that the tear is the cause of the pain. ([Englund, 2008](#)) Arthroscopic meniscal repair results in good clinical and anatomic outcomes. ([Pujol, 2008](#)) Whether or not meniscal surgery is performed, meniscal tears in the knee increase the risk of developing osteoarthritis in middle age and elderly patients, and individuals with meniscal tear were 5.7 times more likely to develop knee osteoarthritis. ([Englund, 2009](#)) AHRQ Comparative Effectiveness Research concluded that arthroscopic lavage for osteoarthritis, with or without debridement, does not improve pain and function for people with OA of the knee. ([AHRQ, 2011](#)) The repair of meniscal tears is significantly improved when performed in conjunction with ACL reconstruction. ([Wasserstein, 2011](#))

Physical therapy vs. surgery: In older patients with degenerative tears and symptoms caused by osteoarthritis, PT/exercise may be an appropriate first option and it may be possible to reserve surgery for those who do not benefit from PT alone. A high quality RCT, the Meniscal Tear in Osteoarthritis Research (METEOR) trial, found similar outcomes from PT versus surgery for meniscal tears

The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

in older individuals. Researchers at seven major universities and orthopedic surgery centers around the U.S. assigned 351 people with arthritis and meniscus tears to get either surgery or physical therapy, nine sessions on average plus exercises to do at home. After six months, both groups had similar rates of functional improvement, and pain scores were also similar. While 30% of patients assigned to physical therapy wound up having surgery before the six months was up, often because they felt therapy wasn't helping them, they ended up the same as those who got surgery right away, as well as the rest of the physical therapy group who stuck with it and avoided having an operation. These results suggest that physical therapy may be an appropriate first option for many patients with osteoarthritis and meniscal tears and that it may be possible to reserve surgery for those who do not benefit from physical therapy alone. ([Katz, 2013](#)) Arthroscopic surgery for knee osteoarthritis offers no added benefit to optimized physical and medical therapy, according to the results of a single-center, RCT reported in the *New England Journal of Medicine*. The study, combined with other evidence, indicates that osteoarthritis of the knee (in the absence of a history and physical examination suggesting meniscal or other findings) is not an indication for arthroscopic surgery and indeed has been associated with inferior outcomes after arthroscopic knee surgery. However, osteoarthritis is not a contraindication to arthroscopic surgery, and arthroscopic surgery remains appropriate in patients with arthritis in specific situations in which osteoarthritis is not believed to be the primary cause of pain. ([Kirkley, 2008](#)) In this RCT, arthroscopic partial medial meniscectomy followed by supervised exercise was not superior to supervised exercise alone in terms of reduced knee pain, improved knee function and improved quality of life, after non-traumatic degenerative medial meniscal tear in ninety patients, mean age 56 years. ([Herrlin, 2007](#)) See also [Arthroscopic surgery for osteoarthritis](#).

ODG Indications for Surgery™ -- Meniscectomy:

Criteria for meniscectomy or meniscus repair (Suggest 2 symptoms and 2 signs to avoid scopes with lower yield, e.g. pain without other symptoms, posterior joint line tenderness that could just signify arthritis, MRI with degenerative tear that is often false positive). Physiologically younger and more active patients with traumatic injuries and mechanical symptoms (locking, blocking, catching, etc.) should undergo arthroscopy without PT.

1. Conservative Care: (Not required for locked/blocked knee.) Exercise/Physical therapy (supervised PT and/or home rehab exercises, if compliance is adequate). AND (Medication. OR Activity modification [eg, crutches and/or immobilizer].) PLUS

2. Subjective Clinical Findings (at least two): Joint pain. OR Swelling. OR Feeling of give way. OR Locking, clicking, or popping. PLUS

3. Objective Clinical Findings (at least two): Positive McMurray's sign. OR Joint line tenderness. OR Effusion. OR Limited range of motion. OR Locking, clicking, or popping. OR Crepitus. PLUS

4. Imaging Clinical Findings: (Not required for locked/blocked knee.) Meniscal tear on MRI (order MRI only after above criteria are met). ([Washington, 2003](#))

For average hospital LOS if criteria are met, see [Hospital length of stay](#) (LOS).

The DYLL REVIEW

We take the worry out of Peer Reviews

25 Highland Park Village #100-177 Dallas TX 75205

Phone: 888-950-4333 Fax: 888-9504-4443

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE, AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)