

ReviewTex, Inc.
1818 Mountjoy Drive
San Antonio, TX 78232
(phone) 210-598-9381 (fax) 210-598-9382
reviewtex@hotmail.com

Notice of Independent Review Decision

Date notice sent to all parties:

May 20, 2014

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Reconsideration – Cervical ESI 62310, 72275, 72020, 01936

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

Board Certified Anesthesiologist; Board Certified Pain Medicine

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

Upheld (Agree)

Provide a description of the review outcome that clearly states whether medical necessity exists for each of the health care services in dispute.

INFORMATION PROVIDED TO THE IRO FOR REVIEW:

PATIENT CLINICAL HISTORY [SUMMARY]:

The patient is a female who reported an injury to her neck when she developed severe interscapular pain described as intermittent. Clinical note dated indicated the patient rating her neck pain as 6/10. The patient performed level one physical therapy that alleviated some pain. The patient also utilized a TENS unit which provided some relief. Upon exam the patient demonstrated 45 degrees of cervical flexion, 50 degrees of extension, and 75 degrees of bilateral rotation. Tenderness was identified at the bilateral paravertebral musculature.

The patient demonstrated 5/5 strength throughout the upper extremities. No reflex deficits were identified. MRI of the cervical spine dated 03/12/14 revealed posterior central and paracentral mild disc bulge at C3-4 with a minimal posterior central disc bulge at C4-5 and C5-6. No evidence of spinal, neural canal, or neural foraminal stenosis was identified. Clinical note dated 03/18/14 indicated the patient continuing with neck pain. The patient continued to demonstrate 5/5 strength at the upper extremities with no identified reflex deficits. Utilization review dated 03/24/14 resulted in a denial as no objective information was submitted regarding failure of conservative treatment. Utilization review dated 04/07/14 resulted in denial as no significant neurocompressive pathology was identified by the submitted MRI.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS, AND CONCLUSIONS USED TO SUPPORT THE DECISION:

Clinical documentation indicated the patient complaining of neck pain. Epidural steroid injection would be indicated in the cervical spine provided that the patient meets specific criteria, including evidence confirmed by imaging studies that the neurocompressive findings and clinical evaluation confirming significant pathology. The submitted MRI revealed no significant neurocompressive findings. No information was submitted regarding reflex, strength, or sensation deficits in the appropriate distributions. Additionally, there is an indication that the patient has undergone some physical therapy. However, no therapy notes were submitted confirming completion of a full course of treatment. Given this, the request is not indicated. Therefore, as such, it is the opinion of this reviewer that the request for cervical epidural steroid injection is not recommended as medically necessary.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

MEDICAL JUDGEMENT, CLINICAL EXPERIENCE, AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
Epidural steroid injection (ESI)
Recommended as an option for treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy). See specific criteria for use below. In a recent Cochrane review, there was one study that reported improvement in pain and function at four weeks and also one year in individuals with chronic neck pain with radiation. (Peloso-Cochrane, 2006) (Peloso, 2005) Other reviews have reported moderate short-term and long-term evidence of success in managing cervical radiculopathy with interlaminar ESIs. (Stav, 1993) (Castagnera, 1994) Some have also reported moderate evidence of management of cervical nerve root pain using a transforaminal approach. (Bush, 1996) (Cyteval, 2004) A recent retrospective review of interlaminar cervical ESIs found that

approximately two-thirds of patients with symptomatic cervical radiculopathy from disc herniation were able to avoid surgery for up to 1 year with treatment. Success rate was improved with earlier injection (< 100 days from diagnosis). (Lin, 2006) There have been recent case reports of cerebellar infarct and brainstem herniation as well as spinal cord infarction after cervical transforaminal injection. (Beckman, 2006) (Ludwig, 2005) Quadriparesis with a cervical ESI at C6-7 has also been noted (Bose, 2005) and the American Society of Anesthesiologists Closed Claims Project database revealed 9 deaths or cases of brain injury after cervical ESI (1970-1999). (Fitzgibbon, 2004) These reports were in contrast to a retrospective review of 1,036 injections that showed that there were no catastrophic complications with the procedure. (Ma, 2005) The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular lumbosacral pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months, and there is insufficient evidence to make any recommendation for the use of epidural steroid injections to treat radicular cervical pain. (Armon, 2007) There is evidence for short-term symptomatic improvement of radicular symptoms with epidural or selective root injections with corticosteroids, but these treatments did not appear to decrease the rate of open surgery. (Haldeman, 2008) (Benyamin, 2009) Epidural steroid injections should be reserved for those who may otherwise undergo open surgery for nerve root compromise. (Bigos, 1999) Intramuscular injection of lidocaine for chronic mechanical neck disorders (MND) and intravenous injection of methylprednisolone for acute whiplash were effective treatments. There was limited evidence of effectiveness of epidural injection of methyl prednisolone and lidocaine for chronic MND with radicular findings. (Peloso-Cochrane, 2006) See the Low Back Chapter for more information and references.

Criteria for the use of Epidural steroid injections, therapeutic:

Note: The purpose of ESI is to reduce pain and inflammation, thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit.

- (1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing.
- (2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants).
- (3) Injections should be performed using fluoroscopy (live x-ray) for guidance
- (4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections.
- (5) No more than two nerve root levels should be injected using transforaminal blocks.
- (6) No more than one interlaminar level should be injected at one session.
- (7) In the therapeutic phase, repeat blocks should only be offered if there is at least 50% pain relief for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year.
- (8) Repeat injections should be based on continued objective documented pain and function response.

(9) Current research does not support a “series-of-three” injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections.

(10) It is currently not recommended to perform epidural blocks on the same day of treatment as facet blocks or stellate ganglion blocks or sympathetic blocks or trigger point injections as this may lead to improper diagnosis or unnecessary treatment.

(11) Cervical and lumbar epidural steroid injection should not be performed on the same day.

Criteria for the use of Epidural steroid injections, diagnostic:

To determine the level of radicular pain, in cases where diagnostic imaging is ambiguous, including the examples below:

(1) To help to evaluate a pain generator when physical signs and symptoms differ from that found on imaging studies;

(2) To help to determine pain generators when there is evidence of multi-level nerve root compression;

(3) To help to determine pain generators when clinical findings are suggestive of radiculopathy (e.g. dermatomal distribution), and imaging studies have suggestive cause for symptoms but are inconclusive;

(4) To help to identify the origin of pain in patients who have had previous spinal surgery.