ReviewTex, Inc.
1818 Mountjoy Drive San Antonio, TX 78232
(phone) 210-598-9381 (fax) 210-598-9382
reviewtex@hotmail.com

Notice of Independent Review Decision

Date notice sent to all parties:

May 27, 2014

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

MRI LUMBAR SPINE w/o Contrast

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION:

Board Certified PM&R; Board Certified Pain Medicine

REVIEW OUTCOME:

Upon independent review, the reviewer finds that the previous adverse determination/adverse determinations should be:

☒ Upheld (Agree)

Provide a description of the review outcome that clearly states whether medical necessity exists for each of the health care services in dispute.

INFORMATION PROVIDED TO THE IRO FOR REVIEW:

PATIENT CLINICAL HISTORY [SUMMARY]:

The patient is a male who reported an injury to his low back from an injury that occurred in xxxx. The clinical note dated xxxxx indicates the patient having previously undergone physical therapy as well as medications. A spinal cord stimulator trial in xxxxx resulted in very minimal benefit. The patient rated his pain at that time as 8-10/10. Upon exam, weakness, numbness, and tingling were identified in the lower extremities. The patient described the ongoing pain as a stabbing sensation. The procedural note dated 12/31/12 indicates the patient undergoing a C7-T1 epidural injection. The clinical note dated 05/06/13 indicates

the primary area of concern was in the cervical region at that time. The patient reported an 80% relief of pain following the most recent procedure. The clinical note dated 09/13/13 indicates the patient continuing with low back pain with radiating pain to the lower extremities. The patient also reported numbness and tingling. Weakness and numbness continued in both lower extremities, right greater than left. The note indicates the patient utilizing Hydrocodone for pain relief. The clinical note dated 12/09/13 indicates the patient complaining of spinous and facet tenderness in both lower extremities. The patient demonstrated range of motion limitations. The clinical note dated 03/06/14 indicates the patient continuing with the use of medications to address the low back complaints. The note indicates the patient having completed physical therapy in addition to the injections. The patient continued to rate the pain as 8-10/10. No strength deficits were identified. Reflexes were 2+ throughout. The clinical note dated 04/03/14 indicates the patient continuing with complaints of numbness and tingling in both lower extremities. 4/5 strength was identified in the left lower extremity. Decreased sensation was also identified in the L4 and L5 distributions on the right. The clinical note dated 05/02/14 indicates the patient continuing with radiating pain from the low back into the lower extremities. The patient had a positive straight leg raise bilaterally at that time.

The utilization review dated 02/03/14 resulted in a denial as the patient had recently undergone an MRI of the lumbar spine.

The utilization review dated 04/10/14 resulted in a denial for an MRI of the lumbar spine as there was no indication for a repeat study.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS, AND CONCLUSIONS USED TO SUPPORT THE DECISION:

The documentation indicates the patient complaining of low back pain with strength and sensation deficits identified in the lower extremities. An MRI of the lumbar spine is indicated provided the patient meets specific criteria to include the patient demonstrating significant changes involving radiculopathy. The clinical notes indicate the patient having recently undergone an MRI of the lumbar spine in June of 2013. There is an indication according to the clinical notes regarding significant findings in the L4-5 level. The recent clinical findings indicated the patient demonstrating radiculopathy that appeared to be in the L4 and L5 distributions. Thus, the present findings do not appear to be progressive in nature. Given these findings, the medical need for a repeat MRI study of the lumbar region has not been established. As such, it is the opinion of this reviewer that the request for an MRI of the lumbar spine without contrast is not recommended as medically necessary.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR

OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

☒ MEDICAL JUDGEMENT, CLINICAL EXPERIENCE, AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS

☒ ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES

MRIs (magnetic resonance imaging)

Recommended for indications below. MRI's are test of choice for patients with prior back surgery, but for uncomplicated low back pain, with radiculopathy, not recommended until after at least one month conservative therapy, sooner if severe or progressive neurologic deficit. Repeat MRI is not routinely recommended, and should be reserved for a significant change in symptoms and/or findings suggestive of significant pathology (eg, tumor, infection, fracture, neurocompression, recurrent disc herniation). (Bigos, 1999) (Mullin, 2000) (ACR, 2000) (AAN, 1994) (Aetna, 2004) (Airaksinen, 2006) (Chou, 2007) Magnetic resonance imaging has also become the mainstay in the evaluation of myelopathy. An important limitation of magnetic resonance imaging in the diagnosis of myelopathy is its high sensitivity. The ease with which the study depicts expansion and compression of the spinal cord in the myelopathic patient may lead to false positive examinations and inappropriately aggressive therapy if findings are interpreted incorrectly. (Seidenwurm, 2000) There is controversary over whether they result in higher costs compared to X-rays including all the treatment that continues after the more sensitive MRI reveals the usual insignificant disc bulges and herniations. (Jarvik-JAMA, 2003) In addition, the sensitivities of the only significant MRI parameters, disc height narrowing and anular tears, are poor, and these findings alone are of limited clinical importance. (Videman, 2003) Imaging studies are used most practically as confirmation studies once a working diagnosis is determined. MRI, although excellent at defining tumor, infection, and nerve compression, can be too sensitive with regard to degenerative disease findings and commonly displays pathology that is not responsible for the patient's symptoms. With low back pain, clinical judgment begins and ends with an understanding of a patient's life and circumstances as much as with their specific spinal pathology. (Carragee, 2004) Diagnostic imaging of the spine is associated with a high rate of abnormal findings in asymptomatic individuals. Herniated disk is found on magnetic resonance imaging in 9% to 76% of asymptomatic patients; bulging disks, in 20% to 81%; and degenerative disks, in 46% to 93%. (Kinkade, 2007) Baseline MRI findings do not predict future low back pain. (Borenstein, 2001) MRI findings may be preexisting. Many MRI findings (loss of disc signal, facet arthrosis, and end plate signal changes) may represent progressive age changes not associated with acute events. (Carragee, 2006) MRI abnormalities do not predict poor outcomes after conservative care for chronic low back pain patients. (Kleinstück, 2006) The new ACP/APS guideline as compared to the old AHCPR guideline is more forceful about the need to avoid specialized diagnostic imaging such as magnetic resonance imaging (MRI) without a clear rationale for doing so. (Shekelle, 2008) A new meta-analysis of randomized trials finds no benefit to routine lumbar imaging (radiography, MRI, or CT) for low back pain without indications of serious underlying conditions, and recommends that

clinicians should refrain from routine, immediate lumbar imaging in these patients. (Chou-Lancet, 2009) Despite guidelines recommending parsimonious imaging, use of lumbar MRI increased by 307% during a recent 12-year interval. When judged against guidelines, one-third to two-thirds of spinal computed tomography imaging and MRI may be inappropriate. (Deyo, 2009) As an alternative to MRI, a pain assessment tool named Standardized Evaluation of Pain (StEP), with six interview questions and ten physical tests, identified patients with radicular pain with high sensitivity (92%) and specificity (97%). The diagnostic accuracy of StEP exceeded that of a dedicated screening tool for neuropathic pain and spinal magnetic resonance imaging. (Scholz, 2009) Clinical quality-based incentives are associated with less advanced imaging, whereas satisfaction measures are associated with more rapid and advanced imaging, leading Richard Deyo, in the Archives of Internal Medicine to call the fascination with lumbar spine imaging an idolatry. (Pham, 2009) Primary care physicians are making a significant amount of inappropriate referrals for CT and MRI, according to new research published in the Journal of the American College of Radiology. There were high rates of inappropriate examinations for spinal CTs (53%), and for spinal MRIs (35%), including lumbar spine MRI for acute back pain without conservative therapy. (Lehnert, 2010) Degenerative changes in the thoracic spine on MRI were observed in approximately half of the subjects with no symptoms in this study. (Matsumoto, 2010) This large case series concluded that iatrogenic effects of early MRI are worse disability and increased medical costs and surgery, unrelated to severity. (Webster, 2010) Routine imaging for low back pain is not beneficial and may even be harmful, according to new guidelines from the American College of Physicians. Imaging is indicated only if they have severe progressive neurologic impairments or signs or symptoms indicating a serious or specific underlying condition, or if they are candidates for invasive interventions. Immediate imaging is recommended for patients with major risk factors for cancer, spinal infection, cauda equina syndrome, or severe or progressive neurologic deficits. Imaging after a trial of treatment is recommended for patients who have minor risk factors for cancer, inflammatory back disease, vertebral compression fracture, radiculopathy, or symptomatic spinal stenosis. Subsequent imaging should be based on new symptoms or changes in current symptoms. (Chou, 2011) The National Physicians Alliance compiled a "top 5" list of procedures in primary care that do little if anything to improve outcomes but excel at wasting limited healthcare dollars, and the list included routinely ordering diagnostic imaging for patients with low back pain, but with no warning flags, such as severe or progressive neurologic deficits, within the first 6 weeks. (Aguilar, 2011) Owning MRI equipment is a strongly correlated with patients receiving MRI scans, and having an MRI scan increases the probability of having surgery by 34%. (Shreibati, 2011) A considerable proportion of patients may be classified incorrectly by MRI for lumbar disc herniation, or for spinal stenosis. Pooled analysis resulted in a summary estimate of sensitivity of 75% and specificity of 77% for disc herniation. (Wassenaar, 2011) (Sigmundsson, 2011) Accurate terms are particularly important for classification of lumbar disc pathology from imaging. (Fardon, 2001) Among workers with LBP, early MRI is not associated with better health outcomes and is associated with increased likelihood of disability and its duration. (Graves, 2012) There is support for MRI, depending on symptoms and signs, to rule out serious pathology such as tumor, infection, fracture, and cauda equina syndrome. Patients with severe or progressive neurologic deficits from

lumbar disc herniation, or subjects with lumbar radiculopathy who do not respond to initial appropriate conservative care, are also candidates for lumbar MRI to evaluate potential for spinal interventions including injections or surgery. For unequivocal evidence of radiculopathy, see AMA Guides. (Andersson, 2000) MRI with and without contrast is best test for prior back surgery. (Davis, 2011) See also ACR Appropriateness Criteria™. See also Standing MRI.

Recent research: More than half of requests for MRI of the lumbar spine are ordered for indications considered inappropriate or of uncertain value, pointing to evidence of substantial overuse of lumbar spine MRI scans. For family physicians, only 34% of their MRI scans were considered appropriate vs 58% of those ordered by other specialties. On the other hand, the vast majority of MRIs ordered for headaches, 83%, were deemed appropriate. (Emery, 2013) This study casts doubt on the value of post-op spinal imaging for patients with sciatica, because it could not distinguish those with a favorable clinical outcome from those with persistent symptoms. Disk herniation was visible in 35% of patients with a favorable outcome and in 33% with an unfavorable outcome, and nerve root compression was present in 24% of those with a favorable outcome and in 26% of those with an unfavorable outcome. They concluded that the MRI scan does not have any discriminatory power at all. Irrelevant findings have the potential to frighten patients and initiate cascades of unnecessary testing or intervention, with occasional risks. The study showed that neither a herniated disk nor the presence of scar tissue on MRI was associated with patient outcome, but these findings may lead to unnecessary further imaging and surgery. (el Barzouhi, 2013) A JAMA article on worsening trends for low back treatment found that there was an escalation in the use of MRI or CT, from 7.2% in 1999 to 11.3% in 2010, while imaging in the acute care setting provides neither clinical nor psychological benefit to patients with routine back pain. The general feeling among physicians was that patients may equate getting MRIs with being synonymous with good medical care, which could drive doctors to try to improve patient satisfaction. (Mafi, 2013)

Indications for imaging -- Magnetic resonance imaging:

- Thoracic spine trauma: with neurological deficit
- Lumbar spine trauma: trauma, neurological deficit
- Lumbar spine trauma: seat belt (chance) fracture (If focal, radicular findings or other neurologic deficit)
- Uncomplicated low back pain, suspicion of cancer, infection, other “red flags”
- Uncomplicated low back pain, with radiculopathy, after at least 1 month conservative therapy, sooner if severe or progressive neurologic deficit.
- Uncomplicated low back pain, prior lumbar surgery
- Uncomplicated low back pain, cauda equina syndrome
- Myelopathy (neurological deficit related to the spinal cord), traumatic
- Myelopathy, painful
- Myelopathy, sudden onset
- Myelopathy, stepwise progressive
- Myelopathy, slowly progressive
- Myelopathy, infectious disease patient
- Myelopathy, oncology patient