



**MEDICAL EVALUATORS
OF TEXAS** ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

Notice of Independent Review Decision

DATE OF REVIEW: August 18, 2014

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

Psychotherapy x6 Sessions

**A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER
HEALTH CARE PROVIDER WHO REVIEWED THE DECISION**

This case was reviewed by a physician who holds a board certification in Physical Medicine and Rehabilitation and is currently licensed and practicing in the state of Texas.

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld (Agree)
- Overturned (Disagree)
- Partially Overturned (Agree in part/Disagree in part)

EMPLOYEE CLINICAL HISTORY [SUMMARY]:

This is a male who sustained an occupational incident on xx/xx/xx. The claimant sustained a brain injury and a skull fracture resulting in headaches, dizziness, blurred vision, hearing loss and depression. He has been treated with Cymbalta, Prozac, and Trazodone. The claimant also complained of facial weakness and has been evaluated for trigeminal neuralgia. MRI of the brain on 09/27/2010 showed nonspecific white matter changes. MRI of the brain on 03/25/2013 showed nonspecific white matter changes and left parietal lobe chronic hemorrhage that was stable from a prior study. In 2013, he had been authorized for 24 sessions of individual psychotherapy. The claimant has seen on a regular basis and states his condition is unchanged. On 07/10/2014, it was noted that his depression had worsened and is wanting to restart counseling. Psychiatric exam showed appropriate affect and demeanor, normal speech pattern, and grossly normal memory. He was diagnosed with moderate depression. He was requested psychotherapy x6 sessions which has been non-certified as there is no sufficient documentation pertaining to the efficacy from previous therapy sessions.



MEDICAL EVALUATORS OF TEXAS ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

The claimant received 24 sessions of psychotherapy in 2013 from which he did not show any improvement. Despite reported worsening of depression per the claimant, his medical evaluation did not exhibit any specific abnormalities. The requested 6 sessions of psychotherapy is not recommended at this time as there is no documentable improvement from previous treatments or abnormalities documented during physical exam. The claimant does not meet necessary requirements as per the ODG guidelines noted below.

ODG – Mental illness & Stress

Cognitive therapy for depression

Recommended. Cognitive behavior therapy for depression is recommended based on meta-analyses that compare its use with pharmaceuticals. Cognitive behavior therapy fared as well as antidepressant medication with severely depressed outpatients in four major comparisons. Effects may be longer lasting (80% relapse rate with antidepressants versus 25% with psychotherapy). (Paykel, 2006) (Bockting, 2006) (DeRubeis, 1999) (Goldapple, 2004) It also fared well in a meta-analysis comparing 78 clinical trials from 1977 -1996. (Gloaguen, 1998) In another study, it was found that combined therapy (antidepressant plus psychotherapy) was found to be more effective than psychotherapy alone. (Thase, 1997) A recent high quality study concluded that a substantial number of adequately treated patients did not respond to antidepressant therapy. (Corey-Lisle, 2004) A recent meta-analysis concluded that psychological treatment combined with antidepressant therapy is associated with a higher improvement rate than drug treatment alone. In longer therapies, the addition of psychotherapy helps to keep patients in treatment. (Pampallona, 2004) For panic disorder, cognitive behavior therapy is more effective and more cost-effective than medication. (Royal Australian, 2003) The gold standard for the evidence-based treatment of MDD is a combination of medication (antidepressants) and psychotherapy. The primary forms of psychotherapy that have been most studied through research are: Cognitive Behavioral Therapy and Interpersonal Therapy. (Warren, 2005) Delivering cognitive behavioral therapy (CBT) by telephone is as effective as delivering it face-to-face in the short term, and telephone therapy is safe and has a higher patient retention rate. The attrition rate from psychotherapy can exceed 50% due to time constraints, lack of available and accessible services, transportation problems, and cost. Significantly fewer participants receiving telephone CBT discontinued their therapy than did those receiving face-to-face CBT. Both treatment groups showed significant improvement in depression, and there were no significant treatment differences when measured at posttreatment between telephone and face-to-face CBT. However, face-to-face CBT was significantly superior to telephone CBT during the follow-up period. The RCT used 18 sessions of either telephone CBT or face-to-face CBT. (Mohr, 2012) Maintenance cognitive-behavioral therapy (CBT) to prevent recurrent depression is most effective in patients at highest risk for relapse, defined as those with 5 or more previous



MEDICAL EVALUATORS OF TEXAS ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

depressive episodes. For individuals at more moderate risk for recurrence (fewer than 5 prior episodes), structured patient psychoeducation may be equally effective. High-risk patients in particular may benefit from specific elements of maintenance CBT by reducing cognitive vulnerability factors for recurrent depression, such as ruminating, negative attributions and memories, and dysfunctional beliefs, or by maintaining positive emotions when experiencing stress. (Stangier, 2013) Studies show that a 4 to 6 session trial should be sufficient to provide evidence of symptom improvement, but functioning and quality of life indices do not change as markedly within a short duration of psychotherapy as do symptom-based outcome measures. (Crits-Christoph, 2001) See Number of psychotherapy sessions for more information. See also Bibliotherapy; Computer-assisted cognitive therapy. Psychotherapy visits are generally separate from physical therapy visits.

ODG Psychotherapy Guidelines:

- Up to 13-20 visits over 7-20 weeks (individual sessions), if progress is being made. (The provider should evaluate symptom improvement during the process, so treatment failures can be identified early and alternative treatment strategies can be pursued if appropriate.)
- In cases of severe Major Depression or PTSD, up to 50 sessions if progress is being made.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES**



**MEDICAL EVALUATORS
OF T E X A S** ASO, L.L.C.

1225 North Loop West • Suite 1055 • Houston, TX 77008
800-845-8982 FAX: 713-583-5943

- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR**
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS**
- TEXAS TACADA GUIDELINES**
- TMF SCREENING CRITERIA MANUAL**
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)**
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)**