



**MEDICAL EVALUATORS
OF T E X A S ASO, LLC.**

2211 West 34th St. • Houston, TX 77018
800-845-8982 FAX: 713-583-5943

Notice of Independent Review Decision

December 4, 2014

IRO CASE #:

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE

Work Hardening Program x 80 hours/units

**A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER
HEALTH CARE PROVIDER WHO REVIEWED THE DECISION**

This case was reviewed by a chiropractor who is currently licensed and practicing in the state of Texas.

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse determination/adverse determinations should be:

- Upheld** (Agree)
 Overtured (Disagree)
 Partially Overtured (Agree in part/Disagree in part)

EMPLOYEE CLINICAL HISTORY [SUMMARY]:

This patient is a male who was injured on xx/xx/xx. Treatment history includes medications, physical therapy and psychotherapy. She had FCE dated 09/30/2014 documented that the evaluatee is unable to perform their regular job duties at this time, the evaluatee required PDL is heavy, however their current physical performance is medium , and that the evaluatee was not able to demonstrate the ability to perform several key functions crucial to the safe performance of their normal work duties.

Follow-Up note dated 10/22/2014 documented the patient complaints of pain and weakness in the left hand, wrist, elbow, and shoulder. Examination of the hand appeared to be completely normal, the patient did endorse tenderness with examination of the left hand, wrist, elbow, and shoulder. ROM and strength were all normal. The patient was diagnosed with strain of the left upper extremity; shoulder, elbow, wrist, and hand and was recommended chronic pain management program or work hardening program.



**MEDICAL EVALUATORS
OF TEXAS** ASO, LLC.

2211 West 34th St. • Houston, TX 77018
800-845-8982 FAX: 713-583-5943

The request was denied twice by UniMed on 10/31/2014 and 11/18/2014 as the physical exam was completely normal.

**ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS,
FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.**

This male sustained superficial injury to his left hand. The patient has undergone beyond reasonable treatment with psychological and physical rehabilitation. The psychological treatment provided minimal benefit. Additional psychological therapy will, most likely, fail. The recent examination by his treating doctor revealed “full ROM and normal strength” on 10/22/2014. Therefore, there is no reason to receive any additional physical rehabilitation since the ROM and strength are normal. FCE revealed his required PDL is heavy but the current physical performance is medium. Comparing the results of the FCE and the examination on 10/22/2014, it revealed improvement to “normal strength” status. Therefore, the FCE should not be considered as supporting documents for the request of work hardening program. There is no documentation of any physical deficits in his hand to prevent him from returning to work or seeking other employment. There is limited evidence to support that the work hardening program would be any benefit in this situation according to ODG. It is unclear why he will not benefit from transition to a home exercise program.

Therefore, I would not recommend the request of work hardening program at this time.

ODG – Forearm, Wrist, & Hand (Acute & Chronic)

Work conditioning, work hardening

Recommended as an option, depending on the availability of quality programs, and should be specific for the job individual is going to return to. (Schonstein-Cochrane, 2003) There is limited literature support for multidisciplinary treatment and work hardening for the neck, hip, knee, shoulder and forearm. (Karjalainen, 2003) Work Conditioning should restore the client’s physical capacity and function. Work Hardening should be work simulation and not just therapeutic exercise, plus there should also be psychological support. Work Hardening is an interdisciplinary, individualized, job specific program of activity with the goal of return to work. Work Hardening programs use real or simulated work tasks and progressively graded conditioning exercises that are based on the individual’s measured tolerances. (CARF, 2006) (Washington, 2006) There is limited, but high quality, evidence that multi-disciplinary rehabilitation for non-specific musculoskeletal arm pain was beneficial for those workers absent from work for at least four weeks. (Dick, 2010) For more information and references, see the Low Back Chapter. The Low Back WH & WC Criteria are copied below.

Criteria for admission to a Work Hardening (WH) Program:

- (1) Prescription: The program has been recommended by a physician or nurse case manager, and a prescription has been provided.
- (2) Screening Documentation: Approval of the program should include evidence of a screening evaluation. This multidisciplinary examination should include the following



MEDICAL EVALUATORS OF TEXAS ASO, LLC.

2211 West 34th St. • Houston, TX 77018
800-845-8982 FAX: 713-583-5943

components: (a) History including demographic information, date and description of injury, history of previous injury, diagnosis/diagnoses, work status before the injury, work status after the injury, history of treatment for the injury (including medications), history of previous injury, current employability, future employability, and time off work; (b) Review of systems including other non work-related medical conditions; (c) Documentation of musculoskeletal, cardiovascular, vocational, motivational, behavioral, and cognitive status by a physician, chiropractor, or physical and/or occupational therapist (and/or assistants); (d) Diagnostic interview with a mental health provider; (e) Determination of safety issues and accommodation at the place of work injury. Screening should include adequate testing to determine if the patient has attitudinal and/or behavioral issues that are appropriately addressed in a multidisciplinary work hardening program. The testing should also be intensive enough to provide evidence that there are no psychosocial or significant pain behaviors that should be addressed in other types of programs, or will likely prevent successful participation and return-to-employment after completion of a work hardening program. Development of the patient's program should reflect this assessment.

(3) Job demands: A work-related musculoskeletal deficit has been identified with the addition of evidence of physical, functional, behavioral, and/or vocational deficits that preclude ability to safely achieve current job demands. These job demands are generally reported in the medium or higher demand level (i.e., not clerical/sedentary work). There should generally be evidence of a valid mismatch between documented, specific essential job tasks and the patient's ability to perform these required tasks (as limited by the work injury and associated deficits).

(4) Functional capacity evaluations (FCEs): A valid FCE should be performed, administered and interpreted by a licensed medical professional. The results should indicate consistency with maximal effort, and demonstrate capacities below an employer verified physical demands analysis (PDA). Inconsistencies and/or indication that the patient has performed below maximal effort should be addressed prior to treatment in these programs.

(5) Previous PT: There is evidence of treatment with an adequate trial of active physical rehabilitation with improvement followed by plateau, with evidence of no likely benefit from continuation of this previous treatment. Passive physical medicine modalities are not indicated for use in any of these approaches.

(6) Rule out surgery: The patient is not a candidate for whom surgery, injections, or other treatments would clearly be warranted to improve function (including further diagnostic evaluation in anticipation of surgery).

(7) Healing: Physical and medical recovery sufficient to allow for progressive reactivation and participation for a minimum of 4 hours a day for three to five days a week.

(8) Other contraindications: There is no evidence of other medical, behavioral, or other comorbid conditions (including those that are non work-related) that prohibits participation in the program or contradicts successful return-to-work upon program completion.

(9) RTW plan: A specific defined return-to-work goal or job plan has been established, communicated and documented. The ideal situation is that there is a plan agreed to by



**MEDICAL EVALUATORS
OF TEXAS ASO, LLC.**

2211 West 34th St. • Houston, TX 77018
800-845-8982 FAX: 713-583-5943

the employer and employee. The work goal to which the employee should return must have demands that exceed the claimant's current validated abilities.

(10) Drug problems: There should be documentation that the claimant's medication regimen will not prohibit them from returning to work (either at their previous job or new employment). If this is the case, other treatment options may be required, for example a program focused on detoxification.

(11) Program documentation: The assessment and resultant treatment should be documented and be available to the employer, insurer, and other providers. There should be documentation of the proposed benefit from the program (including functional, vocational, and psychological improvements) and the plans to undertake this improvement. The assessment should indicate that the program providers are familiar with the expectations of the planned job, including skills necessary. Evidence of this may include site visitation, videotapes or functional job descriptions.

(12) Further mental health evaluation: Based on the initial screening, further evaluation by a mental health professional may be recommended. The results of this evaluation may suggest that treatment options other than these approaches may be required, and all screening evaluation information should be documented prior to further treatment planning.

(13) Supervision: Supervision is recommended under a physician, chiropractor, occupational therapist, or physical therapist with the appropriate education, training and experience. This clinician should provide on-site supervision of daily activities, and participate in the initial and final evaluations. They should design the treatment plan and be in charge of changes required. They are also in charge of direction of the staff.

(14) Trial: Treatment is not supported for longer than 1-2 weeks without evidence of patient compliance and demonstrated significant gains as documented by subjective and objective improvement in functional abilities. Outcomes should be presented that reflect the goals proposed upon entry, including those specifically addressing deficits identified in the screening procedure. A summary of the patient's physical and functional activities performed in the program should be included as an assessment of progress.

(15) Concurrently working: The patient who has been released to work with specific restrictions may participate in the program while concurrently working in a restricted capacity, but the total number of daily hours should not exceed 8 per day while in treatment.

(16) Conferences: There should be evidence of routine staff conferencing regarding progress and plans for discharge. Daily treatment activity and response should be documented.

(17) Voc rehab: Vocational consultation should be available if this is indicated as a significant barrier. This would be required if the patient has no job to return to.

(18) Post-injury cap: The worker must be no more than 2 years past date of injury. Workers that have not returned to work by two-years post injury generally do not improve from intensive work hardening programs. If the worker is greater than one-year post injury a comprehensive multidisciplinary program may be warranted if there is clinical



MEDICAL EVALUATORS OF T E X A S ASO, LLC.

2211 West 34th St. • Houston, TX 77018
800-845-8982 FAX: 713-583-5943

suggestion of psychological barrier to recovery (but these more complex programs may also be justified as early as 8-12 weeks, see Chronic pain programs).

(19) Program timelines: These approaches are highly variable in intensity, frequency and duration. APTA, AOTA and utilization guidelines for individual jurisdictions may be inconsistent. In general, the recommendations for use of such programs will fall within the following ranges: These approaches are necessarily intensive with highly variable treatment days ranging from 4-8 hours with treatment ranging from 3-5 visits per week. The entirety of this treatment should not exceed 20 full-day visits over 4 weeks, or no more than 160 hours (allowing for part-day sessions if required by part-time work, etc., over a longer number of weeks). A reassessment after 1-2 weeks should be made to determine whether completion of the chosen approach is appropriate, or whether treatment of greater intensity is required.

(20) Discharge documentation: At the time of discharge the referral source and other predetermined entities should be notified. This may include the employer and the insurer. There should be evidence documented of the clinical and functional status, recommendations for return to work, and recommendations for follow-up services. Patient attendance and progress should be documented including the reason(s) for termination including successful program completion or failure. This would include noncompliance, declining further services, or limited potential to benefit. There should also be documentation if the patient is unable to participate due to underlying medical conditions including substance dependence.

(21) Repetition: Upon completion of a rehabilitation program (e.g., work conditioning, work hardening, outpatient medical rehabilitation, or chronic pain/functional restoration program) neither re-enrollment in nor repetition of the same or similar rehabilitation program is medically warranted for the same condition or injury.

ODG Work Conditioning (WC) Physical Therapy Guidelines

WC amounts to an additional series of intensive physical therapy (PT) visits required beyond a normal course of PT, primarily for exercise training/supervision (and would be contraindicated if there are already significant psychosocial, drug or attitudinal barriers to recovery not addressed by these programs). See also Physical therapy for general PT guidelines. WC visits will typically be more intensive than regular PT visits, lasting 2 or 3 times as long. And, as with all physical therapy programs, Work Conditioning participation does not preclude concurrently being at work.

Timelines: 10 visits over 4 weeks, equivalent to up to 30 hours.



**MEDICAL EVALUATORS
OF T E X A S** ASO,LLC.

2211 West 34th St. • Houston, TX 77018
800-845-8982 FAX: 713-583-5943

**A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER
CLINICAL BASIS USED TO MAKE THE DECISION:**

- ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE
- AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
- DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
- EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
- INTERQUAL CRITERIA
- MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
- MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
- MILLIMAN CARE GUIDELINES
- ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES**
- PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
- TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
- TEXAS TACADA GUIDELINES
- TMF SCREENING CRITERIA MANUAL
- PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
- OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)