

DECISION AND ORDER

This case is decided pursuant to Chapter 410 of the Texas Workers' Compensation Act and Rules of the Division of Workers' Compensation adopted thereunder.

ISSUES

A contested case hearing was held on May 7, 2009, to decide the following disputed issue:

1. Is the preponderance of the evidence contrary to the decision of the IRO that the claimant is not entitled to 8 sessions of physical therapy to include aquatic therapy for the compensable injury of _____?

PARTIES PRESENT

Petitioner/Claimant appeared and was assisted by TT, ombudsman. Respondent/Carrier appeared and was represented by RR, attorney.

BACKGROUND INFORMATION

The petitioner/claimant sustained a compensable injury on _____ to the lumbar spine. The claimant underwent physical therapy for over three years and underwent epidural steroid injections. From the date of injury through the date of this hearing, the claimant has not missed time off from work for the compensable injury except for a few days when undergoing the epidural steroid injections. The claimant noted that sometime in January 2008, she helped move items for her employer and has since had the need for further therapy. A lumbar spine MRI undertaken on 5/22/08 revealed claimant suffers from degenerative disc disease. The parties noted that the physical therapy being requested is comprised entirely of aquatic therapy. The IRO decision stated that there must first be a justification for additional physical therapy and only then does one proceed to see if the justification for aquatic therapy has been met. The IRO decision noted that physical therapy is utilized initially and then an individual is weaned off of it onto home based exercises, that the claimant's problem is 8 years old, that formal therapy is utilized for people with fear of movement avoidance, which was not documented here, and directed to relatively acute or subacute problems, which is not the case. The IRO decision reasoned that as the claimant does not meet the criteria for physical therapy, the claimant also does not meet the criteria for aquatic therapy. The requesting doctor, Dr. YK, noted in his reports that the request was made for the purpose of improving the claimant's conditioning, strengthening and diminishment of pain. He noted that there was no specific incident that caused the need for the request, but rather, it was based upon his assessment of the claimant's current physical limitations of severe deconditioning, being overweight and significant abnormality in the lumbosacral spine.

Texas Labor Code Section 408.021 provides that an employee who sustains a compensable injury is entitled to all health care reasonably required by the nature of the injury as and when needed. Health care reasonably required is further defined in Texas Labor Code Section 401.011

(22a) as health care that is clinically appropriate and considered effective for the injured employee's injury and provided in accordance with best practices consistent with evidence based medicine or, if evidence based medicine is not available, then generally accepted standards of medical practice recognized in the medical community. Health care under the Texas Workers' Compensation system must be consistent with evidence based medicine if that evidence is available. Evidence based medicine is further defined in Texas Labor Code Section 401.011 (18a) to be the use of the current best quality scientific and medical evidence formulated from credible scientific studies, including peer-reviewed medical literature and other current scientifically based texts and treatment and practice guidelines.

In accordance with the above statutory guidance, the Division of Workers' Compensation has adopted treatment guidelines by Division Rule 137.100. This rule directs health care providers to provide treatment in accordance with the current edition of the Official Disability Guidelines (ODG), and such treatment is presumed to be health care reasonably required as defined in the Texas Labor Code. Thus, the focus of any health care dispute starts with the health care set out in the Official Disability Guidelines (ODG).

Aquatic Therapy

Recommended as an optional form of exercise therapy, where available, as an alternative to land-based physical therapy. Aquatic therapy (including swimming) can minimize the effects of gravity, so it is specifically recommended where reduced weight bearing is desirable, for example extreme obesity. There may be advantages to weightless running in back pain recovery. (Ariyoshi, 1999) (Burns, 2001) For recommendations on the number of supervised visits, see Physical therapy.

Physical therapy (PT)

Recommended. There is strong evidence that physical methods, including exercise and return to normal activities, have the best long-term outcome in employees with low back pain. See also Exercise. Direction from physical and occupational therapy providers can play a role in this, with the evidence supporting active therapy and not extensive use of passive modalities. The most effective strategy may be delivering individually designed exercise programs in a supervised format (for example, home exercises with regular therapist follow-up), encouraging adherence to achieve high dosage, and stretching and muscle-strengthening exercises seem to be the most effective types of exercises for treating chronic low back pain. (Hayden, 2005) Studies also suggest benefit from early use of aggressive physical therapy (“sports medicine model”), training in exercises for home use, and a functional restoration program, including intensive physical training, occupational therapy, and psychological support. (Zigenfus, 2000) (Linz, 2002) (Cherkin-NEJM, 1998) (Rainville, 2002) Successful outcomes depend on a functional restoration program, including intensive physical training, versus extensive use of passive modalities. (Mannion, 2001) (Jousset, 2004) (Rainville, 2004) (Airaksinen, 2006) One clinical trial found both effective, but chiropractic was slightly more favorable for acute back pain and physical therapy for chronic cases. (Skargren, 1998) A spinal stabilization program is more effective than standard physical therapy sessions, in which no exercises are prescribed. With regard to manual therapy, this approach may be the most common physical therapy modality for chronic low back disorder, and it may be appropriate as a pain reducing modality, but it should not be used as an isolated

modality because it does not concomitantly reduce disability, handicap, or improve quality of life. (Goldby-Spine, 2006) Better symptom relief is achieved with directional preference exercise. (Long, 2004) As compared with no therapy, physical therapy (up to 20 sessions over 12 weeks) following disc herniation surgery was effective. Because of the limited benefits of physical therapy relative to "sham" therapy (massage), it is open to question whether this treatment acts primarily physiologically, but psychological factors may contribute substantially to the benefits observed. (Erdogmus, 2007) See also specific physical therapy modalities, as well as Exercise; Work conditioning; Lumbar extension exercise equipment; McKenzie method; & Stretching. [Physical therapy is the treatment of a disease or injury by the use of therapeutic exercise and other interventions that focus on improving posture, locomotion, strength, endurance, balance, coordination, joint mobility, flexibility, activities of daily living and alleviating pain. (BlueCross BlueShield, 2005) As for visits with any medical provider, physical therapy treatment does not preclude an employee from being at work when not visiting the medical provider, although time off may be required for the visit.]

Active Treatment versus Passive Modalities: The use of active treatment modalities instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with acute low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) The most commonly used active treatment modality is Therapeutic exercises (97110), but other active therapies may be recommended as well, including Neuromuscular reeducation (97112), Manual therapy (97140), and Therapeutic activities/exercises (97530).

Patient Selection Criteria: Multiple studies have shown that patients with a high level of fear-avoidance do much better in a supervised physical therapy exercise program, and patients with low fear-avoidance do better following a self-directed exercise program. When using the Fear-Avoidance Beliefs Questionnaire (FABQ), scores greater than 34 predicted success with PT supervised care. (Fritz, 2001) (Fritz, 2002) (George, 2003) (Klaber, 2004) (Riipinen, 2005) (Hicks, 2005) Without proper patient selection, routine physical therapy may be no more effective than one session of assessment and advice from a physical therapist. (Frost, 2004) Patients exhibiting the centralization phenomenon during lumbar range of motion testing should be treated with the specific exercises (flexion or extension) that promote centralization of symptoms. When findings from the patient's history or physical examination are associated with clinical instability, they should be treated with a trunk strengthening and stabilization exercise program. (Fritz-Spine, 2003)

Post Epidural Steroid Injections: ESIs are currently recommended as a possible option for short-term treatment of radicular pain (sciatica), defined as pain in dermatomal distribution with corroborative findings of radiculopathy. The general goal of physical therapy during the acute/subacute phase of injury is to decrease guarding, maintain motion, and decrease pain and inflammation. Progression of rehabilitation to a more advanced program of stabilization occurs in the maintenance phase once pain is controlled. There is little evidence-based research

that addresses the use of physical therapy post ESIs, but it appears that most randomized controlled trials have utilized an ongoing, home directed program post injection. Based on current literature, the only need for further physical therapy treatment post ESI would be to emphasize the home exercise program, and this requirement would generally be included in the currently suggested maximum visits for the underlying condition, or at least not require more than 2 additional visits to reinforce the home exercise program. ESIs have been found to have limited effectiveness for treatment of chronic pain. The claimant should continue to follow a home exercise program post injection. (Luijsterburg, 2007) (Luijsterburg2, 2007) (Price, 2005) (Vad, 2002) (Smeal, 2004)

ODG Physical Therapy Guidelines –

Allow for fading of treatment frequency (from up to 3 or more visits per week to 1 or less), plus active self-directed home PT. Also see other general guidelines that apply to all conditions under Physical Therapy in the ODG Preface, including assessment after a "six-visit clinical trial".

Lumbar sprains and strains (ICD9 847.2):

10 visits over 8 weeks

Sprains and strains of unspecified parts of back (ICD9 847):

10 visits over 5 weeks

Sprains and strains of sacroiliac region (ICD9 846):

Medical treatment: 10 visits over 8 weeks

Lumbago; Backache, unspecified (ICD9 724.2; 724.5):

9 visits over 8 weeks

Intervertebral disc disorders without myelopathy (ICD9 722.1; 722.2; 722.5; 722.6; 722.8):

Medical treatment: 10 visits over 8 weeks

Post-injection treatment: 1-2 visits over 1 week

Post-surgical treatment (discectomy/laminectomy): 16 visits over 8 weeks

Post-surgical treatment (arthroplasty): 26 visits over 16 weeks

Post-surgical treatment (fusion, after graft maturity): 34 visits over 16 weeks

Intervertebral disc disorder with myelopathy (ICD9 722.7)

Medical treatment: 10 visits over 8 weeks

Post-surgical treatment: 48 visits over 18 weeks

Spinal stenosis (ICD9 724.0):

10 visits over 8 weeks

See 722.1 for post-surgical visits

Sciatica; Thoracic/lumbosacral neuritis/radiculitis, unspecified (ICD9 724.3; 724.4):

10-12 visits over 8 weeks

See 722.1 for post-surgical visits

Curvature of spine (ICD9 737)

12 visits over 10 weeks

See 722.1 for post-surgical visits

Fracture of vertebral column without spinal cord injury (ICD9 805):

Medical treatment: 8 visits over 10 weeks

Post-surgical treatment: 34 visits over 16 weeks

Fracture of vertebral column with spinal cord injury (ICD9 806):

Medical treatment: 8 visits over 10 weeks

Post-surgical treatment: 48 visits over 18 weeks

Work conditioning (See also Procedure Summary entry):

10 visits over 8 weeks

The requesting doctor did not present testimony and the claimant relied upon her testimony along with the medical reports in evidence for support of her request. The claimant must first show that she has met the necessary criteria for physical therapy. Here, the evidence revealed that the claimant underwent physical therapy soon after her compensable injury for quite some time and then moved into a home based exercise program. Her epidural steroid injections were undertaken somewhere around that time period. Even the requesting doctor notes that his request is not based upon the undertaking of epidural injections or any acute or subacute trauma. Although the claimant's weight is cited on a few medical records from the date of injury through the current time, the records does not cite any body mass index or indication that it is considered overweight except for one conclusory statement from the requesting doctor. As the claimant must show that she has presented evidence based medicine for her request of physical therapy and has failed to do so, she also in turn has not met the required ODG criteria for aquatic therapy. Based on the evidence presented, the claimant failed to provide evidence based medicine sufficient to contradict the determination of the IRO and the preponderance of the credible evidence is not contrary to the decision of the IRO.

Even though all the evidence presented was not discussed, it was considered. The Findings of Fact and Conclusions of Law are based on all of the evidence presented.

FINDINGS OF FACT

1. The parties stipulated to the following facts:
 - A. Venue is proper in the (City) Field Office of the Texas Department of Insurance, Division of Workers' Compensation.
 - B. On _____, Claimant was the employee of (SELF-INSURED EMPLOYER).
2. Carrier delivered to Claimant and Provider a single document stating the true corporate name of Carrier, and the name and street address of Carrier's registered agent, which document was admitted into evidence as Hearing Officer's Exhibit Number 2.
3. Eight sessions of physical therapy to include aquatic therapy is not health care reasonably required for the compensable injury of _____.

CONCLUSIONS OF LAW

1. The Texas Department of Insurance, Division of Workers' Compensation, has jurisdiction to hear this case.
2. Venue is proper in the (City) Field Office.
3. The preponderance of the evidence is not contrary to the decision of the IRO that 8 sessions of physical therapy to include aquatic therapy is not health care reasonably required for the compensable injury of _____.

DECISION

Claimant is not entitled to 8 sessions of physical therapy to include aquatic therapy for the compensable injury of _____.

ORDER

Carrier is not liable for the benefits at issue in this hearing. Claimant remains entitled to medical benefits for the compensable injury in accordance with §408.021.

The true corporate name of the insurance carrier is **(SELF-INSURED EMPLOYER)** and the name and address of its registered agent for service of process is

**CSC
(STREET ADDRESS), (SUITE)
(CITY), TEXAS (ZIP CODE)**

Signed this 8th day of May, 2009.

Virginia Rodríguez-Gómez
Hearing Officer