

**DECISION AND ORDER**

This case is decided pursuant to Chapter 410 of the Texas Workers' Compensation Act and Rules of the Division of Workers' Compensation adopted thereunder.

**ISSUES**

A contested case hearing was held on May 18, 2010, to decide the following disputed issue:

1. Is the preponderance of the evidence contrary to the decision of the Independent Review Organization (IRO) that a low pressure lumbar discogram with CT at L3-4, L4-5 and L5-S1 is not reasonably required health care for the compensable injury of \_\_\_\_\_?

**PARTIES PRESENT**

Claimant appeared and was assisted by RH, ombudsman. Petitioner/Provider appeared as a witness only without representation. Respondent/Carrier appeared and was represented by JB, adjuster.

**BACKGROUND INFORMATION**

Claimant sustained a compensable injury when the bus she was riding was struck by a train operating in the plant. Her treating doctor is Dr. B, MD, a principal in (Healthcare Provider), Petitioner. Dr. B requested preauthorization to perform a three-level discogram at L3-4, L4-5 and L5-S1.

Dr. B's request was first reviewed by Dr. W, MD, a utilization review agent (URA) for Carrier. Dr. W found that there was no clear documentation of neurologic deficit, muscle atrophy, protective muscle spasm or other abnormality. He went on to say that he found no evidence of a large disc herniation, progressive loss of function, or "anything other than subjective back pain complaints" in the medical records. He found no medical necessity for the requested discogram. The initial denial was appealed. The second URA was Dr. B (2), DO, an orthopedic surgeon. Dr. B (2) concurred with Dr. W that the low pressure lumbar discogram was not medically necessary. Review by an Independent Review Organization (IRO) was requested.

Texas Labor Code Section 408.021 provides that an employee who sustains a compensable injury is entitled to all health care reasonably required by the nature of the injury as and when needed. Health care reasonably required is further defined in Texas Labor Code Section 401.011 (22a) as health care that is clinically appropriate and considered effective for the injured employee's injury and provided in accordance with best practices consistent with evidence based medicine or, if evidence based medicine is not available, then generally accepted standards of medical practice recognized in the medical community. Health care under the Texas Workers' Compensation system must be consistent with evidence based medicine if that evidence is available. Evidence based medicine is defined in Texas Labor Code Section 401.011 (18a) to be the use of the current best quality scientific and medical evidence formulated from credible

scientific studies, including peer-reviewed medical literature and other current scientifically based texts and treatment and practice guidelines. The Commissioner of the Division of Workers' compensation is required to adopt treatment guidelines that are evidence-based, scientifically valid, outcome-focused and designed to reduce excessive or inappropriate medical care while safeguarding necessary medical care. Texas Labor Code Section 413.011(e). Medical services consistent with the medical policies and fee guidelines adopted by the commissioner are presumed reasonable in accordance with Texas Labor Code Section 413.017(1).

In accordance with the above statutory guidance, the Division of Workers' Compensation has adopted treatment guidelines by Division Rule 137.100. This rule directs health care providers to provide treatment in accordance with the current edition of the ODG, and such treatment is presumed to be health care reasonably required as defined in the Texas Labor Code. Thus, the focus of any health care dispute starts with the health care set out in the ODG.

The Low Back section of the ODG has the following recommendation regarding discography:

Not recommended. In the past, discography has been used as part of the preoperative evaluation of patients for consideration of surgical intervention for lower back pain. However, the conclusions of recent, high quality studies on discography have significantly questioned the use of discography results as a preoperative indication for either IDET or spinal fusion. These studies have suggested that reproduction of the patient's specific back complaints on injection of one or more discs (concordance of symptoms) is of limited diagnostic value. (Pain production was found to be common in non-back pain patients, pain reproduction was found to be inaccurate in many patients with chronic back pain and abnormal psychosocial testing, and in this latter patient type, the test itself was sometimes found to produce significant symptoms in non-back pain controls more than a year after testing.) Also, the findings of discography have not been shown to consistently correlate well with the finding of a High Intensity Zone (HIZ) on MRI. Discography may be justified if the decision has already been made to do a spinal fusion, and a negative discogram could rule out the need for fusion (but a positive discogram in itself would not allow fusion). (Carragee-Spine, 2000) (Carragee2-Spine, 2000) (Carragee3-Spine, 2000) (Carragee4-Spine, 2000) (Bigos, 1999) (ACR, 2000) (Resnick, 2002) (Madan, 2002) (Carragee-Spine, 2004) (Carragee2, 2004) (Maghout-Juratli, 2006) (Pneumaticos, 2006) (Airaksinen, 2006) (Manchikanti, 2009) Discography may be supported if the decision has already been made to do a spinal fusion, and a negative discogram could rule out the need for fusion on that disc (but a positive discogram in itself would not justify fusion). Discography may help distinguish asymptomatic discs among morphologically abnormal discs in patients without psychosocial issues. Precise prospective categorization of discographic diagnoses may predict outcomes from treatment, surgical or otherwise. (Derby, 2005) (Derby2, 2005) (Derby, 1999) Positive discography was not highly predictive in identifying outcomes from spinal fusion. A recent study found only a 27% success from spinal fusion in patients with low back pain and a positive single-level low-pressure provocative discogram, versus a 72% success in patients having a well-accepted single-level lumbar pathology of unstable spondylolisthesis. (Carragee, 2006) The prevalence of positive discogram may be increased in subjects with chronic low back pain who have had prior surgery at the level tested for lumbar

disc herniation. (Heggeness, 1997) Invasive diagnostics such as provocative discography have not been proven to be accurate for diagnosing various spinal conditions, and their ability to effectively guide therapeutic choices and improve ultimate patient outcomes is uncertain. (Chou, 2008) Although discography, especially combined with CT scanning, may be more accurate than other radiologic studies in detecting degenerative disc disease, its ability to improve surgical outcomes has yet to be proven. It is routinely used before IDET, yet only occasionally used before spinal fusion. (Cohen, 2005) Provocative discography is not recommended because its diagnostic accuracy remains uncertain, false-positives can occur in persons without low back pain, and its use has not been shown to improve clinical outcomes. (Chou2, 2009) This recent RCT concluded that, compared with discography, injection of a small amount of bupivacaine into the painful disc was a better tool for the diagnosis of discogenic LBP. (Ohtori, 2009) Discography may cause disc degeneration. Even modern discography techniques using small gauge needle and limited pressurization resulted in accelerated disc degeneration (35% in the discography group compared to 14% in the control group), disc herniation, loss of disc height and signal and the development of reactive endplate changes compared to match-controls. These findings are of concern for several reasons. Discography as a diagnostic test is controversial and in view of these findings the utility of this test should be reviewed. Furthermore, discography in current practice will often include injecting discs with a low probability of being symptomatic in an effort to validate other disc injections, a so-called control disc. Although this strategy has never been confirmed to increase test validity or utility, injecting normal discs even with small gauge needles appears to increase the rate of degeneration in these discs over time. The phenomenon of accelerated adjacent segment degeneration adjacent to fusion levels may be, in part, explained by previous disc puncture if discography was used in segments adjacent to the fusion. Similarly, intradiscal therapeutic strategies (injecting steroids, sclerosing agents, growth factors, etc.) have been proposed as a method to treat, arrest or prevent symptomatic disc disease. This study suggests that the injection procedure itself is not completely innocuous and a recalculation of these demonstrated risks versus hypothetical benefits should be considered. (Carragee, 2009) Discography involves the injection of a water-soluble imaging material directly into the nucleus pulposus of the disc. Information is then recorded about the pressure in the disc at the initiation and completion of injection, about the amount of dye accepted, about the configuration and distribution of the dye in the disc, about the quality and intensity of the patient's pain experience and about the pressure at which that pain experience is produced. Both routine x-ray imaging during the injection and post-injection CT examination of the injected discs are usually performed as part of the study. There are two diagnostic objectives: (1) to evaluate radiographically the extent of disc damage on discogram and (2) to characterize the pain response (if any) on disc injection to see if it compares with the typical pain symptoms the patient has been experiencing. Criteria exist to grade the degree of disc degeneration from none (normal disc) to severe. A symptomatic degenerative disc is considered one that disperses injected contrast in an abnormal, degenerative pattern, extending to the outer margins of the annulus and at the same time reproduces the patient's lower back complaints (concordance) at a low injection pressure. Discography is not a sensitive test for radiculopathy and has no role in

its confirmation. It is, rather, a confirmatory test in the workup of axial back pain and its validity is intimately tied to its indications and performance. As stated, it is the end of a diagnostic workup in a patient who has failed all reasonable conservative care and remains highly symptomatic. Its validity is enhanced (and only achieves potential meaningfulness) in the context of an MRI showing both dark discs and bright, normal discs -- both of which need testing as an internal validity measure. And the discogram needs to be performed according to contemporary diagnostic criteria -- namely, a positive response should be low pressure, concordant at equal to or greater than a VAS of 7/10 and demonstrate degenerative changes (dark disc) on MRI and the discogram with negative findings of at least one normal disc on MRI and discogram. See also Functional anesthetic discography (FAD).

**Discography is Not Recommended in ODG.**

**Patient selection criteria for Discography if provider & payor agree to perform anyway:**

- o Back pain of at least 3 months duration
- o Failure of recommended conservative treatment including active physical therapy
- o An MRI demonstrating one or more degenerated discs as well as one or more normal appearing discs to allow for an internal control injection (injection of a normal disc to validate the procedure by a lack of a pain response to that injection)
- o Satisfactory results from detailed psychosocial assessment (discography in subjects with emotional and chronic pain problems has been linked to reports of significant back pain for prolonged periods after injection, and therefore should be avoided)
- o Intended as a screen for surgery, i.e., the surgeon feels that lumbar spine fusion is appropriate but is looking for this to determine if it is not indicated (although discography is not highly predictive) (Carragee, 2006) NOTE: In a situation where the selection criteria and other surgical indications for fusion are conditionally met, discography can be considered in preparation for the surgical procedure. However, all of the qualifying conditions must be met prior to proceeding to discography as discography should be viewed as a non-diagnostic but confirmatory study for selecting operative levels for the proposed surgical procedure. Discography should not be ordered for a patient who does not meet surgical criteria.
- o Briefed on potential risks and benefits from discography and surgery
- o Single level testing (with control) (Colorado, 2001)
- o Due to high rates of positive discogram after surgery for lumbar disc herniation, this should be potential reason for non-certification

The Texas Department of Insurance appointed (Independent Review Organization) as the IRO. (Independent Review Organization) assigned the review to a physician reviewer who was identified as a medical doctor board certified in orthopedic surgery. In upholding Carrier's denial of the discogram, the physician reviewer wrote:

The patient does not meet the ODG criteria for lumbar spine fusion, and therefore the patient does not meet the ODG criteria for discogram. Lumbar instability is not documented in the patient's medical record in flexion/extension views.

Discography is not recommended in ODG. The request does not conform to ODG, and no explanation has been provided as to why the ODG should not be followed in this patient's case. The reviewer finds that medical necessity does not exist for Low Pressure Lumbar Discogram With CT L3-S1.

The physician reviewer certified that he relied upon the Official Disability Guidelines & Treatment Guidelines (ODG) and his medical judgment, clinical experience and expertise in accordance with accepted medical standards.

The ODG entry on lumbar fusion states:

Not recommended for patients who have less than six months of failed recommended conservative care unless there is objectively demonstrated severe structural instability and/or acute or progressive neurologic dysfunction, but recommended as an option for spinal fracture, dislocation, spondylolisthesis or frank neurogenic compromise, subject to the selection criteria outlined in the section below entitled, "Patient Selection Criteria for Lumbar Spinal Fusion," after 6 months of conservative care. For workers' comp populations, see also the heading, "Lumbar fusion in workers' comp patients." After screening for psychosocial variables, outcomes are improved and fusion may be recommended for degenerative disc disease with spinal segment collapse with or without neurologic compromise after 6 months of compliance with recommended conservative therapy. [For spinal instability criteria, see AMA Guides (Andersson, 2000)] For complete references, see separate document with all studies focusing on Fusion (spinal). There is limited scientific evidence about the long-term effectiveness of fusion for degenerative disc disease compared with natural history, placebo, or conservative treatment. Studies conducted in order to compare different surgical techniques have shown success for fusion in carefully selected patients. (Gibson-Cochrane, 2000) (Savolainen, 1998) (Wetzel, 2001) (Molinari, 2001) (Bigos, 1999) (Washington, 1995) (DeBarard-Spine, 2001) (Fritzell-Spine, 2001) (Fritzell-Spine, 2002) (Deyo-NEJM, 2004) (Gibson-Cochrane/Spine, 2005) (Soegaard, 2005) (Glassman, 2006) (Atlas, 2006) According to the recently released AANS/NASS Guidelines, lumbar fusion is recommended as a treatment for carefully selected patients with disabling low back pain due to one- or two-level degenerative disc disease after failure of an appropriate period of conservative care. This recommendation was based on one study that contained numerous flaws, including a lack of standardization of conservative care in the control group. At the time of the 2-year follow up it appeared that pain had significantly increased in the surgical group from year 1 to 2. Follow-up post study is still pending publication. In addition, there remains no direction regarding how to define the "carefully selected patient." (Resnick, 2005) (Fritzell, 2004) A recently published well respected international guideline, the "European Guidelines," concluded that fusion surgery for nonspecific chronic LBP cannot be recommended unless 2 years of all other recommended conservative treatments – including multidisciplinary approaches with combined programs of cognitive intervention and exercises – have failed, or such combined programs are not available, and only then in carefully selected patients with maximum 2-level degenerative disc disease. (Airaksinen, 2006) For chronic LBP, exercise and cognitive intervention may be equivalent to lumbar fusion without

the potentially high surgical complication rates. (Ivar Brox-Spine, 2003) (Keller-Spine, 2004) (Fairbank-BMJ, 2005) (Brox, 2006) In acute spinal cord injury (SCI), if the spine is unstable following injury, surgical fusion and bracing may be necessary. (Bagnall-Cochrane, 2004) (Siebenga, 2006) A study on improving quality through identifying inappropriate care found that use of guideline-based Utilization Review (UR) protocols resulted in a denial rate for lumbar fusion 59 times as high as denial rates using non-guideline based UR. (Wickizer, 2004) The profit motive and market medicine have had a significant impact on clinical practice and research in the field of spine surgery. (Weiner-Spine, 2004) (Shah-Spine, 2005) (Abelson, 2006) Data on geographic variations in medical procedure rates suggest that there is significant variability in spine fusion rates, which may be interpreted to suggest a poor professional consensus on the appropriate indications for performing spinal fusion. (Deyo-Spine, 2005) (Weinstein, 2006) Outcomes from complicated surgical fusion techniques (with internal fixation) may be no better than the traditional posterolateral fusion. (van Tulder, 2006) (Maghout-Juratli, 2006) Despite the new technologies, reoperation rates after lumbar fusion have become higher. (Martin, 2007) According to the recent Medicare Coverage Advisory Committee Technology Assessment, the evidence for lumbar spinal fusion does not conclusively demonstrate short-term or long-term benefits compared with nonsurgical treatment for elderly patients. (CMS, 2006) When lumbar fusion surgery is performed, either with lateral fusion alone or with interbody fusion, unlike cervical fusion, there is no absolute contraindication to patients returning even to contact sports after complete recovery from surgery. Like patients with a thoracic injury, those with a lumbar injury should be pain free, have no disabling neurological deficit, and exhibit evidence of bone fusion on x-ray films before returning. (Burnett, 2006) A recent randomized controlled trial comparing decompression with decompression and instrumented fusion in patients with foraminal stenosis and single-level degenerative disease found that patients universally improved with surgery, and this improvement was maintained at 5 years. However, no obvious additional benefit was noted by combining decompression with an instrumented fusion. (Hallett, 2007) Discography may be supported if the decision has already been made to do a spinal fusion, and a negative discogram could rule out the need for fusion on that disc (but a positive discogram in itself would not justify fusion). Discography may help distinguish asymptomatic discs among morphologically abnormal discs in patients without psychosocial issues. Precise prospective categorization of discographic diagnoses may predict outcomes from treatment, surgical or otherwise. (Derby, 2005) (Derby2, 2005) (Derby, 1999) New research shows that healthcare expenditures for back and neck problems have increased substantially over time, but with little improvement in healthcare outcomes such as functional disability and work limitations. Rates of imaging, injections, opiate use, and spinal surgery have increased substantially over the past decade, but it is unclear what impact, if any, this has had on health outcomes. (Martin, 2008) The efficacy of surgery for nonspecific back pain is uncertain. There may be some patients for whom surgery, fusion specifically, might be helpful, but it is important for doctors to discuss the fact that surgery doesn't tend to lead to huge improvements on average, about a 10- to 20-point improvement in function on a 100-point scale, and a significant proportion of patients still need to take pain medication and don't return to full function. (Chou, 2008) This study showed that

fusion for chronic lower back pain was the least successful common orthopaedic surgery. The study compared the gains in quality of life achieved by total hip replacement, total knee replacement, surgery for spinal stenosis, disc excision for lumbar disc herniation, and arthrodesis for chronic low back pain. For chronic lower back pain, improvements were statistically significant but clinically negligible. Although pain was reduced and function improved slightly, outcomes remained in the moderately affected range, quality of life was not improved and rendered worse, on average. While surgery for spinal stenosis and for disc herniation compare well with archetypical orthopaedic operations, the outcomes of surgery for chronic lower back pain do not even approach those of other orthopaedic procedures, and the data show that patients with back pain are rendered worse off by surgery. (Hansson, 2008) Recent studies document a 220% increase in lumbar spinal fusion surgery rates, without demonstrated improvements in patient outcomes or disability rates. (Deyo, 2009) In a study of 2,378 Washington State workers' compensation claimants who underwent fusion to assess the frequency, timing, and causes of death, the 3-year cumulative mortality rate post-fusion was 1.93% and analgesic-related deaths were responsible for 21% of all deaths and 31.4% of all potential life lost. (Juratli, 2009) A study to compare the surgical experience, clinical outcomes, and effect on body weight between obese and morbidly obese patients undergoing lumbar spine fusion surgery concluded that clinical outcomes were independent of the BMI of the patient, but the incidence of postoperative complications was significant in 45% of morbidly obese and 44% of obese patients. The authors proposed that morbidly obese patients should undergo bariatric surgery before spine surgery in nonemergent situations. (Vaidya, 2009) For nonradicular low back pain with common degenerative changes, there is fair evidence that fusion is no better than intensive rehabilitation with a cognitive-behavioral emphasis for improvement in pain or function, and less than half of patients experience optimal outcomes (defined as no more than sporadic pain, slight restriction of function, and occasional analgesics) following fusion. (Chou, 2009) Posterolateral bone-grafting fusion is not necessary when a Denis type-B thoracolumbar burst fracture associated with a load-sharing score of  $\leq 6$  is treated with short-segment pedicle screw fixation. (Dai, 2009) Discography (and not merely the fusion) may actually be the cause of adjacent segment disc degeneration. This study suggested that the phenomenon of accelerated adjacent segment degeneration adjacent to fusion levels may be, in part, explained by previous disc puncture if discography was used in segments adjacent to the fusion. (Carragee, 2009) Among Medicare recipients, the frequency of complex fusion procedures for spinal stenosis increased 15-fold in just 6 years. The introduction and marketing of new surgical devices and financial incentives may stimulate more invasive surgery. (Deyo-JAMA, 2010) Lumbar spinal fusion surgeries use bone grafts, and are sometimes combined with metal devices, to produce a rigid connection between two or more adjacent vertebrae. The therapeutic objective of spinal fusion surgery for patients with low back problems is to prevent any movement in the intervertebral spaces between the fused vertebrae, thereby reducing pain and any neurological deficits. See also Adjacent segment disease/degeneration (fusion) & Iliac crest donor-site pain treatment.

Lumbar fusion in workers' comp patients: In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may

affect overall success of the procedure, which should be considered. Until further research is conducted there remains insufficient evidence to recommend fusion for chronic low back pain in the absence of stenosis and spondylolisthesis, and this treatment for this condition remains “under study.” It appears that workers’ compensation populations require particular scrutiny when being considered for fusion for chronic low back pain, as there is evidence of poorer outcomes in subgroups of patients who were receiving compensation or involved in litigation. (Fritzell-Spine, 2001) (Harris-JAMA, 2005) (Maghout-Juratli, 2006) (Atlas, 2006) Despite poorer outcomes in workers’ compensation patients, utilization is much higher in this population than in group health. (Texas, 2001) (NCCI, 2006) Presurgical biopsychosocial variables predict patient outcomes from lumbar fusion, which may help improve patient selection. Workers’ compensation status, smoking, depression, and litigation were the most consistent presurgical predictors of poorer patient outcomes. Other predictors of poor results were number of prior low back operations, low household income, and older age. (DeBerard-Spine, 2001) (DeBerard, 2003) (Deyo, 2005) (LaCaille, 2005) (Trief-Spine, 2006) Obesity and litigation in workers’ compensation cases predict high costs associated with interbody cage lumbar fusion. (LaCaille, 2007) A recent study of 725 workers’ comp patients in Ohio who had lumbar fusion found only 6% were able to go back to work a year later, 27% needed another operation, and over 90% were in enough pain that they were still taking narcotics at follow-up. (Nguyen, 2007) A recent case-control study of lumbar fusion outcomes in worker’s compensation (WC) patients concluded that only 9% of patients receiving WC achieved substantial clinical benefit compared to 33% of those not receiving WC. (Carreon, 2009)

Lumbar fusion for spondylolisthesis: Recommended as an option for spondylolisthesis. Patients with increased instability of the spine after surgical decompression at the level of degenerative spondylolisthesis are candidates for fusion. (Eckman, 2005) This study found only a 27% success from spinal fusion in patients with low back pain and a positive single-level low-pressure provocative discogram, versus a 72% success in patients having a well-accepted single-level lumbar pathology of unstable spondylolisthesis. (Carragee, 2006) Unilateral instrumentation used for the treatment of degenerative lumbar spondylolisthesis is as effective as bilateral instrumentation. (Fernandez-Fairen, 2007) Patients with degenerative spondylolisthesis and spinal stenosis who undergo standard decompressive laminectomy (with or without fusion) showed substantially greater improvement in pain and function during a period of 2 years than patients treated nonsurgically, according to the recent results from the Spine Patient Outcomes Research Trial (SPORT). (Weinstein-spondylolisthesis, 2007) (Deyo-NEJM, 2007) For degenerative lumbar spondylolisthesis, spinal fusion may lead to a better clinical outcome than decompression alone. No conclusion about the clinical benefit of instrumenting a spinal fusion can be made, but there is moderate evidence that the use of instrumentation improves the chance of achieving solid fusion. (Martin, 2007) A recent systematic review of randomized trials comparing lumbar fusion surgery to nonsurgical treatment of chronic back pain associated with lumbar disc degeneration, concluded that surgery may be more efficacious than unstructured nonsurgical care but may not be more efficacious than structured cognitive-behavior therapy. Methodological limitations of the randomized trials prevented firm conclusions. (Mirza, 2007) A

comparison of surgical and nonoperative outcomes between degenerative spondylolisthesis and spinal stenosis patients from the SPORT trial found that fusion was most appropriate for spondylolisthesis, with or without listhesis, and decompressive laminectomy alone most appropriate for spinal stenosis. (Pearson, 2010)

Lumbar fusion for Scheuermann's kyphosis: Recommended as an option for adult patients with severe deformities (e.g. more than 70 degrees for thoracic kyphosis), neurological symptoms exist, and pain cannot be adequately resolved non-operatively (e.g. physical therapy, back exercises). Good outcomes have been found in a relatively large series of patients undergoing either combined anterior-posterior or posterior only fusion for Scheuermann's kyphosis. (Lonner, 2007)

#### **Patient Selection Criteria for Lumbar Spinal Fusion:**

For chronic low back problems, fusion should not be considered within the first 6 months of symptoms, except for fracture, dislocation or progressive neurologic loss. Indications for spinal fusion may include: (1) Neural Arch Defect - Spondylolytic spondylolisthesis, congenital neural arch hypoplasia. (2) Segmental Instability (objectively demonstrable) - Excessive motion, as in degenerative spondylolisthesis, surgically induced segmental instability and mechanical intervertebral collapse of the motion segment and advanced degenerative changes after surgical discectomy. [For excessive motion criteria, see AMA Guides, 5th Edition, page 384 (relative angular motion greater than 20 degrees). (Andersson, 2000) (Luers, 2007)] (3) Primary Mechanical Back Pain (i.e., pain aggravated by physical activity)/Functional Spinal Unit Failure/Instability, including one or two level segmental failure with progressive degenerative changes, loss of height, disc loading capability. In cases of workers' compensation, patient outcomes related to fusion may have other confounding variables that may affect overall success of the procedure, which should be considered. There is a lack of support for fusion for mechanical low back pain for subjects with failure to participate effectively in active rehab pre-op, total disability over 6 months, active psych diagnosis, and narcotic dependence. [For spinal instability criteria, see AMA Guides, 5th Edition, page 379 (lumbar inter-segmental movement of more than 4.5 mm). (Andersson, 2000)] (4) Revision Surgery for failed previous operation(s) if significant functional gains are anticipated. Revision surgery for purposes of pain relief must be approached with extreme caution due to the less than 50% success rate reported in medical literature. (5) Infection, Tumor, or Deformity of the lumbosacral spine that cause intractable pain, neurological deficit and/or functional disability. (6) After failure of two discectomies on the same disc, fusion may be an option at the time of the third discectomy, which should also meet the ODG criteria. (See ODG Indications for Surgery -- Discectomy.)

**Pre-operative Surgical Indications Recommended:** Pre-operative clinical surgical indications for spinal fusion should include all of the following: (1) All pain generators are identified and treated; & (2) All physical medicine and manual therapy interventions are completed (sic); & (3) X-rays demonstrating spinal instability and/or myelogram, CT-myelogram, or discography (see discography criteria) & MRI demonstrating disc pathology; & (4) Spine pathology limited to two levels; & (5) Psychosocial screen with confounding issues addressed. (6) For any potential fusion surgery, it is recommended that the injured worker refrain from smoking for at least six weeks prior to surgery and during the period of fusion healing. (Colorado, 2001) (BlueCross BlueShield, 2002)

In accordance with Division Rule 133.308 (t), "A decision issued by an IRO is not considered an agency decision and neither the Department nor the Division are considered parties to an appeal. In a Contested Case Hearing (CCH), the party appealing the IRO decision has the burden of overcoming the decision issued by an IRO by a preponderance of evidence-based medical evidence."

Dr. B disagrees that Claimant does not meet the ODG criteria for a lumbar discogram, contending that it is because he intends it as a confirmatory test – if it is negative it would rule out the need for a fusion. He also disagrees that spinal fusion is not indicated for Claimant. Dr. B testified that Claimant has mechanical low back pain that has not responded to conservative treatment and, therefore, is a candidate for fusion. There is, however, no evidence that Claimant has segmental failure with progressive degenerative changes, loss of height, and disc loading capability. Nor does Claimant's condition meet any other segment of the patient selection criteria for fusion.

In determining the weight to be given to the opinion of an expert, a trier of fact must first determine if the expert is qualified to offer it. The trier of fact must then, however, determine whether the opinion is relevant to the issues at bar and whether it is based upon a solid foundation. An expert's bald assurance of validity is not enough. *See Black vs. Food Lion, Inc.*, 171 F.3rd 308 (5th Cir. 1999); *E.I. Du Pont De Nemours and Company, Inc. v. Robinson*, 923 S.W.2d 549 (Tex. 1995). Evidence is considered in terms of (1) general acceptance of the theory and technique by the relevant scientific community; (2) the expert's qualifications; (3) the existence of literature supporting or rejecting the theory; (4) the technique's potential rate of error; (5) the availability of other experts to test and evaluate the technique; and (7) the experience and skill of the person who applied the technique on the occasion in question. *Kelly v. State*, 792 S.W.2d 579 (Tex.App.-Fort Worth 1990). A medical doctor is not automatically qualified as an expert on every medical question and an unsupported opinion has little, if any, weight. *Black v. Food Lion, Inc.*, 171 F.3rd 308 (5th Cir. 1999).

In weighing the evidence presented in this matter, the hearing officer finds that while there may be some support for Dr. B's assertions and opinions, the preponderance of the evidence supports the IRO physician reviewer's conclusion that a lumbar discogram is not medically necessary in this instance. There has been no agreement between the parties to do a discogram nor is there a pending lumbar fusion. A positive discogram would not support a request for fusion as currently contemplated by Dr. B and the evidence presented fails to show that Claimant meets the patient selection criteria for a lumbar fusion as set forth in the ODG. Dr. B testified that a negative discogram would result in a request for preauthorization of a lumbar fusion and it would appear that the IRO physician reviewer is correct when he says that the medical records do not support the need for lumbar fusion. Although there is evidence that some studies support discography, the ODG explicitly recommends against the procedure. The preponderance of the evidence is not contrary to the IRO recommendation.

Even though all the evidence presented was not discussed, it was considered. The Findings of Fact and Conclusions of Law are based on all of the evidence presented.

### **FINDINGS OF FACT**

1. The parties stipulated to the following facts:

- A. Venue is proper in the (City) Field Office of the Texas Department of Insurance, Division of Workers' Compensation.
  - B. On \_\_\_\_\_, Claimant sustained a compensable injury while the employee of (Employer).
  - C. The Texas Department of Insurance appointed (Independent Review Organization) as the IRO in this matter to review Carrier's denial of the requested low pressure discogram with CT at L3-4, L4-5, and L5-S1.
  - D. The IRO upheld Carrier denial of the requested procedure.
2. Carrier delivered to Claimant a single document stating the true corporate name of Carrier, and the name and street address of Carrier's registered agent, which document was admitted into evidence as Hearing Officer's Exhibit Number 2.
  3. Discography is not recommended in the ODG.
  4. The parties have not agreed to perform a discogram despite the negative recommendation in the ODG.
  5. Claimant has not requested preauthorization of a lumbar fusion in this matter and the discogram requested is not intended to exclude a disc level from a planned spinal fusion surgery.
  6. The discography requested is intended to act as a pre-operative screening tool and its use as such is not recommended in the ODG.
  7. Low pressure discography with CT at L3-4, L4-5 and L5-S1 is not reasonably required medical treatment for the compensable injury of \_\_\_\_\_.

### **CONCLUSIONS OF LAW**

1. The Texas Department of Insurance, Division of Workers' Compensation, has jurisdiction to hear this case.
2. Venue is proper in the (City) Field Office.
3. The preponderance of the evidence is not contrary to the decision of IRO that low pressure discography with CT at L3-4, L4-5 and L5-S1 is not reasonably required medical care for the compensable injury of \_\_\_\_\_.

### **DECISION**

Claimant is not entitled to low pressure discography with CT at L3-4, L4-5 and L5-S1 as reasonably required medical care for the compensable injury of \_\_\_\_\_.

**ORDER**

Carrier is not liable for the benefits at issue in this hearing. Claimant remains entitled to medical benefits for the compensable injury in accordance with §408.021.

The true corporate name of the insurance carrier is **COMMERCE & INDUSTRY INSURANCE COMPANY** and the name and address of its registered agent for service of process is

**CORPORATION SERVICE CO.  
701 BRAZOS STREET, SUITE 1050  
AUSTIN, TX 78701-3232**

Signed this 20th day of May, 2010.

KENNETH A. HUCHTON  
Hearing Officer